

*HIMALAYAN AND  
KASHMIRI BIRDS  
BEING A KEY TO THE BIRDS  
COMMONLY SEEN IN SUMMER  
IN THE HIMALAYAS & KASHMIR  
BY DOUGLAS DEWAR*

---

---

*A COMPANION TO THE  
BIRD VOLUMES OF THE  
FAUNA OF BRITISH INDIA*

---

---

LONDON  
JOHN LANE THE BODLEY HEAD LIMITED

*First Published 1923*

Made and Printed in Great Britain at  
*The Mayflower Press, Plymouth.* William Brendon & Son, Ltd.

---

---

## *PREFACE*

---

THE object of this book is to enable people interested in birds to identify those they meet with while walking in the hill stations of the Himalayas and Kashmir and those they see in the Kashmir valley.

The birds dealt with are the ones commonly seen IN SUMMER at such places. As nine out of ten species of hill birds move to lower levels in winter, the bird population of a hill station in winter differs from that in summer.

Had the scope of this book included birds seen in winter at the various hill stations, and those found in summer only at lower elevations than 5000 feet above the sea-level or higher altitudes than 7500, its bulk would have been considerably increased. It would have attained even greater dimensions had I noticed the rare birds that are sometimes seen in hill stations in summer.

Paradoxical though it may sound, the value of this book lies largely in its omissions !

---

---

## *Preface*

---

If this key had included all the birds of Kashmir and the Himalayas, it would have bewildered the user by its complexity, and so failed in its object. As it stands, it should enable anyone who consults it to identify readily twenty-nine-thirtieths of the birds he sees in summer at elevations between 5000 and 7500 feet.

Except in so far as they may serve as aids to identification, nothing has been said about the habits of birds in this book, which is merely an artificial key.

A treatise on the life-histories and habits of the birds of Kashmir and the Himalayas has yet to be written. Much information, however, about the habits of these birds is contained in Hume's *Nests and Eggs of Indian Birds*, Jerdon's *Birds of India*, the bird volumes of the *Fauna of British India* and my own little book entitled *Birds of the Indian Hills*.

---

---

# CONTENTS

---

	PAGE
1. HOW TO USE THIS BOOK . . . . .	II
2. LOCAL LISTS OF THE BIRDS OF THE HIMALAYAS AND KASHMIR WHICH HAVE APPEARED IN :	
(a) "The Ibis" . . . . .	13
(b) "The Journal of the Asiatic Society" . . . . .	14
(c) "The Journal of the Bombay Natural History Society" . . . . .	15

## PART I

A. CLASSIFICATION ACCORDING TO STRUCTURAL PECULIARITIES . . . . .	19
I. Bill . . . . .	19
II. Crest . . . . .	20
III. Tail . . . . .	22
IV. Sexual Dimorphism . . . . .	25
B. CLASSIFICATION ACCORDING TO COLOUR . . . . .	27
I. Black . . . . .	28
II. Blue . . . . .	42
III. Brown . . . . .	45

---

---

## *Contents*

---

	PAGE
IV. Fawn Colour . . . . .	54
V. Green . . . . .	55
VI. Grey . . . . .	56
VII. Red . . . . .	59
VIII. White . . . . .	67
IX. Yellow . . . . .	71
X. Birds of Many Colours . . . . .	75
C. CLASSIFICATION ACCORDING TO HABITS . . . . .	76
I. Birds having remarkable cries . . . . .	76
II. Birds that commonly visit gardens . . . . .	81
III. Birds that associate in flocks . . . . .	83
IV. Birds that dwell by streams . . . . .	84
V. Birds that occur abundantly on, over, or on the margins of the lakes of the Kashmir valley . . . . .	84
VI. Birds that move about on the trunks of trees in search of insects . . . . .	86
VII. Birds that find much of their food on the ground . . . . .	86
VIII. Birds that capture insects on the wing . . . . .	88
IX. Birds that spend great part of the day on the wing . . . . .	88

## PART II

DESCRIPTIVE LIST OF BIRDS . . . . .	91
-------------------------------------	----

---

---

## HOW TO USE THIS BOOK

---

THIS key has been compiled on the principle followed in *Indian Birds*. Even as the latter has been the means of teaching scores of people the names of the birds commonly seen in the plains of India, so will this, I venture to hope, enable anyone who wishes to do so, to learn in a very short time the names of all the birds seen commonly in summer in Kashmir and at the various Himalayan hill stations.

The student should make a note regarding any strange bird seen, giving the colours in its plumage and any anatomical peculiarity, such as a crest, a long tail or bill. On returning to the house, Part I of the book should be consulted. The bird in question should find place in two or more of the classes of birds in that Part, and so may be set down as being probably one of two or three species. A reference to the descriptions in Part II should complete the identification.

To take an easy example. The observer sees

---

---

## *How to Use This Book*

---

a striking bird. He notes that its body is about the size of that of the myna, that it has a long tail and that there are blue, black, brown and white in the plumage, and the bill is bright red. He consults in Part I of the book, "Birds having the tail long in proportion to the body." He finds there are fifteen of these birds. He then refers to the class "Birds of which the plumage is black, blue, brown and white," and finds that there are only three of these, of which only two have the tail long. He is thus satisfied that the unknown bird is a blue magpie. The fact that the bill is red leads him to infer that it is the red-billed blue magpie. A reference to the description of this species in Part II confirms the diagnosis.

I may add that the descriptions have designedly been made as brief as possible. Too much detail might easily hinder identification, as in that case the descriptions would often include inconspicuous features which the observer had failed to notice.

A full description of every bird found in India is given in the *Fauna of British India*, to which this book is really a companion and a guide for the novice.



**LOCAL LISTS OF THE BIRDS OF THE HIMALAYAS AND KASHMIR WHICH HAVE APPEARED IN "THE IBIS"**

Year of Ibis	Name of Author	Name of Locality
1861.	L. H. Irby	. Oudh and Kumaun.
1868.	R. C. Tytler	. Birds observed on a march from Simla to Mussoorie.
1868.	A. von Pelzeln	. Birds collected by Dr. Stoliczka in Thibet and the Himalayas.
1869.	W. E. Brooks	. Birds observed near Nynee Tal and Almora.
1869.	G. E. Bulger	. Birds obtained in Sikkim, Eastern Himalayas.
1884.	C. H. T. Marshall.	Birds of Chamba in the N.W. Himalayas.
1888.	W. W. Cordeaux	. Birds of Cashmere and the Dras District.
1898.	J. Davidson	. A short trip to Kashmir.

LOCAL LISTS OF THE BIRDS OF THE HIMALAYAS AND KASHMIR WHICH HAVE APPEARED IN THE " JOURNAL OF THE ASIATIC SOCIETY "

Year of Publication	Name of Author	Name of Locality
1868.	F. Stoliczka	. Ornithological Observations in the Sutlej Valley, N.W. Himalayas.
1874.	W. E. Brooks	. Notes on the Ornithology of Cashmir.
1876. } 1878. }	H. H. Godwin- Austen	Lists of Birds from the Hill Ranges of the N.E. Frontier of India.
1900.	H. J. Walton	. Notes on Birds collected in Kumaun.

**LOCAL LISTS OF BIRDS THAT HAVE AP-  
PEARED IN THE "JOURNAL OF THE  
BOMBAY NATURAL HISTORY SOCIETY"**

No. of Vol.	Name of Compiler	Locality
X.	R. H. Rattray	. Notes on nests taken from March to June at Kohat and Mussoorie, N.W.P.
XII.	N. F. T. Wilson	. Nesting in Kashmir.
XVI.	R. H. Rattray	. Birds nesting in the Murree Hills and Gal- lies.
XVII. & XVIII.)	G. E. Ward	. Birds of the Province of Kashmir and Jamu and adjacent districts.
XVIII.		
XIX.	H.A.F. Magrath	. Bird Notes from Murree and the Gallies.
XXI.	H.A.F. Magrath	. Bird Notes by the Way in Kashmir.
XXIV.	H. Whistler	. Some Birds observed at Dalhousie Hill Station in 1915.
XXVI.	A. E. Jones	. A List of Birds found in the Simla Hills.

---

---

## *Local Lists of Birds*

---

No. of Vol.	Name of Compiler	Locality
XXVI.	H. Whistler	. Some Birds observed at Fagoo, near Simla.
XXVII.	R.W.G. Hingston	A List of the Birds of Dharmsala.
XXVII.	H. Whistler	. Further Notes on Birds about Simla.
XXVIII.	A. E. Osmaston	Note on the Nidification and Habits of Some Birds in British Garhwal.

---

---

# *PART I*

## *CLASSIFICATION*

- A. Classification according to Structural Peculiarities*
  - B. Classification according to Colour*
  - C. Classification according to Habits*
-



---

---

*A. CLASSIFICATION ACCORDING TO  
STRUCTURAL PECULIARITIES*

---

- I. Bill.
- II. Crest.
- III. Tail.
- IV. Sexual Dimorphism.

I. BILL

*Birds with Long Bills*

N.B.—The numbers in brackets are those given to the birds in the descriptive list. The capital letters against the bird indicate where it occurs: K=Kashmir. W=Western Himalayas. E=Eastern Himalayas.

1. The Rusty-cheeked Scimitar Babbler (25). E. W. (Fairly long, with a slight downward curve.)
2. The Sunbirds (111 and 112). E. W. (Very small birds with long, slender, downwardly-curved bills.)

---

---

## *Himalayan and Kashmiri Birds*

---

3. The Kingfishers (127 and 128), E. W. K. (Fairly long and straight.)
4. The European Hoopoe (129). E. W. K. (Bill very long and slender, with a slight downward curve.)
5. The Herons (174-176). K. (Fairly long and dagger-shaped.)

### II. CREST

#### *Crested Birds*

1. The Black-throated Jay (7). W. K.
2. The Himalayan Jay (8). E. W. K.
3. The Yellow-cheeked Tit (14). W.
4. The Crested Black Tit (15). W. K.
5. The Brown Crested Tit (16). W. K.
6. The Black-headed Sibia (30). E. W.
7. The Yellow-naped Ixulus (32). E. W.
8. The Himalayan Black Bulbul (37). K.  
E. W.
9. The Striated Green Bulbul (38). E.
10. The Bengal Red-vented Bulbul (39). E.
11. The Punjab Red-vented Bulbul (40). W.
12. The White-cheeked Bulbul (41). E. W.  
K.
13. The Indian Paradise Flycatcher (73).  
W. K.



---

---

*Classification : Structural Peculiarities*

---

14. The West-Himalayan Scaly-bellied Green Woodpecker (114). W. K.

15. The Black-naped Green Woodpecker (11). W. K. (Crest small.)

16. The Northern Pale-headed Woodpecker (116). E. (Crest small.)

17. The Himalayan Pied Kingfisher (127). E. W. K. (Crest small.)

18. The European Hoopoe (129). E. W. K.

19. The Pied Crested Cuckoo (137). W. K.

20. Hodgson's Hawk-Eagle (151). E. W. K. (Crest small.)

21. The Crested Serpent-Eagle (152). E. W.

22. The White-crested Kalij Pheasant (162). W.

23. The Black-backed Kalij Pheasant (163). E.

24. The Large Cormorant (172). K. (Crest small.)

25. The Common Heron (174). K. (Crest recumbent.)

26. The Night Heron (175). K. (Crest recumbent.)

27. The Little Bittern (176). K. (Crest short.)

---

---

# *Himalayan and Kashmiri Birds*

---

## III. TAIL

### 1. *Birds having the tail markedly forked*

1. The Black Drongo (43). W. E.
2. The Indian Ashy Drongo (44). E. W. K.
3. The Western Spotted Forktail (77). W. K.
4. The Eastern Spotted Forktail (78). E.
5. The Swallow (100). E. W. K.
6. The Wire-tailed Swallow (101). W. K.
7. Syke's Striated Swallow (102). E. W.
8. The Common Pariah Kite (154). E. W. K. (The tail is only slightly forked, but this is important for purposes of identification, as all other birds of prey have rounded or wedge-shaped tails.)

### 2. *Birds having the tail long in proportion to the body*

1. The Red-billed Blue Magpie (4). W. (Tail very long.)
2. The Yellow-billed Blue Magpie (5). E. W. K. (Tail very long.)

---

---

*Classification : Structural Peculiarities*

---

3. The Himalayan Tree-pie (6). E. W.
4. The Himalayan Black Bulbul (37). E. W. K. (Tail fairly long.)
5. The Black Drongo (43). W. E.
6. The Indian Ashy Drongo (44). E. W. K.
7. The Indian Paradise Flycatcher (73). W. K. (Tail very long in cock only.)
8. The Western Spotted Forktail (77). W. K.
9. The Eastern Spotted Forktail (78). E.
10. The Wire-tailed Swallow (101). W. K. (The two outer-tail feathers are prolonged far beyond the others and look like wires. They sometimes get broken off.)
11. The European Bee-eater (126). K. (The middle pair of tail feathers are prolonged beyond the others as bristles.)
12. The Cuckoos (132, 133, 135, 136, 137, 138). (Have rather long tails.)
13. The Paroquets (139, 140, 141).
14. Some of the Pheasants (*not* the Kalij or the Monal).
15. The Pheasant-tailed Jaçana (169). K.

---

---

*Himalayan and Kashmiri Birds*

---

3. *Birds having very short tails in proportion to the body*

1. The White-tailed Nuthatch (42). E. W.
2. The Kashmir Wren (47). W. K.
3. The Scaly-breasted Wren (48). E. W.
4. The Brown Dipper (88). E. W. K.
5. The Woodpeckers (114-121). (Fairly short.)
6. The Golden-throated Barbet (124). (Fairly short.)
7. The Kingfishers (127 and 128).
8. The Owls (142-145).
9. The Common Grey Quail (164). E. W.
10. The Chukor (165). W. K.
11. The Black Partridge (166). W. K.
12. The Moorhen (167). K.
13. The Coot (168). E. W. K.
14. The Red-wattled Lapwing (170). E. W. K.
15. The Cormorants (172 and 173). K.
16. The Night Heron (175). K.
17. The Little Bittern (176). K.
18. The Mallard (177). K.
19. The White-eyed Pochard (178). K.
20. The Indian Little Grebe (179). K. (No tail at all.)

#### IV. SEXUAL DIMORPHISM

*Birds in which the sexes differ greatly in appearance*

1. The Fire-cap (35). W.
2. The Spotted-wing (36). W.
3. The Short-billed Minivet (57). E. W.
4. The Indian Oriole (59). W. K.
5. The White-browed Blue Flycatcher (66).  
E. W.
6. The Rufous-bellied Niltava (71). E. W.  
K.
7. The Large Niltava (72). E.
8. The Indian Paradise Flycatcher (73).  
W. K.
9. The Common Pied Bush Chat (74).  
W. K.
10. The Indian Bush Chat (75). W. K.
11. The Dark-grey Bush Chat (76). E. W.  
K.
12. The Blue-fronted Redstart (81). E. W.  
K.
13. The Plumbeous Redstart (82). E. W. K.
14. The Red-flanked Bush-Robin (83). E.  
W. K.

---

---

## *Himalayan and Kashmiri Birds*

---

15. The Magpie-Robin (84). W.
16. The Grey-winged Ouzel (85). E. W.
17. Tickell's Ouzel (86). K.
18. The Blue-headed Rock-Thrush (87). E.  
W. K.
19. The Black-and-yellow Grosbeak (89).  
W. K.
20. The Scarlet Finch (90). E.
21. The Red-mantled Rose-Finch (91).  
W. K.
22. The Red-browed Finch (93). W. K.
23. The House Sparrow (95). W. K.
24. The Cinnamon Tree-Sparrow (97).  
W. K.
25. The White-capped Bunting (98). W. K.
26. The Sunbirds (111 and 112).
27. The Fire-breasted Flower-pecker (113).  
E. W.
28. The Woodpeckers (114-121). (Difference not very great ; the cock has on the head red which the hen lacks.)
29. The Indian Koel (138). W. K.
30. The Western Blossom-headed Paroquet (140). W. K.
31. The Pheasants (162-163).
32. The Black Partridge (166). W. K.
33. The Mallard (177). K.

---

---

*B. CLASSIFICATION ACCORDING TO  
COLOUR*

---

- I. Black.
- II. Blue.
- III. Brown.
- IV. Fawn colour.
- V. Green.
- VI. Grey.
- VII. Red.
- VIII. White.
- IX. Yellow.
- X. Birds of many colours.

Colour affords the easiest means of identifying the great majority of birds, but in many cases the colours displayed are not of a nature to admit of strict classification. There are a great many shades of all the primary colours, and to attempt to distinguish in the lists the various shades of each colour would be confusing to the reader ; in most cases therefore I have not attempted to do so. Many birds have colouring which some people would call

---

---

## *Himalayan and Kashmiri Birds*

---

brown and others grey. In doubtful cases I have included such colouring in both the grey and the brown lists.

In the case of many birds, such as owls and birds of prey, the various species have generally similar colouring.

### I. BLACK

1. Birds of which the plumage is entirely black or appears to be so from a distance.
2. Birds of which the plumage is mainly black.
3. Black-headed birds.
4. Black collaret or gorget or band across the breast.
5. Black band running longitudinally along abdomen.
6. Black streak through eye.
7. Birds having a black moustachial streak.
8. Black and blue birds.
9. Black and brown birds.
10. Black and grey birds.
11. Black and red birds.
12. Black and white birds.
13. Black and yellow birds.
14. Black, blue and white birds.
15. Black, brown and white birds.



---

---

## *Classification according to Colour*

---

16. Black, fawn-coloured and white birds.
17. Black, grey, and white birds.
18. Black, blue, white and chestnut birds.
19. Black, blue, white and brown birds.
20. Black, blue, white and crimson birds.
21. Black, blue, white and red birds.
22. Black, grey, white and chestnut birds.

I. *Birds of which the plumage is entirely black or appears to be so from a distance*

1. The Jungle Crow (1). E. W. K.
2. The Himalayan Whistling Thrush (29). E. W. K. (A good deal of cobalt blue in the plumage which is only noticeable in the sunlight.)
3. The Himalayan Black Bulbul (37). E. W. K. (There is much grey in the plumage, and the bill is red. I have included this among the black birds, because seen from a little distance it looks black.)
4. The Black Drongo (43). W. E.
5. The Indian Ashy Drongo (44). E. W. K. (The plumage of this bird is dark indigo above and slaty below, but I have included the species among black birds because, seen from a distance, it looks black.)
6. The Grey-winged Ouzel (85). E. W.

---

## *Himalayan and Kashmiri Birds*

---

(Cock only. There is some silver-grey in the wings, but the bird looks black from the distance. Bill coral red.)

7. The Indian Koel (138). W. K. (Cock only. Eyes red, bill green.)

8. The Large White Scavenger Vulture (149). W. K. (Young bird only.)

9. The Coot (168). E. W. K. (White bill and shield on forehead.)

10. The Indian Shag (173). K.

### *2. Birds of which the plumage is mainly black*

1. The Indian House Crow (2). E. W. K. (Grey neck.)

2. The Jackdaw (3). K. (Grey neck.)

3. The Himalayan Whistling Thrush (29). E. W. K. (Patches of cobalt blue.)

4. The Himalayan Starling (60). K. (Plumage glossed with purple, red, and green, with many yellow spots.)

5. The Common Pied Bush Chat (74). W. K. (Cock only. White wing-bar and patch on lower back.)

6. The White-capped Redstart (80). E. W. K. (White cap ; tail, rump, and abdomen chestnut.)

---

---

*Classification according to Colour*

---

7. The Magpie-robin (84). Cock only. (Abdomen, outer tail feathers, and wing-bar white.)

8. The Grey-winged Ouzel (85). E. W. (Cock only. Silvery-grey on wing.)

9. The Common Indian Swift (130). W. K. (Plumage brownish-black, with white bar across the lower back.)

10. The White-necked Spine-tail (131). E. W. (Plumage blackish-brown with middle back and chin white.)

11. The Pied Crested Cuckoo (137). W. K. (White wing-bar ; lower plumage white ; tail tipped white.)

12. The Black Vulture (146). E. W. (Breast and thighs white ; bare head and wattles red.)

13. The Indian White-backed Vulture (148). E. W. (White back.)

14. The White-crested Kalij Pheasant (162). W. (Crest white ; wings and tail dark brown glossed with blue.)

15. The Black-backed Kalij Pheasant (163). E. (Wings and tail dark brown glossed with blue.)

16. The Black Partridge (166). W. K. (White on head and white spots and bars on body ; chestnut collar.)

---

---

## *Himalayan and Kashmiri Birds*

---

17. The Red-wattled Lapwing (170). E. W. K. (Some white in wings; lower parts white; white band down side of neck; crimson wattle; yellow legs.)

18. The Large Cormorant (172). K. (White band across throat.)

### *3. Black-headed birds*

(Birds which are *all* black are not included.)

1. The Indian House Crow (2). E. W. K.

2. The Jackdaw (3). K. (Nape and neck grey.)

3. The Red-billed Blue Magpie (4). W.

4. The Yellow-billed Blue Magpie (5). E. W. K.

5. The Black-throated Jay (7). W. K.

6. The Indian Grey Tit (11). E. W. K. (White cheeks.)

7. The Green-backed Tit (12). E. W. K. (White cheeks.)

8. The Yellow-cheeked Tit (14). W. (Cheeks, nape, and tip of crest yellow.)

9. The Crested Black Tit (15). W. K. (Cheeks and patch on nape white.)

10. The Black-headed Sibia (30). E. W.

11. The Red-tailed Minla (34). E. (Broad white eyebrow.)

---

---

*Classification according to Colour*

---

12. The Bengal Red-vented Bulbul (39). E.  
13. The Punjab Red-vented Bulbul (40). W.  
14. The White-cheeked Bulbul (41). E. W.  
K. (Cheeks white.)  
15. The Short-billed Minivet (57). E. W.  
16. The Common Myna (61). K. W.  
(Yellow patch of skin behind the eye.)  
17. The Jungle Myna (62). E. W.  
18. The Indian Paradise Flycatcher (73).  
W. K.  
19. The Common Pied Bush Chat (75).  
W. K. (Cock.)  
20. The Indian Bush Chat (75). W. K.  
(Cock.)  
21. The Dark-grey Bush Chat (76). E. W.  
K. (Cock) (White eyebrow.)  
22. The Western Spotted Forktail (77).  
W. K. (White patch on crown.)  
23. The Eastern Spotted Forktail (78). E.  
(White patch on crown.)  
24. The Little Forktail (79). E. W. K.  
(White patch on crown.)  
25. The White-capped Redstart (80). E. W.  
K. (Crown white.)  
26. The Magpie-Robin (84). W. (Cock.)  
27. The Black-and-yellow Grosbeak (89).  
W. K.

---

---

## *Himalayan and Kashmiri Birds*

---

28. Hodgson's Pied Wagtail (103). K.  
(Face white.)
29. The Common Indian Swift (130).  
W. K.
30. The Pied Crested Cuckoo (137). W. K.
31. The Black Vulture (146). E. W.
32. The Indian White-backed Vulture (148).  
E. W.
34. The Hobby (155). E. W. (Pale eye-  
brow and forehead.)
35. The Black-backed Kalij Pheasant (163).  
E. (Cock.)
36. The Black Partridge (166). W. K.  
(Cock) (Crown buff.)
37. The Red-wattled Lapwing (170). E. W.  
K. (Red wattles and white band on side of  
head.)
38. The Large Cormorant (172). K. (White  
band from eye to eye across throat.)
39. The Night Heron (175). K. (White  
forehead and cheeks.)
4. *Black collaret or gorget or band across the  
breast*
1. White-browed Blue Flycatcher (66). E.  
W. (Collaret is *dark blue* and interrupted in  
middle. Appears black from a distance.)

---

---

*Classification according to Colour*

---

2. The Swallow (100). E. W. K.
3. The European Bee-eater (126). K.
4. The Chukor (165). W. K.

5. *Black band running longitudinally along abdomen*

1. The Indian Grey Tit (11). E. W. K.
2. The Green-backed Tit (12). E. W. K.
3. The Yellow-cheeked Tit (14). W.

6. *Black streak through the eye (i.e. from bill to back of head)*

1. The Himalayan White-crested Laughing Thrush (17). E. W.
2. The Shrikes (55 and 56).
3. The Verditer Flycatcher (67). E. W.
4. The Blue-headed Rock-Thrush (87). E. W. K.
5. The European Bee-eater (126). K.
6. The Lammergeyer (150). W. K. (Black band runs from the eye to the bill and is continued as a "beard.")

7. *Birds having a black moustachial streak*

1. The Himalayan Jay (8). E. W. K.
2. The Yellow-naped Ixulus (32). E. W.

---

---

## *Himalayan and Kashmiri Birds*

---

### *8. Black and blue birds*

1. The Himalayan Whistling Thrush (29).  
E. W. K.

2. The Verditer Flycatcher (67). E. W.  
(A beautiful pale blue bird, with a black band  
through the eye.)

3. The Large Niltava (72). E. (Cock.)

4. The Indian Blue Rock-Pigeon (158).  
E. W. K. (A slaty-blue bird with black wing-  
bars and red legs.)

### *9. Black and brown birds*

(See Brown and Black Birds.)

### *10. Black and grey birds*

1. The Indian House Crow (2). E. W. K.

2. The Jackdaw (3). K.

3. The Himalayan Tree-pie (6). E. W.  
(Back and shoulders reddish-brown; patch  
of chestnut under tail.)

4. The Indian Grey Tit (11). E. W. K.  
(White cheeks and flanks; some white in  
wings.)

5. The Crested Black Tit (15). W. K.  
(Nape and band on sides of neck white.)



---

---

*Classification according to Colour*

---

6. The Himalayan Black Bulbul (37). E. W. K.

7. The Indian Ashy Drongo (44). E. W. K. (Upper plumage dark indigo ; lower plumage dark grey.)

8. The Dark Grey Cuckoo-shrike (58). E. W. (Tail tipped with white.)

9. The Grey-headed Myna (62). W. (Grey washed with red in parts.)

10. The Dark Grey Bush Chat (76). E. W. K. (Cock. Some white in wings, tail and lower plumage.)

11. The Grey-winged Ouzel (85). E. W. (Cock. Bill red.)

12. The Hobby (155). E. W.

13. The Whiskered Tern (171). K. (Bill red.)

*11. Black and red birds*

1. The Short-billed Minivet (57). E. W. (Cock.)

2. The Indian Paradise Flycatcher (73). W. K. (Hen and immature cock. Black and chestnut.)

3. The White-capped Redstart (80). E. W. K. (White cap.)

---

---

## *Himalayan and Kashmiri Birds*

---

4. The Scarlet Finch (90). E. (Cock.)
5. The Red-browed Finch (93). W. K. (Some white in wings and tail.)
6. The Black Vulture (146). E. W. (Breast and patch on thigh white.)

### *12. Black and white birds*

1. The Indian Paradise Flycatcher (73). W. K. (Cock. White bird with very long tail; head and crest black.)
2. The Common Pied Bush Chat (74). W. K. (Cock. Black bird with white wing-bar and patch on back.)
3. The Western Spotted Forktail (77). W. K. (Long-tailed white bird, much spotted and barred with black.)
4. The Eastern Spotted Forktail (78). E. (Like 3.)
5. The Little Forktail (79). E. W. K. (Like 3, but tail short.)
6. The Magpie-Robin (84). W. (Cock. Black and white as in the magpie.)
7. Hodgson's Pied Wagtail (103). K.
8. The Pied Woodpeckers (117-120). (Spotted black-and-white birds, which have a little red or yellow in plumage.)

---

---

## *Classification according to Colour*

---

9. The Himalayan Pied Kingfisher (127). E. W. K. (A spotted black-and-white bird.)
10. The Common Indian Swift (130). W. K. (A black bird with a white rump.)
11. The White-necked Spine-tail (131). E. W. (A blackish-brown bird having middle of back and chin white.)
12. The Pied Crested Cuckoo (137). W. K. (Upper parts black with white wing-bar ; lower parts white.)
13. The Black Vulture (146). E. W. (A black bird with white breast and thighs ; wattles and bare head and neck red.)
14. The Indian White-backed Vulture (148). E. W. (A black bird with white back and some white in the wings.)
15. The Coot (168). E. W. K. (A black duck-like bird with white bill and shield on forehead.)
16. The Large Cormorant (172). K. (A black bird with white band across throat.)

### *13. Black-and-yellow birds*

1. The Yellow-cheeked Tit (14). W. (Some olive-green and brown in tail and wings.)
2. The Short-billed Minivet (57). E. W. (Hen.)

---

---

## *Himalayan and Kashmiri Birds*

---

3. The Indian Oriole (59). W. K.
4. The Black-and-yellow Grosbeak (89). W. K. (Cock.)
5. Hodgson's Yellow-headed Wagtail (105). K. (Some brown in the wings.)

### 14. *Black, blue and white bird*

The Large Niltava (72). E.

### 15. *Black, brown and white birds*

(See Brown, Black and White Birds.)

### 16. *Black, fawn-colour and white bird*

The European Hoopoe (129). E. W. K.

### 17. *Black, grey, and white birds*

1. The Indian Grey Tit (11). E. W. K.
2. The Crested Black Tit (15). W. K.  
(A little chestnut on the wings.)
3. The Dark Grey Cuckoo-Shrike (58). E. W. (Very little white.)
4. The Dark Grey Bush Chat (76). E. W. K.
5. The Indian Cuckoo (135). E. W. (Back brown.)
6. The Indian Ring-Dove (161). W. K.  
(Very little black : only a half collar.)

---

---

*Classification according to Colour*

---

7. The Common Heron (174). K.

8. The Night Heron (175). K.

18. *Black, blue, white and chestnut birds*

1. The White-tailed Nuthatch (42). E. W.

2. The Rufous-bellied Niltava (71). E. W.  
K. (Cock.)

3. The Blue-headed Rock-Thrush (87). E.  
W. K. (Cock.)

4. The Swallow (100). E. W. K. (Very  
little white.)

5. The Wire-tailed Swallow (101). W. K.

6. Sykes's Striated Swallow (102). E. W.  
(Very little white.)

19. *Black, blue, white and brown birds*

1. The Red-billed Blue Magpie (4). W.

2. The Yellow-billed Blue Magpie (5). E.  
W. K.

3. The White-browed Blue Flycatcher (66).  
E. W. (Cock.)

20. *Black, blue, white and crimson bird*

The Red-flanked Bush Robin (83). E. W. K.

---

---

## *Himalayan and Kashmiri Birds*

---

### *21. Black, blue, white, and red bird*

The Common Kingfisher (128). E. W. K.  
(Very little white on side of face.)

### *22. Black, grey, white and chestnut bird*

The Crested Black Tit (15). W. K.

## II. BLUE

1. Birds having blue in the plumage.
2. Blue and red birds.

### *1. Birds having blue in the plumage*

*See* I 8. Black and blue birds.

I 14. Black, blue and white bird.

I 18. Black, blue, white and chestnut birds.

I 19. Black, blue, white and brown birds.

I 20. Black, blue, white and crimson bird.

I 21. Black, blue, white and red bird.

1. The Red-billed Blue Magpie (4).  
*See* I 19.

2. The Yellow-billed Blue Magpie (5).  
*See* I 19.

---

---

*Classification according to Colour*

---

3. The Black-throated Jay (7). W. K.  
(Some blue in wing.)
4. The Himalayan Jay (8). E. W. K.  
(Some blue in the wing.)
5. The Himalayan Whistling Thrush (29).  
*See* I 8.
6. The White-tailed Nuthatch (42). *See*  
I 18.
7. The White-browed Blue Flycatcher (66).  
*See* I 19.
8. The Verditer Flycatcher (67). *See* I 8.
9. The Rufous-bellied Niltava (71). *See* I 18.
10. The Large Niltava (72). *See* I 8.
11. The Blue-fronted Redstart (81). E. W.  
K. (Cock. Tail chestnut-red.)
12. The Plumbeous Redstart (82). E. W. K.  
(Cock. Tail rusty red).
13. The Red-flanked Bush-Robin (83).
14. The Blue-headed Rock-Thrush (87).  
(Cock.) *See* I 18.
15. The Red-browed Finch (93). W. K.  
(Blue-black in wing.)
16. The Swallow (100). *See* I 18.
17. The Wire-tailed Swallow (101). *See* I 18.
18. Sykes's Striated Swallow (102). *See* I 18.
19. The Fire-tailed Yellow-backed Sunbird  
(111). E. (Cock. A metallic-blue crown.)

---

---

## *Himalayan and Kashmiri Birds*

---

20. The Great Himalayan Barbet (123). E. W. (Head and neck violet-blue.)
21. The Golden-throated Barbet (124). E. (Plumage largely green washed with blue.)
22. The European Roller (125). K. (A brilliant light and dark blue bird.)
23. The Common Kingfisher (128). E. W. K. (Plumage largely blue. *See* I 21.)
24. The Western Blossom-headed Paroquet (140). W. K. (Plumage mainly bright green ; some blue in tail.)
25. The Slaty-headed Paroquet (141). E. W. K. (Plumage mainly bright green ; some blue in tail.)
26. The Indian Blue Rock-Pigeon (158). *See* I 8.
27. The White-crested Kalij Pheasant (162). W. (The black in the plumage is glossed with blue.)
28. The Black-backed Kalij Pheasant (163). E. (The black in the plumage is glossed with blue.)

### *2. Blue and red birds*

1. The Blue-fronted Redstart (81). E. W. K. (A little black in the wing.)
2. The Plumbeous Redstart (82). E. W. K.



---

---

## *Classification according to Colour*

---

### III. BROWN

1. Birds of which the general hue is earthy-brown.

- (a) Birds smaller than the sparrow.
- (b) Birds about the size of the sparrow or the bulbul.
- (c) Birds about the size of the myna.

2. Birds of which the plumage is largely deep or rich brown.

- (a) Chocolate.
- (b) Umber.
- (c) Olive-brown.
- (d) Reddish-brown.
- (e) Dark brown.

The number of birds of which the predominating hue is one or more of the many shades of brown is large, and, as most of these have nothing very striking in their plumage, their identification is by no means easy.

Birds which appear to be a uniform earthy-brown are usually of lighter hue below than above. This difference is not noticed when the bird is in its natural surroundings because

---

---

## *Himalayan and Kashmiri Birds*

---

the lower parts are in the shade and so appear darker than they really are. As it is often of advantage to a bird to be inconspicuous, natural selection has doubtless been responsible in many cases for the fact that the lower plumage is of paler hue than the upper.

As this book is intended for the field observer I shall describe these brown birds as they appear to the observer in the field and so describe simply as an earthy-brown bird one which is earth-coloured above and creamy below.

1. *Birds of which the general hue is earthy-brown*

(a) *Birds smaller than the sparrow*

1. The Brown Crested Tit (16). W. K. (Brownish-grey above with a creamy band on each side of the neck. Tinted with yellow below, crested.)

2. The Nepal Babbler (26). E. (Uniform grey-brown with a ring of white feathers round the eye.)

3. The Kashmir Wren (47). K. W. (Reddish-brown barred with black.)

4. The Scaly-breasted Wren (48). E. W. (Rich brown, dotted with buff; black scaly markings on lower plumage.)

---

---

*Classification according to Colour*

---

5. The Sooty Flycatcher (64). E. W. K. (Spotted on head. Ring of white round the eye.)

6. The White-browed Blue Flycatcher (66). E. W. (Hen.) (Olive-brown tinged with dark blue on the lower back.)

7. The Brown Flycatcher (68). W. K. (Ring of white feathers round the eye.)

8. The Rufous-tailed Flycatcher (69). W. K. (Tail chestnut.)

9. The Indian Bush Chat (75). W. K. (Hen.)

(b) *Birds about the size of the Sparrow or the Bulbul*

1. The Himalayan Streaked Laughing-Thrush (23). W. K. (Streaked with black and white; some chestnut on cheeks and lower parts.)

2. The Himalayan Tree-creeper (45). W. K. (Streaked light and dark brown.)

3. The Sikkim Tree-creeper (46). E.

4. The Indian Lesser White-throated Warbler (50). K.

5. The Brown Hill-Warbler (54). E. W. K.

6. The Blue-headed Rock-Thrush (87). E. W. K. (Hen. Olive-brown.)

---

---

## *Himalayan and Kashmiri Birds*

---

7. The Red-mantled Rose-Finch (91). W. K. (Hen.)
8. The House Sparrow (95). W. K. (Hen.)
9. The Cinnamon Tree-Sparrow (97). W. K. (Hen. Reddish-brown.)
10. The Indian Tree-Pipit (106). W. (Some white in tail.)
11. The Brown Rock-Pipit (107). W. (Grey-brown with dark streaks.)
12. The Indian Pipit (108). E. W. K. (Dark streaks.)
13. The Upland Pipit (109). E. W. (Dark brown with buff streaks.)
14. The Skylark (110). W. K. (Dark brown with pale streaks.)
15. The Common Wryneck (122). K. (Grey-brown with dark bars.)
16. The Spotted Himalayan Scops Owl (143). E. W. (Mottled brown; large aigrettes.)
17. The Collared Pigmy Owlet (145). E. W. (Mottled brown. Buff collar.)

### *(c) Birds about the size of the Myna*

1. The Large Hawk-Cuckoo (136). E. W. (Plumage with dark cross-bars.)
2. The Himalayan Wood-Owl (142). E. W.

---

---

## *Classification according to Colour*

---

(A good deal larger than a myna. Plumage mottled.)

3. The Large Barred Owlet (144). E. W. (Smaller than a myna. Plumage has pale cross-bars.)

4. The Little Brown Dove (160). W. (Smaller than Myna. Red-and-black chess-board on each side of neck.)

5. The Common Grey Quail (164). E. W. (A good deal smaller than a myna ; legs short ; tail short ; plumage spotted and streaked with black.)

2. *Birds of which the plumage is largely deep or rich brown*

(a) *Chocolate*

1. The Larger Spotted Nutcracker (10). W. K. (With large white spots.)

2. The Brown Dipper (88). E. W. K.

3. The Crested Serpent-Eagle (152). E. W. (Cream cross-bars on wings and tail. Large white spots on lower parts.)

(b) *Umber*

1. The Himalayan Nutcracker (9). W. (With white spots and streaks.)

---

---

## *Himalayan and Kashmiri Birds*

---

2. The Striated Laughing-Thrush (24). E. W. (Streaked with white; tail chestnut; some chestnut in wing.)

### (c) *Olive-brown*

1. The Himalayan White-crested Laughing-Thrush (17). E. W.

2. The White-throated Laughing-Thrush (18). E. W. (Under parts rust coloured).

3. The Rufous-chinned Laughing-Thrush (19). E. W. K. (Under parts ashy brown.)

4. The Red-headed Laughing-Thrush (20). W. (Some bright golden yellow in wing.)

5. The Eastern Variegated Laughing-Thrush (22). W. (Much golden-yellow in wing.)

6. The Rusty-cheeked Scimitar Babbler (25). E. W. (Bill rather long.)

7. The Indian Great Reed-warbler (49). K.

8. The White-browed Blue Flycatcher (66). E. W. (Hen. Tinged with blue on lower back.)

9. The Rufous-tailed Flycatcher (69). W. K.

10. The Rufous-bellied Niltava (71) E. W. K. (Hen. White patch on fore neck and a blue patch on each side of neck.)

---

---

*Classification according to Colour*

---

11. The Large Niltava (72). E. (Hen.)  
(Blue patch on each side of neck.)
12. The Red-flanked Bush-Robin (83). E.  
W. K. (Hen. White bib ; some blue on tail  
and chestnut on sides of body.)
13. The Grey-winged Ouzel (85). E. W.  
(Hen. Bright red bill.)
14. Tickell's Ouzell (86). K. (Hen. Bill  
yellow.)
15. The Blue-headed Rock-Thrush (87). E.  
W. K. (Hen.)
16. The Fire-breasted flower-pecker (113).  
E. W. (Hen.)
17. The Moorhen (167). K. (Only back  
and shoulders olive-brown. Red shield on  
forehead.)

(d) *Reddish-brown*

1. The Himalayan Streaked Laughing-  
Thrush (23). W. K. (Black and white streaks  
in plumage.)
2. The Kashmir Wren (47). W. K. (Closely  
cross-barred with black ; tail very short.)
3. The Ferruginous Flycatcher (65). E.  
(Cream-coloured ring round eye.)
4. The Cinnamon Tree-Sparrow (97) W.  
K. (Hen.)

---

---

## *Himalayan and Kashmiri Birds*

---

5. The White-capped Bunting (98). W. K.
6. The Eastern Meadow Bunting (99). W. K.
7. The Chukor (165). W. K. (Bill and legs red.)
8. The White-eyed Pochard (178). K. (Head, neck and breast.)

### (e) *Dark brown*

1. The Bengal Red-vented Bulbul (39). E. (Creamy crescentic markings ; red patch under tail.)
2. The Punjab Red-vented Bulbul (40). W. (Creamy crescentic markings ; red patch under tail.)
3. The White-cheeked Bulbul (41). E. W. K. (Creamy crescentic markings ; sulphur-yellow patch under tail.)
4. The Himalayan Tree-creeper (45). W. K. (Yellowish-brown streaks.)
5. The Sikkim Tree-creeper (46). E. (Yellowish-brown streaks.)
6. The Scaly-breasted Wren (48). E. W. (Short tail.)
7. The Brown Hill-warbler (54). E. W. K. (Lower parts pale ruddy-yellow.)
8. The Common Myna (61). K. W. (Yellow legs ; white wing-bar.)



---

---

*Classification according to Colour*

---

9. The Jungle Myna (63). E. W. (Yellow legs ; white wing-bar.)
10. The Indian Bush Chat (75). W. K.
11. The Plumbeous Redstart (82). E. W. K. (Hen. The brown is tinged with blue in parts.)
12. The Scarlet Finch (90). E. (Hen. Rump bright yellow).
13. The Upland Pipit (109). E. W. (Crescentic reddish-buff markings.)
14. The Skylark (110). W. K. (Mottled.)
15. The Common Indian Swift (130). W. K. (Very dark brown ; almost black. White rump.)
16. The White-necked Spine-tail (131). E. W. (Blackish-brown, glossed with green.)
17. The Large Hawk-Cuckoo (136). E. W.
18. The Indian Koel (138). W. K. (Hen. Spotted and barred with white.)
19. The Owls (142-145). (Plumage mottled and barred.)
20. The Indian White-backed Vulture (148). E. W. (Very dark brown ; almost black.)
21. Pallas's Fishing Eagle (153). E. W. K.
22. The Common Pariah Kite (154). E. W. K.
23. The White-crested Kalij Pheasant (162).

---

---

## *Himalayan and Kashmiri Birds*

---

W. (Cock. Wings and tail brown, glossed with green.)

24. The Black-backed Kalij Pheasant (163).

E. (Cock. Wings and tail brown, glossed with green.)

25. The Common Grey Quail (154). E. W.

26. The Black Partridge (166). W. K. (Hen.)

27. The Mallard (177). K. (Duck.) (Buff crescentic markings.)

28. The White-eyed Pochard (178). K.

29. The Indian Little Grebe (179). K. (Very dark rich brown ; chestnut on sides of neck.)

### IV. FAWN COLOUR

#### *Birds having fawn-colour in plumage*

1. The Black-throated Jay (7). W. K. (Back fawn-colour.)

2. The Himalayan Jay (8). E. W. K. (Plumage mainly fawn-colour.)

3. The European Hoopoe (129). E. W. K. (Body plumage fawn ; wings boldly banded black and white.)

---

---

*Classification according to Colour*

---

V. GREEN

1. Birds in the plumage of which bright green occurs.

2. Birds in the plumage of which dull green occurs.

(a) Dull yellow-green.

(b) Olive-green.

1. *Birds in the plumage of which bright green occurs*

1. The Great Himalayan Barbet (123). E. W.

2. The Golden-throated Barbet (124). E. (Plumage mainly grass-green washed with blue.)

3. The European Bee-eater (126). K. (Two middle tail feathers prolonged as bristles.)

4. The Paroquets (139-141). (Plumage mainly grass-green. Tail long.)

5. The Kokla Green Pigeon (157). E. W. K. (Head and lower plumage yellowish-green.)

2. *Birds in plumage of which dull green occurs*

(a) *Dull yellow-green*

1. The Green-backed Tit (12). E. W. K.

2. The Red-tailed Minla (34). E.

3. The Himalayan Greenfinch (94). E. W. K.

---

---

## *Himalayan and Kashmiri Birds*

---

4. The Grey Wagtail (104). K.
5. The Fire-tailed Yellow-backed Sunbird (111). E. (Hen.)
6. The Green Woodpeckers (114 and 115).

### (b) *Olive-green*

1. The Yellow-cheeked Tit (14). W.
2. The Red-headed Babbler (28). E.
3. The Red-billed Liothrix (33). E. W.
4. The Fire-cap (35). W. (Cock.)
5. The Striated Green Bulbul (38). E.
6. Hume's Willow-warbler (51). W. K.
7. Hodgson's Grey-headed Flycatcher-warbler (52). W.
8. Brook's Grey-headed Flycatcher-warbler (53). E.
9. The Nepal Yellow-backed Sunbird (112). E. (Hen.)
10. The Kokla Green Pigeon (157). E. W. K.

## VI. GREY

1. Birds of which the plumage is largely grey.
2. Birds having the head only grey.
3. Grey and black birds.

---

---

## *Classification according to Colour*

---

4. Grey, black and white birds.
5. Grey and red bird.
6. Grey and white birds.

1. *Birds of which the plumage is largely grey*

1. The Indian Grey Tit (11). E. W. K.  
*See I 5.*
2. The Rufous-backed Shrike (55). E. W. K. (Head, neck and upper back pale grey.)
3. The Grey-backed Shrike (56). E. W. K. (Head, neck, shoulders and most of back grey.)
4. The Dark Grey Cuckoo-Shrike (58). E. W. (Body dark grey ; wings and tail black.)
5. The Grey-headed Myna (62). W.
6. The Grey-headed Flycatcher (70). E. W. K. (Head, neck and breast grey ; lower plumage bright yellow.)
7. The Common Pied Bush Chat (74). W. K. (Hen.)
8. The Magpie-Robin (84). W. (Hen.)
9. Tickell's Ouzel (86). K. (Cock. Plumage dark grey.)
10. Black-and-yellow Grosbeak (89). W. K. (Hen.)
11. The Grey Wagtail (104). K. (Head, neck and shoulders bluish-grey ; lower parts bright yellow.)

---

---

## *Himalayan and Kashmiri Birds*

---

12. The Cuckoo (132). E. W. K.
13. The Himalayan Cuckoo (133). E. W.
14. The Small Cuckoo (134). E. W.
15. The Indian Cuckoo (135). E. W.
16. The Slaty-headed Paroquet (141). E. W. K. (A long-tailed, grass-green bird, with a grey head.)
17. The Lammergeyer (150). W. K.
18. The Hobby (155). E. W.
19. The Kestrel (156). E. W. K. (Head, neck and tail grey ; back and wings pale brick-red.)
20. The Indian Turtle-Dove (159). W. K.
21. The Indian Ring-Dove (161). W. K.
22. The Moorhen (167). K. (Head and breast dark slaty grey.)
23. The Whiskered Tern (171). K. (Silvery grey with black cap and abdomen.)
24. The Common Heron (174). K.
25. The Night Heron (175). K.

### *2. Birds having the head only grey*

1. Hodgson's Grey-headed Flycatcher-warbler (52). W.
2. Brooks's Grey-headed Flycatcher-warbler (53). E.
3. The Shrikes (55 and 56). (These have

---

---

## *Classification according to Colour*

---

some of the back grey also, and a black band through eye.) *See* I 6.

4. The Grey-headed Flycatcher (70). E. W. K. (Neck also grey.)

### *3 Grey and black birds*

(*See* Black and grey birds.)

### *4. Grey, black and white birds*

*See* (Black, grey, and white birds.)

### *5. Grey and red birds*

The Kestrel (156). E. W. K. (Wings dull red.)

### *6. Grey and white birds*

1. The Magpie Robin (184). W. (Hen.)

2. The Common Heron (174). K.

## VII. RED

1. Birds having red in the plumage or red wattles or red skin on the head.

(a) Bright red.

(b) Dull red (chestnut, bay, maroon.)

2. Red-billed birds.

---

---

## *Himalayan and Kashmiri Birds*

---

3. Red-legged birds.
4. Birds having patch of bright red feathers under the tail.
5. Bird having patch of chestnut feathers under the tail.
6. Birds having patch of red on shoulder.
7. Red and black birds.

1. *Birds having red in the plumage or red wattles  
or red skin on the head*

(a) *Bright red*

1. The Red-billed Liothrix (33). E. W. (Cock. Wing narrowly margined with crimson.)
2. The Red-tailed Minla (34). E. (Some crimson in wings and tail.)
3. The Fire-cap (35). W. (Cock. Cap flaming red.)
4. The Bengal Red-vented Bulbul (39). E. (Crimson patch under tail.)
5. The Punjab Red-vented Bulbul (40). W. (Crimson patch under tail.)
6. The Short-billed Minivet \* (57). E. W. (Cock. Much crimson in plumage.)
7. The White-capped Redstart \* (80). E. W. K. (Tail bright chestnut-red.)

\* These birds are conspicuously red.



---

---

*Classification according to Colour*

---

8. The Blue-fronted Redstart \* (81). E. W. K. (Tail chestnut-red.)
9. The Plumbeous Redstart \* (82). E. W. K. (Cock. Tail rusty red.)
10. The Red-flanked Bush-Robin (83). E. W. K. (A patch of bright chestnut on each flank.)
11. The Scarlet Finch \* (90). E. (Cock.)
12. The Red-mantled Rose-Finch (91). W. K. (Cock.)
13. The Himalayan Goldfinch (92). W. K. (Some crimson on head.)
14. The Red-browed Finch (93). W. K. (Cock. Some crimson on head.)
15. The Fire-tailed Yellow-backed Sunbird (111). E. (Cock. Much crimson on upper plumage.)
16. The Fire-breasted Flower-pecker (113). E. W. (Cock. Patch of crimson on breast.)
17. The Woodpeckers (114-121). (Cocks have bright red on head and some cocks and hens have red patch under tail.)
18. The Great Himalayan Barbet (123). E. W. (Red patch under tail.)
19. The Golden-throated Barbet (124). E. (Forehead crimson.)

\* These birds are conspicuously red.

---

---

## *Himalayan and Kashmiri Birds*

---

20. The Common Kingfisher (128). E. W. K. (Some red on face; lower parts rust-coloured.) *See* I 21 and II.

21. The Paroquets (139-141). (Red patch on Wing.) *See* Green birds.

22. The Black Vulture (146). E. W. (Wattles and bare head and neck red.)

23. The Red-wattled Lapwing (170). E. W. K. (Wattles crimson.)

### (b) *Dull red (chestnut, bay, maroon)*

1. The Himalayan Tree-pie (6). E. W. (Patch of chestnut under tail.) *See* Birds having tail long in proportion to body.

2. The Red-headed Tit (13). E. W. (Top of head and nape chestnut.)

3. The White-throated Laughing-Thrush (18). E. W. (Lower plumage rust-coloured.)

4. The Rufous-chinned Laughing-Thrush (19). E. W. K. (Cheeks, chin and upper throat red.)

5. The Red-headed Laughing-Thrush (20). W. (Top of head chestnut.)

6. The Himalayan Streaked Laughing-Thrush (23). W. K. (Cheeks, chin, throat and breast chestnut.)

---

---

*Classification according to Colour*

---

7. The Striated Laughing-Thrush (24). E. W. (Tail chestnut ; chestnut in wing.)
8. The Rusty-cheeked Scimitar Babbler (25). E. W. (Forehead and cheeks rust-coloured.) *See* Birds with long bills.
9. The Red-headed Babbler (28). E. (Head chestnut.)
10. The Black-headed Sibia (30). E. W. (Lower plumage rust-coloured.) *See* I 3.
11. The Yellow-naped Ixulus (32). E. W. (Chestnut half-collar.)
12. The White-tailed Nuthatch (42). E. W. (Lower parts chestnut.)
13. The Rufous-backed Shrike (55). E. W. K. (Shoulders and lower back rusty red.)
14. The Grey-backed Shrike (56). E. W. K. (Lower back rusty red.)
15. The Ferruginous Flycatcher (65). E.
16. The Rufous-tailed Flycatcher (69). W. K. (Tail chestnut.)
17. The Rufous-bellied Niltava (71). E. W. K. (In cock lower plumage chestnut ; in hen tail chestnut.)
18. The Large Niltava (72). E. (Hen has deep rufous in tail.)
19. The Indian Paradise Flycatcher (73). W. K. (Hen and young cock have body and

---

---

## *Himalayan and Kashmiri Birds*

---

tail pale chestnut.) *See* Birds having tail long in proportion to body.

20. The Indian Bush Chat (75). W. K. (Lower plumage russet.)

21. The Dark Grey Bush Chat (76). E. W. K. (Hen has some dull red on rump and tail.)

22. The Redstarts (80–82).

23. The House-Sparrow (95). W. K. (Cock has dark chestnut on upper back.)

24. The Tree-Sparrow (96). E. (Pale chestnut on shoulders and upper back.)

25. The Cinnamon Tree-Sparrow (97). W. K. (Cock has light cinnamon upper plumage.)

26. The Swallow (100). E. W. K. (Forehead, chin and throat chestnut.) *See* Blue birds.

27. The Wire-tailed Swallow (101). W. K. (Head chestnut.) *See* Blue birds.

28. Sykes's Striated Swallow (102). E. W. (Chestnut collar and rump.) *See* Blue birds.

29. The Nepal Yellow-backed Sunbird (112). E. (Cock.) (Upper back maroon.)

30. The Kestrel (156). E. W. K. (Back and wings brick-red.) *See* Grey birds.

31. The Kokla Green-pigeon (157). E. W. K. (Cock has middle back maroon.)

---

---

## *Classification according to Colour*

---

32. The White-eyed Pochard (178). K.  
(Head, neck and breast dull dark chestnut.)

33. The Little Grebe (179). K. (Chestnut  
on sides of neck.)

### *2. Red-billed birds*

1. The Red-billed Blue Magpie (4). W.

2. The Red-billed Liothrix (33). E. W.

3. The Himalayan Black Bulbul (37). E.  
W. K.

4. The Grey-winged Ouzel (85). E. W.

5. The Large Indian Paroquet (139). E. W.

6. The Chukor (65). W. K.

7. The Moorhen (167). K. (Tip of bill  
greenish-yellow.)

8. The Red-wattled Lapwing (170). E. W.  
K.

9. The Whiskered Tern (171). K.

### *3. Red-legged birds*

1. The Red-billed Blue Magpie (4). W.

2. The Himalayan Black Bulbul (37). E.  
W. K.

3. The Common Kingfisher (128). E. W. K.

4. The Indian Blue Rock-Pigeon (158).  
E. W. K. (Legs reddish-pink.)

5. The Doves (159-161.)

---

---

## *Himalayan and Kashmiri Birds*

---

6. The Chukor (165). W. K.

7. The Black-bellied Tern (171). (Orange-red.)

4. *Birds having patch of bright red feathers under the tail*

1. The Bengal Red-vented Bulbul (39). W.

2. The Punjab Red-vented Bulbul (40). E.

3. The Woodpeckers (114-121).

4. The Great Himalayan Barbet (123). E. W.

5. *Bird having patch of chestnut feathers under the tail*

The Himalayan Tree-pie (6). E. W.

6. *Birds having patch of red on shoulder*

1. The Tree-Sparrow (96). E.

2. The Large Indian Paroquet (139). E. W.  
*See Green birds.*

3. The Western Blossom-headed Paroquet (140). W. K. *See Green birds.*

4. The Slaty-headed Paroquet (141). (Cock.)  
*See Green birds.*

7. *Red and black birds*

(*See Black and red birds.*)

---

---

*Classification according to Colour*

---

VIII. WHITE

1. Birds having much white in the plumage.
2. Birds having numerous white spots in the plumage.
3. Birds having a white bar in the wing ; seen during flight.
4. Birds having white cheeks.
5. Birds having ring of white feathers round the eye.
6. Bird having a snow-white cap.

1. *Birds having much white in the plumage*

1. The Himalayan White-crested Laughing-Thrush (17). E. W. (Crested head and throat white with black streak through eye.)
2. The Indian Paradise Flycatcher (73). W. K. (Cock. All white except crested head ; tail very long.)
3. The Forktails (77-79). *See* Black and white birds.
4. The Magpie-Robin (84). W.
5. Hodgsons' Pied Wagtail (103). K.
6. The Himalayan Pied Kingfisher (127). E. W. K.
7. The European Hoopoe (129). E. W. K.

---

---

## *Himalayan and Kashmiri Birds*

---

(Wings and tail boldly banded black and white.)

*See* Crested birds.

8. The Pied Crested Cuckoo (137). W. K.  
*See* Crested Birds.

9. The Large White Scavenger Vulture (149). W. K.

10. Pallas's Fishing Eagle (153). E. W. K.  
(Head, chin and throat white. White band across tail.)

11. The Pheasant-tailed Jaçana (169). K.

12. The Red-wattled Lapwing (170). E. W. K.

13. The Grey Heron (174). K. (Lower parts white; some white in the head.)

14. The Night Heron (175). K. (Forehead, cheeks and under parts white.)

### *2. Birds having numerous white spots in the plumage*

1. The Himalayan Nutcracker (9). W.  
(On the wing body looks white and wings black.)

2. The Larger Spotted Nutcracker (10). W. K.

3. The Spotted-wing (36). W. (Cock.)

4. The Forktails (77-79). *See* Black and white birds.



---

---

*Classification according to Colour*

---

5. The Pied Woodpecker (117-120).

6. The Himalayan Pied Kingfisher (127).

E. W. K. *See* Black and white birds.

7. The Indian Koel (138). W. K. (Hen.)

8. The Black Partridge (166). W. K.  
(Cock.)

3. *Birds having a white bar in the wing ; seen  
during flight*

1. The Himalayan Tree-pie (6). E. W.

2. The Black-headed Sibia (30). E. W.

3. The Rufous-backed Shrike (55). E.  
W. K.

4. The Common Myna (61). K. W.

5. The Jungle Myna (63). E. W.

6. The Common Pied Bush Chat (74).  
W. K. (Cock.)

7. The Magpie-Robin (84). W.

8. The Blue-headed Rock-Thrush (87). E.  
W. K. (Cock.)

9. The Pied Crested Cuckoo (137). W. K.

10. The Indian White-backed Vulture (148).  
E. W. (The wing bar runs longitudinally, not  
transversely as in other cases.)

11. The Red-wattled Lapwing (170). E.  
W. K.

---

---

## *Himalayan and Kashmiri Birds*

---

### *4. Birds having white cheeks*

1. The Indian Grey Tit (11). E. W. K.
2. The Green-backed Tit (12). E. W. K.
3. The Crested Black Tit (15). W. K.
4. The White-throated Laughing-Thrush (18). E. W.
5. The Black-throated Babbler (27). E.
6. The White-cheeked Bulbul (41). E. W. K.
7. The House-Sparrow (95). W.K. (Cock.)
8. The Tree-Sparrow (96). E.
9. The Black Partridge (166). W. K. (Cock.)
10. The Red-wattled Lapwing (170). E. W. K. (A white band runs from eye down the neck.)

### *5. Birds having ring of white feathers round the eye*

1. The Nepal Babbler (26). E.
2. The Indian White-eye (31). E. W. K.
3. The Yellow-naped Ixulus (32). E. W.
4. The Sooty Flycatcher (64.) E. W. K.
5. The Ferruginous Flycatcher (65). E. (Ring cream-coloured.)

---

---

*Classification according to Colour*

---

6. The Brown Flycatcher (68). W. K.
7. The Blue-fronted Redstart (81). E. W. K. (Hen. Ring cream-coloured.)

6. *Bird having a snow-white cap*

The White-capped Redstart (80). E. W. K.

## IX. YELLOW

1. Birds having bright yellow or gold in the plumage.

2. Birds having bright yellow or orange legs.

3. Birds having dull-coloured upper plumage and bright yellow under parts.

4. Birds having patch of yellow feathers under the tail.

1. *Birds having bright yellow or gold in the plumage*

1. The Green-backed Tit (12). E. W. K.

2. The Yellow-cheeked Tit (14). W.

3. The Red-headed Laughing-Thrush (20). W.	} Golden-yellow in wing.
4. The Western Yellow-winged Laughing-Thrush (21). E.	
5. The Eastern Variegated Laughing-Thrush (22). W.	

---

---

*Himalayan and Kashmiri Birds*

---

6. The Black-throated Babbler (27). E.  
(Lower plumage bright reddish-yellow.)
7. The Indian White-eye (31). E. W. K.
8. The Red-billed Liothrix (33). E. W.
9. The Fire-cap (35). W.
10. The Striated Green Bulbul (38). E.  
(Yellow patch under tail.)
11. The White-cheeked Bulbul (41). E. W. K.  
(Yellow patch under tail.)
12. Hodgson's Grey-headed Flycatcher-warbler (52). W.
13. Brooks's Grey-headed Flycatcher-warbler (53). E.
14. The Short-billed Minivet (57). E. W.  
(Hen.)
15. The Indian Oriole (59). W. K.
16. The Grey-headed Flycatcher (70). E. W. K.
17. The Black-and-yellow Grosbeak (89). W. K.
18. The Scarlet Finch (90). E. (Hen. Rump bright yellow.)
19. The Himalayan Goldfinch (92). W. K.  
(Some gold in wing.)
20. The Himalayan Greenfinch (94). E. W. K.
21. The Grey Wagtail (104). K.

---

---

*Classification according to Colour*

---

22. Hodgson's Yellow-headed Wagtail (105). K.
23. The Fire-tailed Yellow-backed Sunbird (111). E. (Cock.)
24. The Nepal Yellow-backed Sunbird (112). E. (Cock.)
25. The Brown-fronted Pied Woodpecker (120). W. K. (Cock. Some gold in the crest.)
26. The Golden-throated Barbet (124). E.
27. The Pheasant-tailed Jaçana (169). K. (Back of neck golden-yellow.)

2. *Birds having bright yellow or orange legs*

1. The Common Myna (61). K. W.
2. The Jungle Myna (63). E. W.
3. Tickell's Ouzel (86). K.
4. The Cuckoo (132). E. W. K.
5. The Himalayan Cuckoo (133). E. W.
6. The Small Cuckoo (134). E. W.
7. The Indian Cuckoo (135). E. W.
8. The Large Hawk-Cuckoo (136). E. W.
9. The Hobby (153). E. W.
10. The Kestrel (156). E. W. K.
11. The Red-wattled Lapwing (170). E. W. K.
12. The Mallard (177). K. (Orange red.)

---

---

## *Himalayan and Kashmiri Birds*

---

### *3. Birds having dull-coloured upper plumage and bright-yellow under parts*

1. The Green-backed Tit (12). E. W. K. (Deep yellow with black median band.)
2. The Yellow-cheeked Tit (14). W. (Black median band.)
3. The Black-throated Babbler (27). E. (Reddish-yellow.)
4. Hodgson's Grey-headed Flycatcher-warbler (52). E.
5. Brooks's Grey-headed Flycatcher-warbler (53). E.
6. The Short-billed Minivet (57). E. W. (Hen.)
7. The Grey-headed Flycatcher (70). E. W. K.
8. The Himalayan Greenfinch (94). E. W. K.
9. The Grey Wagtail (104). K.

### *4. Birds having patch of yellow feathers under the tail*

1. The Striated Green Bulbul (38). E.
2. The White-cheeked Bulbul (41). E. W. K.

---

---

*Classification according to Colour*

---

---

X. BIRDS OF MANY COLOURS

1. The Fire-tailed Yellow-backed Sunbird (111). E. (Cock.)
2. The Nepal Yellow-backed Sunbird (112). E. (Cock.)
3. The Great Himalayan Barbet (123). E. W.
4. The Golden-throated Barbet (124). E.
5. The European Bee-eater (126). K.
6. The Common Kingfisher (128). E. W. K.
7. The Kokla Green Pigeon (157). E. W. K.
8. The Mallard (177). K. (Drake.)

---

## C. CLASSIFICATION ACCORDING TO HABITS

---

### I. BIRDS HAVING REMARKABLE CRIES

Description of Call	Name of Bird	No. of Bird in Part II
A short, sharp "Jack" . . .	{	The Jack- daw . . . 3
A sibilant <i>ti-ti-chee</i> or <i>pretty-pretty</i> . . . . .	}	The Indian Grey Tit . 11
(All the tits have similar calls.)		
A plaintive <i>poor-wee-wee</i> , the second <i>wee</i> very high-pitched. . . . .	{	The Red- headed Laughing- Thrush . 20
Call like the ringing of a little bell . . . . .	{	The Red- headed Babbler . 28
Loud, pleasant whistle . . .	{	The Hima- layan Whist- ling-Thrush 29



## Classification according to Habits

High-pitched, loud, penetrating metallic <i>titeree, titeree, tweeyo</i> . . . . .	{	The Black-headed Sibia . . . . .	30
Loud, pleasant, cheeping call . . . . .	{	The Indian White-eye . . . . .	31
<i>Tee-tee-tee-tee-tee</i> . (In breeding season cock sings well.) . . . . .	{	The Red-billed Liotrith . . . . .	33
Pleasant, tinkling, twittering . . . . .	{	The Bulbuls . . . . .	39-41
A cheery whistle, heard often at dawn, which has been described as “ <i>chēyk, chěchi, chēyk, chēȳk, chěchi, chēȳ, chēyh</i> ” . . . . .	{	The Black Drongo . . . . .	43
Like the above, but less harsh . . . . .	{	The Indian Ashy Drongo . . . . .	44
A shrill <i>pee-pee</i> . . . . .	{	The Himalayan Tree-creeper . . . . .	45
A loud call of four or five high-pitched notes. Very common in spring in the Western Himalayas . . . . .	{	Hodgson's grey-headed Flycatcher-warbler . . . . .	52
Like noise of sawing. . . . .	{	The Brown Hill-warbler . . . . .	54

---

## *Himalayan and Kashmiri Birds*

---

Soft, mellow <i>pee-ho</i> . . . .	{	The Indian Oriole . . . .	59
<i>Keeky, keeky, keeky . . . churr, churr, kok, kok, kok</i> . . . .	{	The Common Myna . . . .	61
A pleasant trill of about a dozen notes . . . . .	{	The Verditer Flycatcher . . . .	67
A loud, pleasant call of three notes: <i>Think-of-me</i> . . . . .	{	The grey-headed Flycatcher . . . .	70
Sweet little warble . . . . .	{	The Common Pied Bush Chat . . . .	74
A sweet and powerful song . . . . .	{	The Magpie-Robin . . . .	84
A fine, blackbird-like song . . . . .	{	The Grey-winged Ouzel . . . .	85
Ditto	{	Tickell's Ouzel . . . .	86
A sweet, soft song . . . . .	{	The blue-headed Rock-Thrush . . . .	87
<i>Kui-Kree</i> . . . . .	{	The Black-and-Yellow Grosbeak . . . .	89

---



---

## Classification according to Habits

---

Sweet song in air . . . .	The Sky-lark 110
Loud, plaintive <i>pee-you, pee-you</i> . . . . .	{ The Great Himalayan Barbet . 123
<i>Katak-katak-katak</i> . . . . .	
A soft <i>uk-uk-uk</i> . . . . .	{ The Golden throated Barbet . 124
A "shivering" scream . . . . .	{ The Common Indian Swift . . 130
<i>Cuckoo-cuckoo</i> . . . . .	{ The Cuckoo 132
<i>Coo-coo-coo</i> . . . . .	{ The Himalayan Cuckoo . . 133
<i>Pot-you-chick-chick-chick</i> . . . . .	{ The Small Cuckoo . 134
Melodious <i>wherefore-wherefore</i> . . . . .	{ The Indian Cuckoo . 135
A <i>crescendo</i> "brain-fever, brain-fever, BRAIN-FEVER" . . . . .	{ The Large Hawk-Cuckoo . . 136
A high-pitched, wild, metallic call . . . . .	{ The Pied Crested Cuckoo . 137

## *Himalayan and Kashmiri Birds*

- |   |   |  |
|---|---|--|
| 1. A <i>crescendo</i> “ku-il, ku-il,<br>KU-IL” . . . . .                          | } | The Indian<br>Koel . . . 138               |
| 2. <i>Ku-y-o</i> . . . . .  |   |  |
| 3. <i>Kekaree, kekaree, karee,<br/>karee</i> . . . . .                            |   |  |
| Loud screams uttered during<br>flight . . . . .                                   | } | The Paro-<br>quets . 139-141               |
| A soft, low whistle repeated<br>at intervals of thirty<br>seconds . . . . .       | } | The Collar-<br>ed Pigmy<br>Owlet . . . 145 |
| Rather a mournful, queru-<br>lous <i>chee-he-he-he-he-he</i>                      | } | The Com-<br>mon Pariah<br>Kite . . . 154   |
| <i>Kokla-kokla</i> . . . . .  | } | The Kok-<br>la Green<br>Pigeon . . . 157   |
| A soft, subdued <i>cuk-cuk-coo-<br/>coo-coo</i> . . . . .                         | } | The Little<br>Brown<br>Dove . . . 160      |
| <i>Ku-ku</i> . . . <i>ku</i> . . . . .  | } | The Indian<br>Ring Dove 161                |
| <i>Juk-juk-tee-tee-tur.</i> Very<br>high-pitched and rapidly<br>uttered . . . . . | } | The Black<br>Partridge . 166               |
| A loud, shrill “Did-he-do-<br>it? Pity-to-do-it!” . . . . .                       | } | The Red-<br>wattled<br>Lapwing . 170       |

---



---

*Classification according to Habits*

---

Harsh Jay-like cries . . . .		The Magpies 4-6
Do.		The Jays .7 & 8
Do.	{	The Nut-crackers 9 & 10
Do.	{	The Wood-peckers 114-121
Do.	{	The European Roller 125
Chattering cries . . . .	{	The Babblers . 17-25
Shrill Screams . . . .	{	The Himalayan Whistling-Thrush 29
Do.	{	The Fork-tails . 77-79
Do.	{	The Kingfishers 127-128

## II. BIRDS THAT COMMONLY VISIT GARDENS

1. The Indian House Crow (2). (Srinagar and Almora.)
2. The Jackdaw (3). (Srinagar.)
3. The Black-throated Jay (7). (Naini Tal and Mussoorie.)

---

---

## *Himalayan and Kashmiri Birds*

---

4. The Tits (11-16.)
5. The Himalayan Streaked Laughing-Thrush (23).
6. The Black-headed Sibia (30).
7. The Indian White-eye (31).
8. The Red-billed Liothrix (33). (Darjeeling.)
9. The Red-tailed Minla (34). (Darjeeling.)
10. The Red-vented Bulbuls (39 and 40).
11. The White-cheeked Bulbul (41). (Almora and Srinagar.)
12. The Indian Ashy Drongo (44).
13. The Tree-creepers (45 and 46).
14. The Grey-headed Flycatcher-warblers (52 and 53).
15. The Short-billed Minivet (57).
16. The Indian Oriole (59). (Almora and Srinagar.)
17. The Himalayan Starling (60). (Srinagar.)
18. The Common Myna. (Almora, Dharm-sala, Srinagar.)
19. The Ferruginous Flycatcher (65). (Darjeeling.)
20. The White-browed Blue Flycatcher (66).
21. The Verditer Flycatcher (67).
22. The Rufous-bellied Niltava (71).

---

---

*Classification according to Habits*

---

23. The Indian Paradise Flycatcher (73).  
(Almora.)
24. The Dark-grey Bush Chat (76).
25. The Grey-winged Ouzel (85).
26. Tickell's Ouzel (86). (Srinagar.)
27. The Sparrows (95-97).
28. The Eastern Meadow Bunting (99).
29. Sykes's Striated Swallow (102).
30. Hodgson's Yellow-headed Wagtail (105).  
(Srinagar.)
31. The Sunbirds (111 and 112). (Darjeeling.)
32. The Common Indian Swift (130).
33. The Common Pariah Kite (154).
34. The Indian Ring-Dove (161). (Srinagar.)

### III. BIRDS THAT ASSOCIATE IN FLOCKS

1. The Tits (11-16).
2. The Babblers (17-28).
3. The Indian White-eye (31).
4. The Yellow-naped Ixulus (32).
5. The Red-billed Liothrix (33).
6. The Spotted-Wing (35).
7. The White-tailed Nuthatch (42).
8. Minivets (57).

---

## *Himalayan and Kashmiri Birds*

---

9. The Himalayan Starling (60).
10. The Jungle Myna (63).
11. The Paroquets (139-141).
12. The Common Grey Quail (164).
13. The Chukor (165).
14. The Whiskered Tern (171).
15. The Ducks (177 and 178).
16. The Black Bulbul (37).

### IV. BIRDS THAT DWELL BY STREAMS

1. The Himalayan Whistling-Thrush (29).
2. The Fork-tails (77-79).
3. The Redstarts (80-82).
4. The Brown Dipper (88).
5. The Grey Wagtail (104).
6. The Kingfishers (127 and 128).

### V. BIRDS THAT OCCUR ABUNDANTLY ON, OVER, OR ON THE MARGINS OF THE LAKES OF THE KASHMIR VALLEY

1. The White-cheeked Bulbul (41).  
(Makes sallies into the air over the water to catch insects.)



---

---

## *Classification according to Habits*

---

2. The Indian Great Reed-warbler (49).  
(Lives amongst reeds.)
3. The Swallow (100). (Hawks insects over the water and rests on lotus flowers.)
4. Hodgson's Pied Wagtail (103). (Runs about on the floating lotus leaves.)
5. Hodgson's Yellow-headed Wagtail (105).  
(Runs about on the floating lotus leaves.)
6. The European Bee-eater (126). (Hawks insects in the air over the water.)
7. The Kingfishers (127 and 128). (Dive for fish.)
8. Pallas's Fishing-Eagle (153). (A fishing bird of prey.)
9. The Moorhen (167). (Swims like a duck.)
10. The Coot (168). (Swims like a duck.)
11. The Pheasant-tailed Jaçana (169).  
(Swims like a duck, runs about on floating vegetation.)
12. The Whiskered Tern (171). (Flies over the water and catches fish at the surface.)
13. The Cormorants (172 and 173). (Swimmers and divers.)
14. The Herons (174 and 175). (Stand in shallow water and catch fish.)

---

---

## *Himalayan and Kashmiri Birds*

---

15. The Little Bittern (176). (Stands in shallow water and catches fish.)

16. The Ducks (177 and 178).

17. The Indian Little Grebe (179). (Swims like a duck ; frequently dives.)

### VI. BIRDS THAT MOVE ABOUT ON THE TRUNKS OF TREES IN SEARCH OF INSECTS

1. The White-tailed Nuthatch (42).

2. The Tree-creepers (45 and 46).

3. The Woodpeckers (114-121).

4. The Common Wryneck (122).

### VII. BIRDS THAT FIND MUCH OF THEIR FOOD ON THE GROUND

1. The Crows (1 and 2). (Omnivorous and feed anywhere and everywhere.)

2. The Jackdaw (3).

3. The Babblers (17-25).

4. The Red-billed Liothrix (33).

5. The Shrikes (55 and 56). (Descend from perch to ground, seize their quarry, and return to perch to devour it.)

---

---

## *Classification according to Habits*

---

6. The Starlings and Mynas (60, 61, 63).
7. The Chats (74-76).
8. The Magpie-Robin (84).
9. The Blue-headed Rock-Thrush (87).
10. The Black-and-yellow Grosbeak (89).
11. The Finches and Sparrows (90-97).
12. The Buntings (98 and 99).
13. The Wagtails (103-105).
14. The Pipits (106-109).
15. The Skylark (110).
16. The European Roller (125). (Descends from perch to ground, seizes its quarry and returns to perch.)
17. The European Hoopoe (129). (Probes earth with the long bill.)
18. The Vultures (146-150).
19. The Common Pariah Kite (154).
20. The Kestrel (156). (Hovers on rapidly vibrating wings and drops on to its prey.)
21. The Blue Rock-Pigeon (158).
22. The Doves (159-161).
23. The Pheasants (162 and 163).
24. The Common Grey Quail (164).
25. The Partridges (165 and 166).
26. The Red-wattled Lapwing (170).

*See also* : Birds that spend the day by streams.

---

---

*Himalayan and Kashmiri Birds*

---

VIII. BIRDS THAT CAPTURE INSECTS  
ON THE WING

1. The Drongos (43 and 44).
2. The Flycatchers (64-73).
3. The Dark Grey Bush Chat (76)
4. The Swallows (100-102).
5. The European Bee-eater (126).
6. The Swifts (130 and 131).

IX. BIRDS THAT SPEND GREAT PART  
OF THE DAY ON THE WING

1. The Swallows (100-102).
2. The Skylark (110). (Soars and sings in the air.)
3. The Himalayan Pied Kingfisher (127). (This bird quarters the waters for its quarry, often hovering Kestrel-like.)
4. The Swifts (130 and 131).
5. The Vultures (146-150). (Sail for hours together on outstretched wings.)
6. The Common Pariah Kite (154). (Sails in circles.)
7. The Kestrel (156). (Hovers on rapidly moving wings.)
8. The Whiskered Tern (176).

---

---

*PART II*

*DESCRIPTIVE LIST OF BIRDS*

---



---

---

# DESCRIPTIVE LIST OF BIRDS

---

## EXPLANATION OF ABBREVIATIONS

F. STANDS for the *Fauna of British India*, and the number which follows indicates the number of the bird in the Fauna. Thus (F. 63) against a bird indicates that it is the species bearing the number 63 in the bird volumes of the *Fauna of British India* (First Edition).

J. stands for Jerdon's *Birds of India*, and the number that follows indicates the number of the bird in Jerdon's work.

The above two works contain descriptions far more detailed than those given in this little book ; the reader may wish in some cases to refer to these volumes to check his identification of a bird.

I means the bird described is 6 inches in length, or the size of a sparrow.

---

## *Himalayan and Kashmiri Birds*

---

—I means that the bird is smaller than a sparrow.

+I means that the bird is slightly larger than a sparrow.

II means that the bird is  $8\frac{1}{2}$  inches in length, or the size of a bulbul.

—II means that the bird is slightly smaller than a bulbul.

+II means that the bird is slightly larger than a bulbul.

III means that the bird is 10 inches long, or the size of a myna.

+III and —III, that it is slightly larger or smaller.

IV means that the bird is  $17\frac{1}{2}$  inches long, or the size of a house-crow.

+IV and —IV mean that it is a little larger or smaller.

V means that the bird is 23 inches in length, or the size of a kite.

+V and —V mean that it is a little smaller or larger.

N.B.—Ornithologists usually measure birds from the tip of the bill to the tip of the tail; hence a small bird with a long tail



---

---

## *Descriptive List of Birds*

---

will measure as much as a larger one with a short tail.

While I shall adopt the usual measurement, I shall, in the case of a bird having a long tail, measure only part of the tail, i.e. treat the bird as though it had an average tail, and add the words, "bird has long tail."

E. means Eastern Himalayas, i.e. Nepal and eastwards.

W. means Western Himalayas, i.e. West of Nepal.

K. means Kashmir.

To illustrate: the entry

"*Corvus macrorhynchus*: The Jungle Crow or Corby. F. 4, J. 660, + IV. E. W. K." means that the number of the species in the *Fauna of British India* is 4, and in Jerdon's *Birds of India* 660. That the species is rather larger than the house-crow and that it is found in the Eastern, the Western Himalayas, and Kashmir.

---

---

## *Himalayan and Kashmiri Birds*

---

### *The Crow Family*

1. *Corvus macrorhynchus* : The Jungle Crow or Indian Corby. F. 4, J. 660, +IV. E. W. K. Glossy black all over.

The common crow of all the hill stations, except Almora and Srinagar, where it is largely replaced by *Corvus splendens* (the House Crow).

2. *Corvus splendens* : The Indian House Crow. F. 7, J. 663, IV. E. W. K.

A glossy black bird with the neck, upper back and breast grey. Thus it is not unlike a jackdaw (q.v.) in appearance.

It is the commonest crow of Almora and Srinagar. It is occasionally seen at Simla. I have not observed it at any other hill station.

3. *Corvus monedula* : The Jackdaw. F. 9, J. 665, +III. K.

A glossy black bird with grey nape and neck. This is rather like the house-crow in appearance, but may be easily distinguished by

(1) Its short, sharp *caw*.

(2) The grey of the neck is paler and extends up the nape.

(3) The white eye.

(4) Its much smaller size.

---

---

## *Descriptive List of Birds*

---

This bird does not occur in the Himalayas, but is a permanent resident in Kashmir. It is abundant at Srinagar.

### *The Magpies*

4. *Urocissa occipitalis*: The Red-billed Blue Magpie. F. 12, J. 671, +III, with a tail nearly 19 inches long. W.

Head, neck and breast black. Conspicuous white patch on nape and neck. Rest of upper plumage is purplish-blue. Tail blue tipped with white; the middle pair of tail feathers has a black band next to the white tip. Wings brown, blue and white. Lower parts whitish. Bill and legs bright red.

This fine bird is common at Naini Tal, but does not occur at Murree, Dharmsala, or Darjeeling. I have seen it at Mussoorie and at Annandale (Simla).

5. *Urocissa flavirostris*: The Yellow-billed Blue Magpie. F. 13, J. 672, +III, with a tail nearly 18 inches long. E. W. K.

Very like the last species in colouring, but readily distinguished by the yellow bill and the much smaller nuchal patch of white feathers. The blue of the upper parts is greyer than in the red-billed species.

---

---

## *Himalayan and Kashmiri Birds*

---

This bird is less often seen than the red-billed species in the western hill stations, as it seems to keep to higher elevations in summer. About Simla it begins to get abundant at 8000 feet ; but at Darjeeling it occurs from 6000 feet upwards. Occurs in Kashmir in hills and valley.

6. *Dendrocitta himalayensis*. The Himalayan Tree-pie. F. 18, J. 676, -III, with tail 9 inches long. E. W.

Forehead black ; crown of head, nape and back of neck dark ashy-grey. Sides of head, chin, throat and upper breast brownish-black. Belly grey ; a patch of chestnut under the tail. Back and shoulders reddish-brown. Wings black, with a white bar. Tail black and grey.

This may be described as a small and dingy edition of the tree-pie of the plains.

This species is less common at most hill stations than the blue magpie.

### *Jays*

7. *Garrulus lanceolatus* : The Black-throated Jay. F. 24, J. 670, +III. W. K.

The crested head is black ; the remainder of upper plumage is fawn-coloured. The feathers of the chin and throat are black with white

---

---

## *Descriptive List of Birds*

---

shafts. Tail jay-blue with narrow black cross-bars; the tip is white, but there is a broad black subterminal bar. The wings are coloured blue, fawn, black, white and grey.

Not common at Dharmsala; fairly so at Murree and Simla, and abundant at Naini Tal and Mussoorie, where it seeks its food in gardens and among outhouses. Sometimes it goes about in flocks. The flight is somewhat heavy, like that of a crow, but the flaps of the wings are more rapid. Its call is harsh.

8. *Garrulus bispecularis*: The Himalayan Jay. F. 26, J. 669, +III. E. W. K.

A fawn-coloured bird with a conspicuous black moustache. Tail black. Wings jay-blue, fawn, grey, black and white.

Often goes about in flocks. A shier bird than the black-throated jay. Harsh voice.

Rare at Dharmsala and Darjeeling. Common at Naini Tal; common at Simla in winter.

### *Nutcrackers*

These members of the crow family are large arboreal birds, which feed largely on seeds of pine and cedar trees. They emit harsh, jay-like notes.

---

---

## *Himalayan and Kashmiri Birds*

---

9. *Nucifraga hemispila*. The Himalayan Nutcracker. F. 27, J. 666, —IV. W.

Plumage brown, heavily spotted and streaked with white. Wings black with white bars.

Rare at Dharmsala; not very common at Simla and Naini Tal. It is rarely seen in gardens, being a forest bird.

10. *Nucifraga multipunctata*: The Larger Spotted Nutcracker. F. 28, J. 667, —IV. W. K.

Head and hind neck chocolate-brown; rest of body plumage dark chocolate-brown with many large white spots; these spots occupy more space than the brown on the lower plumage. Wings black, spotted with white. Tail black, tipped with white.

During flight the body appears mainly white and the wings black. The cry is harsh, like that of the magpie.

This fine bird is not often seen below 8000 feet in summer. It is fairly common in the fir forests of Gulmarg.

### *Tits*

Small birds which usually go about in flocks. Although unlike crows in appearance and size, they show their affinity to the corvidae by the

---

## *Descriptive List of Birds*

---

way they have of holding their quarry under their claws.

11. *Parus atriceps*. The Indian Grey Tit. F. 31, J. 645, —I. E. W. K.

Head, neck and breast black, except cheeks, which are white. Back grey, tinged with blue near the tail. Wings and tail black, white and grey. A broad black band runs down the middle of the white abdomen.

Common at Dharmsala, Almora, and in the Kashmir valley. In most hill stations it is not seen, as about 6000 feet elevation is ordinarily the highest limit of its range.

Less gregarious than most tits.

12. *Parus monticola*. The Green-backed Tit. F. 34, J. 644, —I. E. W. K.

Head, neck and breast black, with the cheeks white. A black band runs down the middle of the deep-yellow abdomen. Back and shoulders greenish-yellow. Wings and tail black, slaty-blue and white.

The commonest tit of all the hill stations from Murree to Darjeeling. It occurs at Gulmarg, but is not very abundant there.

13. *Ægithaliscus erythrocephalus*. The Red-headed Tit. F. 35, J. 634, —I. E. W.

A midget, only 4 inches long.

---

---

## Himalayan and Kashmiri Birds

---

Top of head and nape chestnut. Chin, eyebrow and moustache white. Sides of head and round patch on the throat black. Upper plumage blue-grey with brown in the wings and brown and white in the tail. Lower plumage is rust-coloured. This species lacks the black abdominal band which is so conspicuous in most tits.

Very common at every hill station (except Almora) from Dharmsala to Darjeeling. Usually seen in flocks.

14. *Machlolophus xanthogenys*: The Yellow-cheeked Tit. F. 42, J. 647, —I. W.

Has rather a long pointed crest. Head and neck black, except the eyebrow, a patch on the nape, the cheeks and the tip of the crest, which are bright yellow. Back, wings and tail are olive-green, brown, black and white. Lower parts yellow, except for a broad black band running down the middle of the breast and abdomen.

Occurs from Murree to Naini Tal. Not so common as the red-headed and green-backed tits.

15. *Lophophanes melanolophus*: The Crested Black Tit. F. 44, J. 638, —I. W. K.

A tiny bird. Head, crest and breast black,



---

---

## *Descriptive List of Birds*

---

except for a patch on the nape, the sides of the head and a band down the side of the neck, which are white. Rest of the plumage is iron-grey, suffused with chestnut on the sides of the body and under the tail. A row of white spots and a little chestnut on the wings.

Common at Gulmarg and fairly so at Murree, Dharmsala, Simla, and Naini Tal.

16. *Lophophanes dichrous* : The Brown Crested Tit. F. 49, J. 637, —I. W. K.

A small bird. Brownish-grey above, yellowish below. Cream-coloured band on each side of neck. It might well pass for a young crested black tit.

Usually keeps to rather higher elevations than the hill stations. I have, however, seen it on several occasions at Naini Tal.

### *Babblers*

Babblers are noisy birds which usually go about in small flocks seeking for their insect quarry among the foliage and fallen leaves. The "Seven Sisters" of the plains belong to this family.

17. *Garrulax leucocephalus* : The Himalayan White-crested Laughing Thrush. F. 69, J. 407, +III. E. W.

---

## *Himalayan and Kashmiri Birds*

---

Crest, head, throat, breast, and neck snowy-white, except for a broad black streak running through the eye and an ashy band on the nape. Remainder of plumage brown.

Goes about in flocks of about twenty uttering loud and discordant cries.

Not found west of Garhwal. In the Eastern Himalayas it is not often seen higher than 6000 feet.

18. *Garrulax albigularis* : The White-throated Laughing-Thrush. F. 76, J. 411, +III. E. W.

Upper plumage olive-brown, lower rust-coloured. Cheeks, chin and throat white. Feathers round eye black. Some of the tail feathers have white tips.

Goes about in noisy flocks in the woods. It has a number of cries; one is plaintive, a second a trill, a third a squeak.

Abundant at Mussoorie, fairly common at Naini Tal and Simla, rare at Darjeeling. Not found at Dharmsala or Murree.

19. *Ianthocincla ruficularis* : The Rufous-chinned Laughing-Thrush. F. 80, J. 421, -III E. W. K.

Forehead and crown black. Sides of head black and white, with a red patch on each

---

---

## *Descriptive List of Birds*

---

cheek. Grey ring round eye. Rest of upper plumage olive-brown. Narrow white bar near edge of wing. Chin and upper throat red; lower throat white. Rest of lower parts ashy-brown. Gregarious. Mussoorie is the only hill station at which I have found it common. It occurs in the vicinity of Darjeeling and in the dense woods of the lower hills of Kashmir.

20. *Trochalopterym erythrocephalum*: The Red-headed Laughing-Thrush. F. 82, J. 415, +III. W. Top of head chestnut. Lower part of cheeks, chin and upper throat dark chocolate—almost black. Rest of plumage olive-brown, with some bright golden-yellow in the wing.

This gregarious bird has a curious plaintive call, which sounds like *poor-wee-wee*. The second *wee* is very high-pitched.

In summer this bird is not often seen below 7000 feet. I have noticed it on Lariakanta at Naini Tal. It occurs in the higher woods near Simla and Dharmsala. A great skulker.

21. *Trochalopterym nigrimentum*: The Western Yellow-winged Laughing-Thrush. F. 85, J. 416, III. E.

Feathers of forehead reddish with black shafts; those of rest of head grey with black

---

---

## *Himalayan and Kashmiri Birds*

---

shafts. Nape bright chestnut. Back and shoulders reddish, spotted with black. Rump olive-green. Tail grey, suffused with golden-yellow. There is much golden-yellow in the wings. Sides of head are black and grey. Chin and throat black with reddish mottlings.

This is one of the commonest babblers at Darjeeling. It often feeds on the paths, diving into the nearest thicket at the approach of man.

22. *Trochalopteryum variegatum* : The Eastern Variegated Laughing-Thrush. F. 90, J. 418, +III. W.

An olive-brown bird with *much* golden-yellow in the wings. It may be distinguished from the two species described above by the facts that : (1) It has no red on the head. The crown and nape are greyish-brown. (2) Tail feathers are tipped with white. Sides of head black and white. Chin and upper throat black.

In summer it is not found below 7000 feet. It nests at Simla and is abundant at Dharm-sala. A noisy, gregarious species.

23. *Trochalopteryum lineatum*. The Himalayan Streaked Laughing-Thrush. F. 99, J. 425, II. W. K.

---

---

## *Descriptive List of Birds*

---

Forehead, crown, nape and shoulders ashy brown, streaked with black. Lower back reddish-brown, with white streaks. Cheeks, chin, throat and breast chestnut. Wings and tail reddish-brown. The tail has a black band across it near the tip. The tip of the tail is grey. A greyish-white ring round the eye.

This is the common babbler of the western hill stations, where it takes the place of the "seven sisters" of the plains. They often visit gardens. They sometimes nest in gardens. Occasionally two or three of them will hold squeaking competitions. Their movements are jerky, like those of the "seven sisters."

24. *Grammatoptila striata*: The Striated Laughing-Thrush. F. 101, J. 382, +III. E. W.

The plumage of this fine bird is mainly umber-brown, each feather having a bold, white, median streak. The tail is chestnut and there is some chestnut in the wing.

Not often seen below 7000 feet in the Western Himalayas, but at Darjeeling it occurs from 6000 feet upwards. As Jerdon remarks, it is more often heard than seen. One of its calls resembles the clucking of a hen that has just laid an egg.

---

---

## *Himalayan and Kashmiri Birds*

---

25. *Pomatorhinus erythrogeus*: The Rusty-cheeked Scimitar Babbler. F. 129, J. 405, +III. E. W.

Bill  $1\frac{1}{2}$  inches long, slender and curved like a scimitar. Upper plumage olive-brown. Fore-head and sides of head rust-coloured. Chin, throat, and lower parts streaked white and grey. Rusty patch under the tail.

Goes about in pairs. Call a loud whistle.

Fairly common at Naini Tal, Simla, and Darjeeling, but not, I believe, at other hill stations.

26. *Alcippe nepalensis*: The Nepal Babbler. F. 163, J. 388, —I. E.

A tiny grey-brown bird with a conspicuous ring of white feathers round the eye. Its sober colouring led Jerdon to call it the Quaker Thrush. It occurs in small parties in dense jungle. The call is a low twitter.

27. *Stachyrhis nigriceps*: The Black-throated Babbler. F. 169, J. 391, —I. E.

Head black, streaked with white. Cheeks white. Rest of upper plumage olive-brown, washed with red on wings and tail. Lower plumage bright-reddish yellow.

A tiny bird which goes about in small parties in the upper foliage.

---

---

## *Descriptive List of Birds*

---

28. *Stachyridopsis ruficeps*: The Red-headed Babbler. F. 174, J. 393, —I. E.

Head chestnut. Rest of upper plumage olive-green. Lower plumage white, tinted with red.

Its note is said to be "clear and musical, like the ringing of a little bell."

Common at Darjeeling in parties in tall trees.

29. *Myiophoneus temmincki*: The Himalayan Whistling-Thrush. F. 187, J. 343, +III. E. W. K.

From a little distance it looks like a cock blackbird; the yellow bill adding to the similitude.

There is, however, a good deal of cobalt blue in the black plumage, only noticeable in the sunlight.

One of the commonest birds of the streams of the Himalayas and Kashmir. It likes to perch on boulders in mountain streams, by the side of which it nests. It feeds on insects, which it catches on boulders. It has a loud and pleasant whistling call. When disturbed it flies off, uttering a shrill cry. In this respect it resembles other birds which frequent mountain streams.

---

---

## *Himalayan and Kashmiri Birds*

---

30. *Lioptila capistrata*: The Black-headed Sibia. F. 204, J. 429, II. E. W.

A bulbul-like bird with a short crest. Head and crest black. A reddish collar. Rest of upper plumage greyish-brown, the wings and tail having some chestnut in them. A white bar in the wing. Lower plumage rust-coloured. The tail is rather long, the middle feathers being the longest and the outer ones the shortest.

It usually keeps to the upper parts of trees. Sometimes it attaches itself to a wandering party of tits. Its call is so loud, high-pitched, and penetrating, that it cannot fail to attract notice. The call resembles one of the louder cries of the streaked laughing-thrush. Hutton syllabises the call: *titteree, titteree, tweeyo*.

Common at Simla, Naini Tal, Mussoorie, and Darjeeling.

31. *Zosterops palpebrosa*: The Indian White-eye. F. 226, J. 631, —I. E. W. K.

Known also as the girdle-eye and spectacle-bird, because of a conspicuous ring of white feathers round the eye. A tiny bird, that goes about in little flocks among the foliage. Upper plumage golden-yellow. Wings and tail greenish-brown. Chin and throat bright yellow, rest of lower plumage grey.



---

---

## *Descriptive List of Birds*

---

It utters a loud cheeping note—at breeding season, in spring, it sings a sweet little lay.

Common at Murree, Dharmsala, Simla, Naini Tal, Almora, also in the woods of the lesser ranges of the Kashmir hills; but in the Eastern Himalayas does not ascend as high as the hill stations.

32. *Ixulus flavicollis*: The Yellow-naped Ixulus. F. 232, J. 623, —I. E. W.

Forehead and crest rich brown. Nape and sides of head grey-brown. Black moustache. White ring round eye. Chestnut collar round hind neck, broad in the middle. Back olive-brown. Wings and tail brown suffused with olive. Chin and throat white. Rest of lower plumage pale yellow.

Goes about in twittering flocks seeking for insects among the foliage.

Not common in the Western Himalayas but abundant at Darjeeling.

33. *Liothrix lutea*: The Red-billed Liothrix. F. 235, J. 614, I. E. W.

The Pekin Robin of bird-fanciers. Upper plumage olive-green. Forehead and crown tinged with yellow. Yellow ring round eye. Bill coral red. Chin and upper throat bright yellow, lower throat rich orange. Rest of

---

---

## *Himalayan and Kashmiri Birds*

---

lower plumage yellow. Many of the wing feathers are edged with yellow and, in the cock, with crimson also. A very lively sprightly bird. The ordinary call is *tee-tee-tee-tee-tee*. In the spring the cock sings beautifully. Gregarious.

One of the commonest birds of Darjeeling, but in the Western Himalayas it does not usually ascend as high as the hill stations. It has been seen near Kausauli and is common at Bhim Tal below Naini Tal.

34. *Minla igneitinca* : The Red-tailed Minla. F. 258, J. 618, —I. E.

Head black with broad white eyebrows, which almost meet at the back of the head. Back yellowish-green suffused with dull red. Wings and tail black and crimson. Throat white, rest of lower plumage yellow. The hen is duller and her underparts are white.

This species is often seen at Darjeeling along with parties of tits.

35. *Cephalopyrus flammiceps* : The Fire-cap. F. 260, J. 633, —I. W.

A midget 4 inches long.

*Cock* : Upper parts olive-green. Throat and breast golden-yellow ; rest of lower parts dull yellow ; a cap of flaming red. Chin suffused

---

---

## *Descriptive List of Birds*

---

with bright red. Wings and tail brown edged with yellow.

*Hen*: Has no red on head or chin and the yellow parts of her plumage are duller and tinged with green.

36. *Psaroglossa spiloptera*: The Spotted-wing. F. 261, J. 691, —II. W.

The classification of this bird has puzzled ornithologists. Jerdon classes it with the starlings; but the hen is not coloured like the cock as is the case with the starlings.

*Cock*: The ground colour of the head and back is grey, each feather being edged with black on the head and shoulders and brown on the back, so that the bird has markings like the scales of a fish. Rump reddish-brown. Chin and throat deep chestnut maroon. Rest of lower parts reddish brown. The wings are black glossed with green or blue and spotted white, like the plumage of a starling.

*Hen*: Upper plumage brown with grey spots; lower parts brown with cream edges to the feathers. Wings dark brown with some grey in them.

Not found west of Dharmsala. Not found as high as Simla in summer, but occurs at

---

---

## *Himalayan and Kashmiri Birds*

---

Naini Tal. A flock of half a dozen or more birds having a starling-like appearance, which twitter like starlings and keep to the topmost branches of trees, from which they make sallies into the air, may be set down as a company of spotted-wings.

### *Bulbuls*

Bulbuls are larger than sparrows. The sexes are alike. The typical ones have the head crested and a patch of red or yellow under the tail. They frequent gardens and have cheerful, tinkling calls.

37. *Hypsipetes psaroides*: The Himalayan Black Bulbul F. 269, J. 444, II, with tail  $4\frac{1}{2}$  inches long. E. W. K.

Head and untidy loose crest black; cheeks grey; bill and legs coral red. Upper plumage dark grey; wings and tail dark brown, edged with grey. Lower parts grey, pale on the abdomen. Seen from a distance this bird looks black, hence its popular name. A party of dull black-looking birds, with red bills and legs, which keeps to the tops of trees and chatters harshly may be set down with certainty as black bulbuls.

These birds are common at the various

---

---

## *Descriptive List of Birds*

---

western hill stations ; but at Darjeeling are rarely seen at elevations above 5000 feet.

Found in the dense jungles of the lower hills and in the valley of Kashmir.

38. *Alcurus striatus* : The Striated Green Bulbul. F. 277, J. 449, II. E.

Crested head and upper plumage olive-green, streaked with yellow. Wings and tail brown with olive green and yellow markings. Cheeks brown with creamy streaks. Chin bright yellow. Throat yellow spotted with black. Sides of neck and breast black, streaked with yellow. Yellow patch under tail. Rest of lower plumage yellow in middle and olive-brown streaked with yellow at sides.

Goes about in parties which keep to tree-tops and utter frequently pleasing notes.

Common at Darjeeling.

39. *Molpastes bengalensis* : The Bengal Red-vented Bulbul. F. 282, J. 461, II. E.

This is the bulbul which is so common in the plains. The short crest, head, neck and breast are black. The body plumage is brown, each feather having a narrow cream margin which gives the bird a scaly appearance. Wings and tail brown, the tail being narrowly tipped with white. There is a small patch

---

## *Himalayan and Kashmiri Birds*

---

of white just above the tail. A conspicuous crimson patch under the tail.

Common at Darjeeling.

40. *Molpastes intermedius* : The Punjab Red-vented Bulbul. F. 283, J. 283, II. W.

This is the western form of the last-named species. It is merely a race and not a separate species. It is very like the Bengal race in appearance, the chief difference being that the cheeks are chocolate-brown instead of black.

It is a common bird at Almora, but, as it rarely ascends the hills higher than 5500 feet, it is only seen at the lowest parts of most hill stations.

41. *Molpastes leucogenys* : The White-cheeked Bulbul. F. 284, J. 458, II. E. W. K.

In shape and colouring this is not unlike the two species described above. It differs, however, in the crest being longer and ending in a forwardly directed point. It has white cheeks and the patch of feathers under the tail is bright sulphur-yellow instead of crimson.

It is very common at Almora and in the Kashmir valley. At Srinagar it hawks insects over the backwaters and canals just as fly-catchers do.

As 7000 feet is about as high as it is found,

---

---

## *Descriptive List of Birds*

---

it is not common at most hill stations, but may occasionally be seen at Dharmsala, Simla and other western stations. In the Eastern Himalayas it does not seem to ascend higher than 5000 feet.

### *Nuthatches*

Small, short-tailed climbing birds which often go about in small flocks searching trunks and branches of trees for insects.

42. *Sitta himalayensis*: The White-tailed Nuthatch. F. 315, J. 284, —I. E. W.

Upper parts slaty-blue. Forehead and a broad line running down the sides of the head and neck black. There is some white in the tail. Under parts chestnut.

The hen is duller than the cock.

The common nuthatch of Naini Tal and Darjeeling. Rare at Simla and Dharmsala. Does not occur west of Kangra.

### *Drongos*

Drongos are active, long-tailed birds which have many of the habits of flycatchers. Their plumage is largely black and both sexes are arrayed alike.

43. *Dicrurus ater*: The Black Drongo, or

---

---

## *Himalayan and Kashmiri Birds*

---

King Crow. F. 327, J. 278, II, with a tail 6 inches long. W. E.

A glossy jet-black bird with a forked tail.

It is fond of sitting on telegraph wires and on the backs of cattle.

It is very common everywhere in the plains of India. It is common at Almora and on the Kashmir Road. It is fairly abundant at Dharmsala and Mussoorie and has been seen at Naini Tal. In the vicinity of Simla it has not been observed at elevations of over 5000 feet, and in the Eastern Himalayas it does not ascend as high as this.

44. *Dicrurus longicaudatus*: The Indian Ashy Drongo. F. 328, J. 280, II, with a tail  $6\frac{1}{2}$  inches long. E. W. K.

Upper plumage metallic dark indigo, lower plumage dark grey. Very like the King Crow in habits and appearance; but its cry is less harsh.

This is the commonest drongo of most hill stations, but at Dharmsala its place is taken by the King Crow, this species living at altitudes of 9000 feet in summer at that locality.

It is common in the Kashmir valley and at Murree, Simla, Mussoorie and Darjeeling.



---

---

## *Descriptive List of Birds*

---

### *Creepers*

Creepers are small brown birds with rather long bills. They feed on insects, which they catch in the same way as nuthatches do, but their hunting grounds are not confined to trees; they seek their quarry on rocks and walls.

45. *Certhia himalayana*: The Himalayan Tree-creeper. F. 341, J. 243, I. W. K.

Upper plumage dark brown, heavily streaked with yellowish-brown. Short, broad, pale eyebrow. Rump, wings and tail reddish-brown, streaked with yellowish-brown on the wings and with black cross-bars on the tail. Chin and throat white.

Usually go about in pairs. Run about with great address on vertical surfaces, using the stiff tail feathers as supports after the manner of woodpeckers. The call is a shrill *pee-pee*.

Common at Gulmarg and the West Himalayan hill stations.

46. *Certhia discolor*: The Sikkim Tree-creeper. F. 344, J. 245, I. E.

Like the last species in appearance and habits, but the chin and throat are earthy-

---

---

## *Himalayan and Kashmiri Birds*

---

brown instead of white and there are no black cross-bars on the tail.

An eastern race of the Himalayan tree-creeper. Common at Darjeeling.

### *Wrens*

Wrens are mottled-brown, short-tailed mid-gets which creep about in brushwood.

47. *Anorthura neglecta*: The Kashmir Wren. F. 352, (Not recorded by Jerdon.) —I. K. W.

Less than 4 inches long. Plumage reddish-brown; the lower back, wings and short tail being closely cross-banded with black.

Common at Gulmarg and Dharmsala. At Simla and other stations it lives at high elevations in summer.

48. *Pnæpyga squamata*: The Scaly-breasted Wren. F. 356, J. 329, —I. E. W.

Upper plumage rich brown, dotted with buff. Lower plumage in cock white, dotted and scaled with black; in hen ruddy, dotted and scaled with black.

The tail of this tiny bird is so short as to be almost invisible.

In the western hill stations it is seen only in winter. At Darjeeling it occurs in summer.

---

---

## *Descriptive List of Birds*

---

### *Warblers*

A very large family of "tiny brownie birds," of which nearly one hundred members occur in India. As these birds are all small and have very little distinctive colouring, they are the most difficult birds to identify.

49. *Acrocephalus stentoreus*: The Indian Great Reed-warbler. F. 363, J. 515, —II. K.

A giant among warblers. An olive-brown bird with a cream-coloured eyebrow. Chin and throat nearly white, the throat sometimes has brown streaks; rest of lower plumage creamy.

This bird lives among reeds in which it moves about with much address.

Fairly common on the margins of the Kashmir lakes.

50. *Sylvia affinis*: The Indian Lesser White-throated Warbler. F. 402, J. 583, I. K.

Upper plumage earthy-brown, tinged with grey on head and nape. Lower plumage pale grey. Tail feathers tipped with white.

Occurs in the Kashmir valley in open country near streams.

51. *Phylloscopus humii*: Hume's Willow-

---

---

## *Himalayan and Kashmiri Birds*

---

warbler. F. 418 (not described by Jerdon),  
—I. W. K.

A midget 4 inches long. Upper plumage light olive-green, with a faint grey patch on head and a creamy eyebrow. A yellowish bar in the wing. Sides of head and lower plumage faint buff colour.

This bird is said to leave Simla in the middle of May, but it is found at Dharmsala and in the Kashmir valley throughout the summer.

52. *Cryptolopha xanthoschista*: Hodgson's Grey-headed Flycatcher-warbler. F. 434, J. 573, —I. W.

A small bird, just over 4 inches long. Head and neck grey; a broad band of pale grey runs from the bill over the head to the nape; this band is more distinct in some individuals than in others. Eyebrow white. Rest of upper plumage olive-green. Lower plumage bright yellow. Wings brown, the feathers having yellowish-green margins. Tail brown.

The commonest warbler at all the West Himalayan hill stations, including Naini Tal and Almora. Its loud call of four or five high-pitched, not very musical notes, emanates from every tree in spring. It nests in gardens.

---

---

## *Descriptive List of Birds*

---

53. *Cryptolopha jerdoni*: Brooks's Grey-headed Flycatcher-warbler. F. 435, J. 572, —I. E.

This is the eastern form of Hodgson's species. It differs from this in that the head, except for the median grey band, is dark grey, almost black, and the rest of the upper plumage is greyish-brown instead of olive-green.

This species occurs at Darjeeling.

54. *Suya crinigera*: The Brown Hill-warbler. F. 458, J. 547, —II. E. W. K.

There is nothing distinctive about the colouring of this bird. The upper plumage is dark brown, each feather having an olive-green margin. Cheeks dark brown. Lower parts pale ruddy yellow, with faint brown markings on the breast.

It seeks its food on the ground, but sometimes perches on the summit of long grass or other point of vantage and utters a rasping call, like the noise made by a saw.

Fairly common at Simla, but rare in the Eastern Himalayas. It is said to be fairly common on the lower ranges of Kashmir. I have not had the opportunity of testing this statement.

---

---

## *Himalayan and Kashmiri Birds*

---

### *Shrikes, or Butcher Birds*

Shrikes form a well-marked family of birds. When perched, they sit upright with the tail pointing downwards. Their habit is to sit exposed on a perch and from there pounce on some insect on the ground. Many species have a broad black band running backwards from the beak through the eye.

Shrikes are common birds in the plains, but do not ascend the Himalayas to any great altitude, they are therefore rarely seen at any hill station. One species, however, is very abundant in the Kashmir valley.

55. *Lanius erythronotus* : The Rufous-backed Shrike. F. 476, J. 257, — III. E. W. K.

Head, neck and upper back pale grey, with a broad black band running across the forehead and through the eyes. Shoulders and lower back rusty red. Chin, throat and upper breast white. Wings and tail black, some of the feathers having reddish edges. White wing-bar.

Very common in the Kashmir valley. Dozens may be seen on the journey from Baramula to Srinagar. They perch in the poplars. It breeds as high as 6000 feet near Simla. Jones says it is a dupe of the cuckoo. In most parts

---

---

## *Descriptive List of Birds*

---

of the Himalayas this species does not ascend above 5000 feet. It occurs at Almora.

56. *Lanius tephronotus*: The Grey-backed Shrike. F. 477, J. 258, II. E. W. K.

Very like the rufous-backed shrike in appearance. It differs from it in that (1) the grey of the head extends more than half-way down the back, and the shoulders are grey; (2) the red of the lower back is chestnut rather than rusty; (3) it has *no* white wing-bar.

This species is common in the Kashmir valley but not in the hill stations, at any rate in summer.

### *Minivets*

Little birds that go about in flocks and keep to the tops of trees. The cocks are usually bright red and black and the hens vivid yellow and black.

57. *Pericrocotus brevirostris*: The Short-billed Minivet. F. 495, J. 273, —I, but with a tail 4 inches long. E. W.

*Cock*: Head and shoulders glossy black. Back and lower parts crimson. Wings black with a crimson band running along (not across) each. Middle tail feathers black, the others crimson.

---

---

## *Himalayan and Kashmiri Birds*

---

*Hen*: Crown of head, neck and shoulders greyish-green. Forehead, remainder of upper plumage, and lower plumage bright yellow. Wings brown with yellow band. Tail black and yellow.

Common at Dharmsala and Darjeeling. Fairly common at other hill stations.

### *Cuckoo-Shrikes*

Cuckoo-Shrikes have the habits of minivets: but they lack their bright colouring.

58. *Campophaga melanoschista*: The Dark Grey Cuckoo-Shrike. F. 505, J. 269, II. E. W.

A dark grey bird having wings and tail black. The latter tipped with white.

In the Western Himalayas this species is not often seen above 5000 feet; but it goes higher in the eastern part of the range and occurs at Darjeeling.

### *Orioles*

Orioles are birds of brilliant plumage, nearly as large as mynas. They have melodious calls.

59. *Oriolus kundoo*: The Indian Oriole, or Mango Bird. F. 518, J. 470, —III. W. K.

*Cock*: Plumage mainly bright yellow. Some black on the sides of the head and in the wings



---

---

## *Descriptive List of Birds*

---

and tail. During flight this bird looks golden-yellow. Bill pink. Eyes red.

*Hen*: The yellow of the plumage is duller and the back is tinged with green.

Keeps to leafy trees. Call a soft, rich, mellow *peebo*.

Common at Almora and in the Kashmir valley. As it does not commonly ascend above 5000 feet, it is not likely to be seen at other hill stations.

### *Starlings*

60. *Sturnus humii*: The Himalayan Starling. F. 529, J. 682, II. K.

Very like the English starling in appearance. A glossy black bird, with yellow specks and spots. The sheen on the plumage looks purple, green or red according to the incidence of sunlight on the plumage. Abundant in the Kashmir valley, equally at home in grass land or among the willows in which the bird nests.

61. *Acridotheres tristis*: The Common Myna. F. 549, J. 684, III. K. W.

Head, neck and upper breast black; rest of body plumage rich brown; wings black, showing conspicuous white bar during flight.

---

## *Himalayan and Kashmiri Birds*

---

Tail feathers black, all except the middle pair tipped with white : thus the white is visible only during flight.

Bill and legs bright yellow ; bright yellow patch of skin behind the eye.

This is the common myna of the plains. It is also the common myna of the Kashmir valley, Simla, Dharmsala and Almora. At the other hill stations it is largely, if not entirely, replaced by the Jungle Myna. Brooks observed it at Naini Tal in 1869 ; I do not remember seeing it there.

62. *Sturnia malabarica* : The Grey-headed Myna. F. 538, J. 688, II. W.

Forehead pale grey, almost white in some specimens. Remainder of upper plumage dark grey, tinged with reddish-yellow on the lower back ; the feathers of the head and neck have white shafts, giving this part of the plumage a streaked appearance. Wings black and grey. Chin almost white ; rest of lower parts reddish-grey, the red being deepest towards the tail. Many of the feathers have white shafts.

This bird keeps to the tops of trees. Common at Almora ; but I have not seen it at other hill stations.

---

---

## *Descriptive List of Birds*

---

63. *Æthiopsar fuscus*: The Jungle Myna. F. 552, J. 686, III. E. W.

Very like the common myna and is doubtless often mistaken for it.

It may, however, be readily distinguished by the fact that there is no yellow patch of bare skin behind the eye. The entire head is black. There is a little tuft of black feathers on the forehead which the common myna has not. The base of the bill is blue-black.

Common at Mussoorie, rare at Simla. At Dharmsala and, I think, Darjeeling it does not ascend the mountains so high as the hill stations. Brooks observed it at Ramgarh, between Naini Tal and Almora.

### *Flycatchers*

Flycatchers are well-represented in the Himalayas. They feed exclusively on insects which they catch on the wing. Their habit is to make sallies into the air from a perch in order to capture their quarry. All flycatchers have this habit, but some birds which are not flycatchers behave in this way, e.g. bee-eaters, drongos.

64. *Hemichelidon sibirica*: The Sooty flycatcher. F. 558, J. 296, —I. E. W. K.

---

## *Himalayan and Kashmiri Birds*

---

Upper plumage brown, with dark spots on the head. Ring of white feathers round eye. Tail brown. Lower plumage smoky-brown with greyish patches. Small white patch on throat, indistinct in some specimens.

Common at Darjeeling and Dharmsala, but rare at intermediate hill stations. It occurs abundantly on the pine-covered hills of Kashmir.

65. *Hemichelidon ferruginea* : The Ferruginous Flycatcher. F. 559, J. 299, —I. E.

A brown bird in which the greater part of the plumage is tinted with red. The red on the rump almost entirely obscures the brown, so that this part of the bird appears chestnut. A ring of cream-coloured feathers round the eye.

Common at Darjeeling. It usually makes its flights after insect quarry from a low branch or the stump of a tree.

66. *Cyornis superciliaris* : The White-browed Blue Flycatcher. F. 568, J. 310, —I. E. W.

*Cock* : Upper plumage dull Prussian blue, with some brown in the wings. A conspicuous white eyebrow. Tail black, edged with blue ; all feathers, except middle pair, tipped with

---

---

## *Descriptive List of Birds*

---

white. Lower plumage white save for a broad dark blue collaret interrupted in the middle.

*Hen*: Olive-brown above, tinged with dark blue on the lower back. Lower parts pale buff, which fades to white near the tail.

*Very* common at Simla, where it sometimes nests in houses. Abundant at most hill stations except Darjeeling.

67. *Stoparola melanops*: The Verditer Flycatcher. F. 579, J. 301, I. E. W.

This beautiful bird is the commonest flycatcher at *all* the Himalayan hill stations.

The cock is arrayed in Cambridge blue—blue of that peculiar shade known as verditer blue. There is a black band running through the eye.

The hen's plumage is duller than that of the cock. Young birds are greyish-blue and their lower plumage is spotted with cream.

This flycatcher likes to sit on a telegraph wire or a high branch, from which the cock emits a pleasant trill or warble of about a dozen notes.

This species nests in gardens and verandahs.

68. *Alseonax latirostris*: The Brown Flycatcher. F. 588, J. 297, 298, —I. W. K.

The bird may pass for a hen sparrow unless

---

## *Himalayan and Kashmiri Birds*

---

closely scrutinised. The upper plumage is earthy-brown, with darker markings on the head. The lower parts are white tinged with grey on the breast. There is a ring of white feathers round the eye.

This bird is nowhere common. It is rare at Simla and Dharmsala. Fairly common at Naini Tal and Almora.

It may be difficult to distinguish this bird from the Sooty Flycatcher (64). It is a little larger than No. 64 and the edges of the flight feathers have paler edges, the bill is a little longer. This is a commoner bird than No. 64.

69. *Alseonax ruficaudatus*: The Rufous-tailed Flycatcher. F. 589, J. 307, -I. W. K.

Dull olive-brown above. Tail chestnut. Lower plumage pale earthy-brown.

This inconspicuous bird is nowhere common. I have not noticed it at Simla, Mussoorie, or Naini Tal: but it has been recorded from Kumaun. I have seen it at Gulmarg. At Dharmsala it is found up to 6000 feet.

70. *Culicicapa ceylonensis*: The Grey-headed Flycatcher. F. 592, J. 295, +I. E. W. K.

Head, neck and breast grey. Wings and tail brown. Back dull yellow. Lower plumage

---

---

## *Descriptive List of Birds*

---

bright yellow. The yellow in the plumage is not very noticeable except when the bird is on the wing.

It is often found near streams.

It has a loud, pleasant call of three notes, which may be syllabised : *Think-of-me*.

It is common at Naini Tal, Simla and Dharmsala : but not at Darjeeling.

71. *Niltava sundara* : The Rufous-bellied Niltava. F. 594, J. 314, +I. E. W. K.

Jerdon calls this bird the rufous-bellied, fairy blue chat, and the cock indeed is the kind of bird one would expect to meet in fairyland.

*Cock* : Front and sides of head, chin and throat rich velvety-black. Crown, nape, lower back, a spot on each cheek and one on each wing brilliant, glistening cobalt blue. Rest of lower plumage chestnut-red. Tail black and bright blue.

*Hen* : Upper plumage olive-brown, with a white patch on fore-neck and a brilliant blue patch on each side of the neck. Tail and feathers above it chestnut-red. Lower plumage pale olive-brown.

Very common at Darjeeling, less common in the western hill stations and the Kashmir valley.

---

---

## *Himalayan and Kashmiri Birds*

---

72. *Niltava grandis*: The Large Niltava. F. 593, J. 316, —II. E.

Like the last species, but larger, and there is no red on the under plumage, the red being replaced by dull blue. No white patch on fore-neck of hen.

This bird is common near Darjeeling, but does not come up as high as the station.

73. *Terpsiphone paradisi*: The Indian Paradise Flycatcher. F. 598, J. 288, II, but the cock has a very long tail. W. K.

Perhaps the most beautiful and most striking bird in the world. It is sometimes called the Ribbon Bird.

*Hen and Young Cock*: like a bulbul in size and shape. The crested head is metallic blue-black. Back, wings and tail chestnut. Lower parts white.

*Second-year Cock*: As above, but the two middle tail feathers are much longer than the others, being 16 inches long.

*Old Cock*: Chestnut parts of plumage white, with some black in the wing.

It is impossible to mistake this bird as it flits about among trees displaying its long, thin chestnut or white ribbon-like tail feathers which flutter in the air like satin streamers.



---

---

## *Descriptive List of Birds*

---

Very common at Almora and in the Kashmir valley. Common also *below* the other western hill stations, e.g. at the Brewery below Naini Tal.

### *Chats*

Chats are robin-like birds, that feed entirely on insects; these they often catch on the wing after the manner of flycatchers. They frequent open country.

74. *Pratincola caprata* : The Common Pied Bush Chat. F. 608, J. 481, —I. W. K.

*Cock* : Plumage black, except for white wing-bar and large patch on lower back.

*Hen* : Plumage grey, streaked with brown and, on the lower back, tinged with red.

The cock has a sweet little song.

This species is not seen at most hill stations, as it does not seem to ascend the Himalayas above 6000 feet. It is common in the Kashmir valley and at Almora.

75. *Pratincola maura* : The Indian Bush Chat. F. 610, J. 483, —I. W. K.

*Cock* : Upper plumage dark brown, almost black. Large white patch on each side of neck; these patches sometimes unite to form a collar. Lower back white. Breast orange: rest of lower plumage russet.

---

---

## *Himalayan and Kashmiri Birds*

---

*Hen* : Upper plumage brown, without any white. Lower plumage pale-reddish yellow.

Breeds in spring in Kashmir and at Almora, Simla and Naini Tal. At Dharmsala it occurs up to 6000 feet.

76. *Oreicola ferrea* : The Dark Grey Bush Chat. F. 615, J. 486, I. E. W. K.

*Cock* : Upper plumage black with a white eyebrow and some white in the wings and tail. Lower plumage white washed with grey on breast and thighs.

*Hen* : Upper plumage reddish-grey with faint dark spots ; the red is pronounced on the rump and tail. Lower plumage as in cock.

This species habitually makes sallies into the air after its insect quarry and doubtless is often mistaken for a flycatcher.

Very common at Naini Tal, Simla and Dharmsala, and fairly so at other hill stations, except Almora. Occurs in Kashmir valley.

### *Forktails*

Forktails are glorified wagtails. The startling black-and-white plumage renders their identification easy. They live by mountain streams, and utter a loud scream during flight.

77. *Henicurus maculatus* : The Western

---

---

## *Descriptive List of Birds*

---

Spotted Forktail. F. 630, J. 584, II, with a tail of 6 inches. W. K.

A very beautiful bird.

Head and neck jet black, save for a white patch extending from the centre of the crown to the bill. Shoulders black, copiously spotted with white. Wings, black and white. Tail black, each feather tipped with white. As the tail feathers are graduated, the middle pair being the shortest and the outer pair the longest, the tail, when closed, appears to be barred with broad V-shaped black, and narrow white, bars. Legs pale pink. Lower back and rump white.

Inhabits mountain streams, feeding on tiny insects which it picks off the boulders. Nest often within a yard of the water's edge.

Occurs on every hill stream in Kashmir and the Western Himalayas, at places where the water rushes down the hillside.

78. *Henicurus guttatus*: The Eastern Spotted Forktail. F. 631, J. 584, II, with tail  $5\frac{1}{4}$  inches long. E.

This is the East Himalayan race of the last species. Jerdon (rightly in my opinion), does not distinguish between the two.

This is a trifle smaller than the western

---

## *Himalayan and Kashmiri Birds*

---

form. It has fewer white marks on the shoulder and these are smaller and round instead of being oval.

Habits identical with those of the last species. Common on every mountain torrent in the Eastern Himalayas.

79. *Microcichla scouleri*: The Little Fork-tail. F. 637, J. 587, —I. E. W. K.

This is just a short-tailed miniature of the Western Spotted Forktail. Its plumage, markings and habits are the same.

It is less common than the larger species, but is a common bird. In Kashmir and the Western Himalayas it is more abundant at higher elevations than the spotted species, being most common in summer between 6000 and 8000 feet, but in the Eastern Himalayas it is most often seen between 2000 and 5000 feet.

### *Redstarts*

Redstarts are robin-like birds, having red tails which they constantly flirt. Most of them frequent mountain streams.

80. *Chimmarrhornis leucocephalus*: The White-capped Redstart. F. 638, J. 506, +I. E. W. K.

---

---

## *Descriptive List of Birds*

---

A feathered exquisite.

A snow-white cap; remainder of head, whole neck, breast, wings and upper back rich velvety black. Tail, rump and abdomen bright chestnut red, so that, as the bird leaps from a boulder in mid-stream into the air, it looks as though it were on fire.

Common in Kashmir and the Western Himalayas at all elevations above 6000 feet; in summer up to the glaciers. In the Eastern Himalayas, it does not occur as a rule at elevations lower than 7000 feet.

A frequenter of mountain streams.

81. *Ruticilla frontalis*: The Blue-fronted Redstart. F. 639, J. 503, I. E. W. K.

*Cock*: Head and shoulders blue, brightest on forehead. Rump, breast and abdomen chestnut, as is the tail except the middle pair of feathers which are black. Wings brown with reddish margins.

*Hen*: A brown bird with a ring of pale feathers round the eye. Tail as in cock. Lower plumage suffused with orange. This redstart is usually found away from water. In the summer it usually keeps to elevations higher than those of the hill stations, but it is occa-

---

---

## *Himalayan and Kashmiri Birds*

---

sionally seen at Naini Tal and Dharmasala in rhododendron forests.

82. *Rhyacornis fuliginosus*: The Plumbeous Redstart. F. 646, J. 505, —I. E. W. K.

Very robin-like in appearance—Jerdon calls it the water-robin. It picks its food off the boulders of mountain torrents and streams.

*Cock*: Plumage dull dark blue with tail and abdomen rusty-red.

*Hen*: Upper plumage dull bluish-brown. Wings brown. Lower back and base of tail white. Lower plumage greyish-brown, mottled with white.

Common on all streams in Kashmir and the Himalayas, from 6000 to 13,000 feet elevation, except in the Eastern Himalayas where its usual range is from 1500 to 5000 feet.

### *Robins*

83. *Ianthia rufilata*: The Red-flanked Bush Robin. F. 654, J. 508, I. E. W. K.

*Cock*: Forehead, eyebrow and rump ultramarine blue, a patch of this colour on the wings. Cheeks black. Rest of upper plumage and sides of breast deep purple-blue. Tail black and blue. Lower plumage white except the blue sides of the breast and a large showy

---

---

## *Descriptive List of Birds*

---

patch of bright chestnut on each side of the body.

*Hen*: Olive-brown suffused with blue on the rump and tail, a white bib and a bright chestnut patch on the sides of the body.

A forest-haunting bird having a noisy call. It is fairly common at Gulmarg and Dharm-sala in summer, but at other hill stations it seems to occur only in winter.

84. *Copsychus saularis*: The Magpie-Robin. F. 663, J. 475, —II. W.

*Cock*: All plumage glossy black, except the following which are white: Abdomen, outer tail feathers, and wing-bar.

*Hen*: Grey-brown where cock is black; white as in cock.

This species frequently elevates the tail.

This is essentially a bird of the plains, but in summer some individuals ascend the Himalayas.

It occurs in summer at Almora, Dharm-sala and Simla, where it visits gardens. I have seen the nest at Almora. 6500 feet appears to be the limit of its upward range and 6000 feet of the nesting range in the Western Himalayas. In the Eastern Himalayas it is rarely, if ever, seen at altitudes over 3000 feet.

---

---

## *Himalayan and Kashmiri Birds*

---

### *Blackbirds and Thrushes*

85. *Merula bouboul* : The Grey-winged Ouzel. F. 676, J. 361, +III. E. W.

The *Kastura* of Indians. Perhaps the finest songster of the Himalayas. From Murree to Darjeeling, between 6000 and 9000 feet the hills ring with the blackbird-like song of this species.

*Cock* : Black with silvery-grey on the wings.

*Hen* : Olive-brown. The bill is coral-red in both sexes.

86. *Merula unicolor* : Tickell's Ouzel. F. 678, J. 356, +II. K.

*Cock* : An ashy-grey bird.

*Hen* : Upper plumage olive-brown. Wings and tail dark brown. Chin, throat and abdomen white. Breast ashy-brown with dark spots. Bill yellow.

The common thrush of the Kashmir valley. Its blackbird-like song is to be heard in spring in every garden and grove. Nests in poplar trees.

87. *Petrophila cinclorhyncha* : The Blue-headed Rock-Thrush. F. 691, J. 353, -II. E. W. K.

*Cock* : A handsome and striking bird. Head,



---

---

## *Descriptive List of Birds*

---

chin, throat and wing-patch bright cobalt-blue. Sides of neck, shoulders, upper back and a broad band through the eye black. Rump and lower parts chestnut. Wings black and blue with a white bar.

*Hen* : Olive-brown above, cream below, with dark bars on rump and breast.

Cock has a sweet, soft song, rather like that of the white-eye. When alarmed he utters a curious harsh cry. As its name implies this bird frequents rocky ground, but likes to have trees near at hand to which it can repair when alarmed.

Common at all hill stations, including Almora. Occurs, but is not common, in Kashmir.

### *Dippers*

Dippers are small, short-tailed, dark-plumaged birds that inhabit mountain streams, into the waters of which they dive for their food.

88. *Cinclus asiaticus* : The Brown Dipper. F. 709, J. 347, II, tail 2 inches long. E. W. K.

Plumage chocolate-brown, darkest on wings and tail, more or less flecked with white.

Dippers occur on most mountain streams, but they are not so common as forketails. In

---

## *Himalayan and Kashmiri Birds*

---

the Western Himalayas and Kashmir dippers spend the summer at elevations between 6000 feet and the snow-line. In the Eastern Himalayas they are more commonly found between 1000 and 5000 feet.

### *Finches*

Finches, of which the sparrow and canary are familiar examples, are thick-billed birds that feed largely on grain. The stout bill is an organ adapted to husking seeds. The family is well represented in the Himalayas.

89. *Pycnorhamphus icteroides*: The Black-and-Yellow Grosbeak. F. 741, J. 725, -III. W. K.

*Cock*: Coloured very like a black-headed Oriole.

Head, throat, wings, shoulders and a patch above the tail black. Rest of plumage rich yellow tinged with orange on the hindneck. Bill very stout.

*Hen*: Wings black and grey. Rest of plumage ashy-grey, tinged faintly with yellowish-red on the back and abdomen.

This species goes about in couples or small flocks and feeds largely on the ground. Call sounds like *kiu-kree*.

---

---

## *Descriptive List of Birds*

---

Fairly common at Murree and Dalhousie, but less so at the other western hill stations and in Kashmir.

90. *Hæmatospiza sipahi*: The Scarlet Finch. F. 751, J. 735, -II. E.

*Cock*: Plumage brilliant scarlet with some black in wings and tail.

*Hen*: Upper plumage brown with greenish-yellow crescent-shaped markings. Rump bright yellow. Lower plumage dull pale yellow with dark crescent-shaped markings. Bill stout.

Darjeeling is the only hill station at which this fine finch occurs.

91. *Propasser grandis*: The Red-mantled Rose Finch. F. 757, J. 741, +I. W. K.

*Cock*: Most of the plumage brown tinged with rosy red. Eyebrow, sides of head, chin and throat shining rosy pink. Rest of lower plumage rosy-red.

*Hen*: Grey-brown, darkest in the upper plumage and everywhere streaked with dark brown. Wings and tail brown.

I have not noticed this bird at any of the Himalayan hill stations, as it seems to live at higher elevations, but it is fairly common at Gulmarg.

---

## *Himalayan and Kashmiri Birds*

---

92. *Carduelis caniceps* : The Himalayan Goldfinch. F. 767, J. 749, —I. W. K.

Very like the European Goldfinch.

Forehead, chin and part of face in front of eye crimson, rather dull in the hen. Wings gold and black. Rest of upper plumage grey, merging into white on the lower back. Tail black and white.

Fairly common in Kashmir, less so in the Western Himalayas. Does not occur east of Kumaun.

93. *Callcanthis burtoni* : The Red-browed Finch. F. 768, J. 748, +I. W. K.

Both cock and hen are striking birds.

Crown and nape black in cock, dark brown in hen. Cock has forehead and large patch of feathers round the eye crimson. In the hen these parts of the plumage are buff. Wings and tail blue-black with a good deal of white, which is very conspicuous during flight. When the wings are closed the white takes the form of square spots, so that the birds have then somewhat the appearance of miniature jays.

These birds go about in pairs, sometimes in company with tits.

As they live at high elevations in summer,

---

---

## *Descriptive List of Birds*

---

they are not likely to be seen at any Himalayan station, but they are common at Gulmarg.

94. *Hypacanthis spinoides* : The Himalayan Greenfinch. F. 772, J. 750, —I. E. W. K.

A little bird that keeps largely to the tops of trees and utters a rather mournful *pee-pee-pee-pee*.

Head and upper back dull green, with a yellow eyebrow and a patch of this hue (which sometimes becomes a collar) on each side of the neck. Wings and tail brown and yellow. Lower plumage bright yellow.

The hen is a dull edition of the cock.

This bird in summer lives at higher altitudes than most hill stations ; but may be observed occasionally at Simla, Gulmarg and Naini Tal in summer.

95. *Passer domesticus* : The House Sparrow. F. 776, J. 706, I. W. K.

*Cock* : Top of head and lower back ashy-grey. Throat black. Cheeks and sides of head pure white. Streak over the eye and upper back chestnut. Wings and tail brown, the former with a narrow white bar. Lower plumage sullied white.

*Hen* : a dull brown bird with sullied white

---

---

## *Himalayan and Kashmiri Birds*

---

under-parts. Cream-coloured eyebrow and narrow white wing bar.

Abundant at Kasauli, Almora, Simla, Dharm-sala and Srinagar. Brooks in 1868, found it common at Naini Tal. I saw it nesting there in the Club verandah twenty years later ; but it seems to be becoming scarcer there, rather than more abundant.

96. *Passer montanus* : The Tree-Sparrow. F. 779, J. 710, —I. E.

Sexes alike. Head chestnut with sides of face white and a black patch below and behind eye. Chin and throat black. Shoulders and upper back are pale chestnut, which fades into yellowish-brown on the rump. Tail brown. Wings brown, chestnut, black and white. Lower parts pale ashy-grey.

The only sparrow found at Darjeeling. I have not noticed it at any other hill station.

97. *Passer cinnamomeus* : The Cinnamon Tree-Sparrow. F. 780, J. 708, —I. W. K.

*Cock* : Upper plumage bright cinnamon, with pale yellow cheeks and a small black patch round the eye. Wings black and white with reddish streaks. Tail brown. Chin and throat black, with bright yellow patch on either side. Rest of lower plumage grey tinged with yellow.

---

---

## *Descriptive List of Birds*

---

*Hen*: Upper plumage reddish-brown, with dark streaks. Eyebrow yellow above, dark brown below. Tail brown. Wings brown, black and white. Sides of head and neck and lower plumage grey tinged with yellow.

This sparrow is common in Kashmir, both in the valley and on the hills, and at the West Himalayan hill stations, including Naini Tal, but not Almora.

### *Buntings*

The English yellow-hammer is a familiar member of the bunting family. Buntings are finch-like birds somewhat larger than the sparrow. They are characterised by having a conical bill which terminates in a sharp point.

98. *Embiriza stewarti*: The White-capped Bunting. F. 793, J. 718, +I. W. K.

*Cock*: Light-grey cap. Upper plumage chestnut. Wings and tail dark brown, with a good deal of white in the tail. Chin and throat black with white patch in middle.

*Hen*: A brown bird streaked with darker brown, washed with chestnut on shoulders and rump. Some white in the tail.

Fairly common in Kashmir on lower slopes

---

---

## *Himalayan and Kashmiri Birds*

---

of hills. In the Himalayan stations this species is not often seen as in the summer it lives rather above their level.

99. *Emberiza stracheyi*: The Eastern Meadow Bunting. F. 794, J. 714, +I. W. K.

This is the common bunting of Kashmir and the western hill stations.

Easily recognised by two longitudinal black bands (with a grey one between them) running from the bill over the head to the neck, and two black moustachial stripes.

Upper plumage reddish-brown streaked with black. Throat and breast grey with brown spots. Rest of lower plumage chestnut brown. Tail black, with a good deal of white in the outer feathers, visible only during flight.

This bird feeds on the ground and is common in gardens and on open ground. Abundant at Gulmarg, Murree, Dharmsala, and Simla.

### *Swallows*

100. *Hirundo rustica*: The Swallow. F. 813, J. 82, I, with tail  $4\frac{1}{2}$  inches long. E. W. K.

This is the common swallow of England.

Forehead, chin and throat chestnut; rest of upper plumage steel-blue. Wings and tail



---

---

## *Descriptive List of Birds*

---

black, glossed with green. Some white in the tail. Lower plumage reddish-cream with a broad black band across the breast. This bird does not usually ascend as high as the hill stations, but is common in the Kashmir valley.

101. *Hirundo smithii*: The Wire-tailed Swallow. F. 818, J. 84, -I, with two outer feathers, the ends of which look like wires, 7 inches long. W. K.

Forehead, crown, and nape chestnut. Lower plumage white. Rest of colouring like that of the swallow. The outer tail feathers are long with wire-like ends.

This species, like the last, rarely comes as high as the Himalayan hill stations, but is occasionally seen in the Kashmir valley.

102. *Hirundo erythropygia*: Sykes's Striated Swallow. F. 823, J. 85, +1, with tail 3 inches long. E. W.

Upper parts steel-blue, except the collar and rump which are chestnut-red. Lower plumage very pale red, narrowly streaked with brown. Wings and tail brown, glossed with blue. A little white in the tail.

The common swallow of all the Himalayan hill stations.

---

## *Himalayan and Kashmiri Birds*

---

### *Wagtails*

Wagtails are slightly built, elegant, rather long-tailed birds, which frequent the water's edge and run about over stones seeking their insect quarry. The tail constantly wags up and down.

Wagtails usually frequent streams in which the water does not flow very rapidly, hence are not often seen at hill stations; but three species are common in Kashmir.

103. *Motacilla hodgsoni*: Hodgson's Pied Wagtail. F. 830, J. 590, II. K.

Face white; rest of upper plumage, throat and breast black. Rest of lower plumage white. Wings dark brown and white. Tail black and white.

Common on the Kashmir lakes. It runs about on the leaves of the lotus and its flight is undulating.

104. *Motacilla melanope*: The Grey Wagtail. F. 832, J. 592, +I, with tail  $3\frac{1}{2}$  inches long. K.

Head, neck and shoulders bluish-grey with cream-coloured eyebrow. Rest of upper plumage greenish-yellow and black. Chin and throat black, rest of lower plumage bright yellow.

---

---

## *Descriptive List of Birds*

---

This species frequents the stream that runs through the golf links at Gulmarg.

105. *Motacilla citreoloides* : Hodgson's Yellow-headed Wagtail. F. 838, J. 594, +I, with tail over 3 inches long. K.

Head, neck and lower plumage rich deep yellow. Rest of upper plumage velvety black. Wings dark brown and white. Tail black and white.

Common on the Kashmir lakes. It runs about on the floating leaves of the lotus. Occurs also in gardens on lawns.

### *Pipits*

Pipits are dull-plumaged birds, somewhat like wagtails in shape; they feed on the ground, but not at the water's edge as wagtails do.

Several species should be placed among the common birds of Kashmir and the Himalayas, but it is not easy to describe such dull birds so as to enable the reader readily to distinguish them from one another.

106. *Anthus maculatus* : The Indian Tree-pipit. F. 841, J. 596, I. W.

Upper plumage sandy brown suffused with

---

## *Himalayan and Kashmiri Birds*

---

green, with indistinct dark streaks and inconspicuous creamy eyebrow. Tail brown and white, Lower plumage pale cream with conspicuous dark streaks on the throat.

107. *Anthus similis*: The Brown Rock-Pipit. F. 841, J. 604, II. W.

Upper plumage grey-brown with dark streaks. No white in tail. Eyebrow sandy yellow. Breast sandy yellow with a few ill-defined dark streaks.

108. *Anthus rufulus*: The Indian Pipit. F. 847, J. 600, +I. E. W. K.

Very like Indian Tree-Pipit, but upper plumage is not suffused with green, and lower plumage is darker.

This species does not often come as high as Indian hill stations, especially in the west; but it occurs in the Kashmir valley.

109. *Oreocorys sylvanus*: The Upland Pipit. F. 853, J. 606, +I. E. W.

Upper plumage very dark brown with reddish-buff edges to the feathers. Indistinct pale eyebrow. Tail as in Indian Tree-Pipit. Chin and throat white, rest of lower plumage buff with black streaks.

---

---

## *Descriptive List of Birds*

---

### *Skylarks*

Skylarks are in appearance dull-brown birds ; in shape they resemble pipits ; but, unlike pipits, they rarely perch anywhere save on the ground and they sing while soaring in the air.

110. *Alauda arvensis* : The Skylark. F. 860, J. 766, +I. W. K.

Upper parts dark brown with pale streaks. An indistinct pale eyebrow. Lower parts pale sandy with dark brown and black streaks. Wings and tail brown ; some white in the two outer tail feathers.

This is the skylark of England.

I have not noticed it at any Himalayan station, but it is common in the Kashmir valley, where it sings in the air just as it does in England.

### *Sunbirds*

Sunbirds are the Asiatic counterparts of the South American Humming-birds. They are small birds, of which the plumage of the cocks is iridescent. The bill is long, slender, and curved and the tongue tubular. The birds live largely on the nectar of flowers, which they often sip while hovering on vibrating wings.

---

## *Himalayan and Kashmiri Birds*

---

They also eat insects. The hen is a bird of dull plumage.

111. *Æthopyga ignicauda*: The Fire-tailed, Yellow-backed Sunbird. F. 887, J. 228, —I. E.

*Cock*: Upper plumage crimson, with metallic-blue crown and yellow rump. Lower plumage yellow, with purple on throat and some crimson on breast. Middle pair of tail feathers elongated.

*Hen*: A dull yellow-green bird with brown wings and tail and some yellow on lower parts. Middle tail feathers not lengthened.

It is only likely to be seen at Darjeeling in April, while it is migrating towards its breeding grounds at high elevations. In April the cocks have not got their full plumage. They are then like the hen, but with bright patches of colour.

112. *Æthopyga nepalensis*: The Nepal Yellow-backed Sunbird. F. 892, J. 229, —I. E.

*Cock*: Head, neck and throat metallic-green, with black on sides of head and deep red on sides of neck. Upper back deep red or maroon. Rest of back yellow, brightest on rump. Tail, of which middle pair of feathers is longer than the others, metallic-green and black; all feathers except the two middle

---

---

## *Descriptive List of Birds*

---

pairs have pale tips. Lower plumage yellow, washed and streaked with crimson.

*Hen*: Olive-green. Middle tail feathers not lengthened. The common sunbird of Darjeeling. Found in gardens.

The Western form of this species, *Æthopyga scheriæ* (The Himalayan Yellow-backed Sunbird), does not ascend the mountains as high as the hill stations, but I have seen it at the Brewery below Naini Tal.

It may be distinguished from the Eastern form by the fact that the chin and throat are crimson.

### *Flowerpeckers*

Flowerpeckers are minute birds which keep to the topmost branches of trees.

113. *Dicæum ignipectus*: The Fire-breasted Flowerpecker. F. 915, J. 241, —I. E. W.

One of the smallest birds in India. As it does not ascend the hills above 6000 feet, Almora is the only hill station at which it is likely to be seen.

*Cock*: Upper parts black glossed with green. Lower parts buff, with large patch of crimson on breast and a smaller black one.

*Hen*: Olive-brown above, buff below.

---

## *Himalayan and Kashmiri Birds*

---

### *Woodpeckers*

Woodpeckers are birds that feed on insects which they pick off the trunks of trees. They have strong claws by means of which they are able to cling to the bark. The feathers of the tail are stiff and serve to support the woodpecker when it is climbing. The sharp chisel-like bill enables the bird to dig out insects that lurk in the crevices of the bark, and also to excavate holes in the trunk in which the eggs are laid. When a woodpecker moves about a tree-trunk, the head always points upwards. Its movements on tree-trunks are jerky, like those of a mechanical toy. The flight from tree-trunk to tree-trunk is undulating. When on the wing a woodpecker often emits a loud scream.

114. *Gecinus squamatus*: The West Himalayan Scaly-bellied Green Woodpecker. F. 946, J. 170, —IV. W. K.

*Cock*: Upper plumage dull green, tinged with yellow on the lower back. Crown and nape crimson. A black band round the face and one from below the eye to the bill. Wings and tail brownish-black, spotted and banded with white and cream.



---

---

## *Descriptive List of Birds*

---

*Hen*: Like cock, but head and nape black instead of crimson. Common at Simla, Mussoorie, and Naini Tal. Not very common at Dalhousie, Dharmsala, and Murree, and in Kashmir.

115. *Gecinus occipitalis*: The Black-naped Green Woodpecker. F. 950, J. 172, +III. W. K.

Like the last species, but in the cock the nape is black instead of crimson.

Common at Simla and Dharmsala.

116. *Gecinuculus grantia*: The Northern Pale-headed Woodpecker. F. 958, J. 177, +II. E.

*Cock*: Forehead, cheeks and chin light brown. Crown red. Nape yellow. Rest of upper plumage dull red. Wings brown spotted with cream. Tail brown, cream, and red.

*Hen*: Like cock, but whole head is dull yellow.

Darjeeling is the only hill station at which the woodpecker occurs.

117. *Hypopicus hyperythrus*: The Rufous-bellied Pied Woodpecker. F. 960, J. 161, II. E. W.

Upper plumage spotted black and white. Lower plumage dull red. A patch of crimson

---

---

## *Himalayan and Kashmiri Birds*

---

under the tail. Crown red in cock, black spotted with white in hen.

Although this woodpecker ranges all along the Himalayas, Murree is the only station at which I have seen it.

118. *Dendrocopus himalayensis*: The Western Himalayan Pied Woodpecker. F. 961, J. 154, +II. W. K.

*Cock*: Crown crimson. Upper plumage black. Face and neck white, separated by a black band. There is also a black band behind the ear. Lower parts grey.

*Hen*: As cock, but her crown and nape are black like the back. Common at Dalhousie, Murree, and Gulmarg, less so at the other western hill stations, where the Brown-fronted Pied Woodpecker is the commonest species.

119. *Dendrocopus cathpharius*: The Lesser Pied Woodpecker. F. 965, J. 156, +I. E.

A small eastern form of the last species. It is distinguished by both the cock and hen having crimson on the sides of the neck. The grey lower plumage is streaked longitudinally with brown.

120. *Dendrocopus auriceps*: The Brown-fronted Pied Woodpecker. F. 969, J. 159, II. W. K.

---

---

## *Descriptive List of Birds*

---

This is the commonest woodpecker of all the West Himalayan hill stations, except perhaps Dalhousie, where the Western Himalayan pied species is commoner. It is distinguished from the other pied woodpeckers by the umber-brown forehead and crown. The cock's crest is golden in front and crimson behind; the crest of the hen is pale yellow.

121. *Pyrrhopicus pyrrhotis*: The Red-eared Bay Woodpecker. F. 978, J. 176, +III. E.

Head brown. Cock only has crimson patch behind each ear; patches sometimes unite behind to form a half collar. Rest of upper plumage barred black and dull red. Lower parts olive-brown washed with red.

Fairly common at Darjeeling.

122. *Lynx torquilla*: The Common Wryneck. F. 1003, J. 188, +I. K.

An inconspicuous grey-brown bird with dark bars and streaks all over the body, giving the plumage a mottled appearance.

This bird runs up and down the trunks of trees after the manner of woodpeckers. It has a peculiar habit of twisting its head, hence the name Wryneck.

Fairly common in Kashmir.

---

---

## *Himalayan and Kashmiri Birds*

---

### *Barbets*

Barbets are stout-billed, fruit-eating, tree-haunting birds. They have monotonous calls of two or three notes, which they utter frequently. They lay their eggs in holes which they excavate in trees.

123. *Megalæma marshallorum*: The Great Himalayan Barbet. F. 1006, J. 191, +III. E. W.

A fine bird. Head and neck violet-blue. Upper back olive-brown, streaked longitudinally with pale green. Lower back and tail bright green. Wings green, with splashes of blue, brown and yellow. Upper breast brown. Red patch under tail. Rest of lower parts yellow, with median blue band. The massive bill is yellow.

This bird sometimes goes about in small flocks. Its call, which is a loud, plaintive *Pee-you, pee-you*, resounds on the hill-sides and in the valleys.

Common at Murree, Dharmsala, Mussoorie and Naini Tal; less so at Dalhousie, Simla and Darjeeling.

124. *Cyanops franklini*. The Golden-throated Barbet. F. 1017, J. 196, II. E.

---

---

## *Descriptive List of Birds*

---

A grass-green bird washed with blue. The head is gaudily coloured. Top of head, chin and upper throat golden-yellow. Forehead crimson, a black streak on each side of head.

Its cry is *Katak-katak-katak*.

Common at Darjeeling.

### *Rollers*

Heavy-billed, bright-plumaged birds, which take up a position on a perch and descend to the ground to secure their insect quarry. They sometimes eat small snakes. Their cries are harsh.

125. *Coracias garrula* : The European Roller. F. 1024, J. 125, +III. K.

This is a near relative of the roller, or "blue jay" of the plains.

Its plumage is mainly light and dark blue; the two shades of blue being arranged in broad bands on the wings and tail. A gorgeous bird.

Fairly common in the Kashmir valley.

### *Bee-eaters*

Bee-eaters are elegant emerald-green birds. The middle pair of tail feathers project as bristles beyond the others.

Bee-eaters catch their insect quarry in the

---

## *Himalayan and Kashmiri Birds*

---

air after the manner of flycatchers. The expanded wings are triangular in shape. Bee-eaters nest in holes in sandbanks.

126. *Merops apiaster*: The European Bee-eater. F. 1029, J. 121, II, with a tail 6 inches long. K.

Forehead white, rest of head, neck, shoulders and upper back chestnut, darkest on the head. Wings chestnut, green and black. Tail bronze-green, the long middle pair of feathers are black at the ends. Chin and throat yellow. Rest of lower parts bluish-green. Black band through eye. Black gorget. Bill black. Wing lining buff.

This fine bird is often to be seen hawking insects over the lakes in the Kashmir valley.

### *Kingfishers*

Kingfishers live on fish for which they dive: the bill is long and the tail short. They nest in holes which they excavate in river-banks.

127. *Ceryle lugubris*: The Himalayan Pied Kingfisher. F. 1034, J. 137, —IV. E. W. K.

Plumage white, boldly barred, spotted and streaked with black. Bill black.

This fine bird is found on the Kashmir lakes and the larger hill streams up to 6000 feet.

---

---

## *Descriptive List of Birds*

---

It quarters the water in search of fish. Every now and then it hovers on rapidly vibrating wings. If it espies a fish in the water below, its pinions cease quivering and it drops like a stone into the water. Sometimes it checks its fall and flies to another spot where it repeats the operation. It is impossible to mistake this bird. Its colouring and habits are unique.

128. *Alcedo ispida*: The Common Kingfisher. F. 1035, J. 134, II, but tail very short. E. W. K.

Head and nape blue with faint black cross-bars. Back bright, glistening pale blue. Tail dark blue. Wings greenish-blue. Sides of head show streaks of red, blue, black and white. Chin cream-coloured. Rest of lower parts rust-coloured. Bill black. Legs coral-red. Neck and tail short.

This bird squats on a bank, or a branch overhanging water, watching for fish. As it sits its head bobs up and down. When it espies a fish it dives into the water. Its call is a shrill scream.

This bird occurs on all the larger hill streams up to about 5500 feet. It is very common on the Kashmir lakes, and is fond of perching on the heads of lotus flowers and on house-boats.

---

---

## *Himalayan and Kashmiri Birds*

---

### *Hoopoes*

Hoopoes are birds that feed on the ground, into which they probe with their long, slender, slightly-curved bills. The crest is conspicuous. When folded it looks like a backward continuation of the bill. When a hoopoe is disturbed, and invariably when the bird alights, the crest is expanded like a fan.

129. *Upupa epops*: The European Hoopoe. F. 1066, J. 254, III. E. W. K.

Body plumage sand-coloured or pale fawn. Crest reddish-fawn with black tips to feathers. There is also some white in the crest. Wings and tail white with very bold black bars. Legs short. On the wing this bird looks like a huge black-and-white butterfly. A bird that cannot be mistaken for any other.

It is fairly common in the Kashmir valley and at Almora. Simla is the only other hill station at which I have noticed it.

### *Swifts*

Swifts are birds which never descend to the ground or perch in trees. Their claws are so arranged that they cannot support themselves on a perch. When they require rest,



---

---

## *Descriptive List of Birds*

---

they betake themselves to their nests, which are saucer-shaped structures made of straw, feathers, etc., cemented together and to the wall of a building by the birds' saliva. Their wings are very long and slender. As a swift dashes through the air at great speed it is shaped like an anchor with very large flukes.

130. *Cypselus affinis*: The Common Indian Swift. F. 1073, J. 100, —I. W. K.

A very dark brown, almost black, bird with a white bar across the lower back.

This is the common swift of the West Himalayan hill stations, and of Kashmir. It nests under the eaves of houses. It utters what Jerdon describes as a "shivering scream," which is sometimes heard at night. Most dak bungalows in the W. Himalayas have nests of this swift. A colony nests in the Landour bazaar at Mussoorie.

A larger bird, sometimes seen in Kashmir and the Himalayas, with no white bar across back and a white abdomen, is *Cypselus melba*, the Alpine Swift.

131. *Chætura nudipes*: The White-necked Spinetail. F. 1077, J. 97, II. E. W.

Upper parts blackish-brown, glossed with green. Middle of back and chin white.

---

---

## *Himalayan and Kashmiri Birds*

---

Breast and abdomen brown. Tail short, square, and spiny.

The only hill station at which this bird is common is Darjeeling.

### *Cuckoos*

Cuckoos are remarkable birds. Most of them are parasitic on other birds. They fly swiftly. They have loud, distinctive calls.

132. *Cuculus canorus* : The Cuckoo. F. 1104, J. 199, —III, with rather long tail. E. W. K.

Upper plumage, chin, throat and upper breast grey. Rest of lower plumage white, with numerous narrow dark brown cross-bars. Tail tipped with white. Eyes and legs yellow.

This is the common cuckoo of England. Its call, from which it derives its name, is familiar to every Englishman. From April to June the call is heard at every hill station and throughout the Kashmir valley. A bird more often seen than heard.

133. *Cuculus saturatus* : The Himalayan Cuckoo. F. 1105, J. 200, —III, with the tail rather long. E. W.

Like the last species in appearance, but the grey of the upper plumage is much darker

---

---

## *Descriptive List of Birds*

---

and the lower parts have the background pale buff and the dark cross-bars more regular.

The call sounds like the *uk-uk-uk* of the hoopoe, and is doubtless often mistaken for it. As a matter of fact the call has five syllables and may be rendered *cuck-coo-coo-coo-coo*. The first two syllables are like the cuckoo of the common cuckoo, but are uttered so softly as to be inaudible at greater distances than a few yards. At a distance the call sounds like *coo-coo-coo*.

This cuckoo is very common at Naini Tal; fairly so at Mussoorie, Dharmsala, Dalhousie and Darjeeling. I do not remember hearing it at Simla or Murree.

134. *Cuculus poliocephalus* : The Small Cuckoo. F. 1106, J. 201, +II. E. W.

A small edition of the cuckoo, with the lower parts tinged with buff and broader black bars on the breast.

The call of this bird has been rendered *Pot-you-chick-chick-chick*. It is harsh and uttered both by day and by night.

Darjeeling is the only hill station at which this cuckoo is common.

135. *Cuculus micropterus* : The Indian Cuckoo. F. 1107, J. 203, -III, with rather a long tail. E. W.

---

---

## *Himalayan and Kashmiri Birds*

---

Head grey. Back brown. Tail barred black and white, thin throat and upper breast ashy-grey. Rest of lower parts cream-coloured with black crossbars, about half as broad as the pale interspaces.

Call very distinctive. It may be rendered by the words *wherefor*, *wherefor* musically intoned. In imitation of its call the bird is often called the Kyphul pakka, or broken-pekie bird.

Common at Darjeeling, Naini Tal and Mussoorie, but further west it does not ascend the Himalayas as far as the hill stations; the bird is, however, sometimes heard at Dalhousie.

136. *Hierococcyx sparveroides*: The Large Hawk-Cuckoo. F. 1108, J. 207, III, with rather a long tail. E. W.

Head grey. Back brown. Tail brown with broad dark cross bands and tipped with white. Lower plumage pale rusty with brown crossbars.

This is a large race of the Hawk-Cuckoo, or brain-fever bird of the plains. I am not able to distinguish between the calls of the two species. It is a loud crescendo scream, which may be rendered, "brain-fever, *brain-fever*, BRAIN-FEVER."

---

---

## *Descriptive List of Birds*

---

Very common at Mussoorie and Darjeeling : less so at Naini Tal and Dharmsala. I do not remember hearing the bird at Simla or Murree.

137. *Coccytes jacobinus* : The Pied Crested Cuckoo. F. 1118, J. 212, +II, with tail 6 inches long. W. K.

A handsome bird.

Upper plumage glossy black. Broad white wing-bar. Tail tipped with white. Lower plumage white. A conspicuous black crest.

The call is difficult to describe, but easy to recognise. Jerdon well describes it as "high-pitched, wild and metallic."

This bird rarely comes as high as the hill stations. Jones states that it has been seen at 8000 feet near Simla and that it is common up to 5500 feet. It is fairly abundant in the Kashmir valley.

138. *Eudynamis honorata* : The Indian Koel. F. 1120, J. 214, III, with tail 8 inches long. W. K.

*Cock* : A glossy black bird with a green bill and brilliant crimson eye. As he flies he looks like a slenderly-built crow with an unusually long tail.

*Hen* : A brown bird heavily spotted and

---

---

## *Himalayan and Kashmiri Birds*

---

barred with white all over. Bill and eye as in cock.

A very noisy bird, especially at dawn. It has three distinct calls. The commonest is a melodious crescendo, "Ku-il, *Ku-il*, KU-IL," whence its name. The second may be syllabised *Ku-y-o*. The third is a series of loud *Kekarees*.

Common in the Kashmir valley and at Almora. I have never seen or heard the bird at any other hill station.

### *Paroquets*

Paroquets are long-tailed parrots, of which the plumage is mainly grass-green. They associate in flocks, and feed on grain and fruit. They are noisy and destructive to crops.

139. *Palæornis nepalensis*: The Large Indian Paroquet. F. 1135, J. 147, +III, with a tail over a foot long. E. W.

A beautiful grass-green bird with some blue in the tail and a red patch on each shoulder. The cock has a pink collar round the back of his neck, which is connected with the beak on each side by a black band.

I have not seen this bird at any hill station, but it is tolerably common in Kuman, Nepal

---

---

## *Descriptive List of Birds*

---

and Garhwal, and there seems no reason why it should not appear sometimes in the hill stations. It is a common bird in the plains of Northern and Central India, particularly in the Punjab.

140. *Palæornis cyanocephalus* : The Western Blossom-headed Paroquet. F. 1139, J. 149, II, with tail 8 inches long. W. K.

*Cock* : General colour bright grass-green, with some blue on the tail. Red patch on wing. Head red with a bloom like that on a plum.

*Hen* : Head dull grey-blue. Rest of plumage as in cock. The only hill station at which this bird is likely to be seen is Dharmsala. It occurs in the Kashmir valley.

141. *Palæornis schisticeps* : The Slaty-headed Paroquet. F. 1141, J. 150, II, with tail 9 inches long. E. W. K.

Distinguished from the last species by the head being slate-coloured in both sexes, and the cock alone having a red patch on the wing.

The common green parrot of the West Himalayan hill stations. Abundant at Murree, Dharmsala, Simla, Mussoorie and Naini Tal. Not common at Darjeeling. Occurs on Kashmir hills, but not in the valley.

---

---

## *Himalayan and Kashmiri Birds*

---

### *Owls*

Owls are nocturnal birds of prey. Their structure is well-adapted to their nocturnal, predacious habits: they have soft, silky plumage, large forwardly-directed eyes, big ears (often surrounded by aigrettes or tufts of feathers), and feathered legs. The colouring of all is very similar—mottled brown. They have distinctive calls, which, however, are not easy to assign to the birds that emit them because these are so nocturnal in their habits. Owing to the uniform colouring that runs through the family, it is difficult so to describe owls as to make it easy for the casual observer to identify each species he sees.

142. *Syrnium nivicola*: The Himalayan Wood-Owl. F. 1158, J. 66, —IV. E. W.

No aigrettes or ear tufts. Tail rather long for an owl. Plumage brown with creamy spots and thin wavy lines. Lower plumage cream-coloured with dark longitudinal streaks and crossbars.

I have not identified its call. Mr. A. E. Jones writes, "This owl has a variety of notes besides the usual 'Who-hoo.' One is a peculiar squawk which can easily be imitated by placing



---

---

## *Descriptive List of Birds*

---

a blade of grass between the two thumbs of one's hands and blowing sharply on the edge of the grass. Another is similar to the note of the domestic pigeon when he is showing off before the female."

This owl usually keeps to higher levels than those of the hill stations ; but is fairly common at Simla.

143. *Scops spilocephalus* : The Spotted Himalayan Scops Owl. F. 1175, J (not described), —II. E. W.

Large aigrettes. Mottled brown above. Lower plumage white, speckled and spotted with brown.

I have not noticed this owl at any hill station, but Jones states that it is fairly plentiful up to 8000 feet in the Simla rhododendron forests, where it breeds in April. He describes the call as a double whistle—*who-who*.

144. *Glaucidium cuculoides* : The Large Barred Owlet. F. 1183, J. 79, +II. E. W.

No aigrettes. Upper plumage brown with white, cream or reddish crossbars. These crossbars serve to distinguish this from most of the other owls, which have the upper plumage mottled rather than barred.

A noisy bird. Blanford describes its call as

---

---

## *Himalayan and Kashmiri Birds*

---

“ a peculiar cackle, like a laugh ” ; Hume speaks of it as “ a clucking, vibrating call.”

This owl is often out in daylight. Hingston has noticed that it is often mobbed by bulbuls at Dharmsala, where it is the commonest owl.

Fairly common at Simla ; not common at other hill stations.

145. *Glaucidium brodiei* : The Collared Pigmy Owlet. F. 1186, J. 80, +I. E. W.

Very like the last species, but is much smaller and distinguishable by having a buff collar.

More diurnal than most owls. I once saw one moving about at noon at Mussoorie.

There is some difference of opinion regarding the call of this bird. I believe it is the soft, low, single whistle repeated at intervals of about 30 seconds, which is often heard at Mussoorie, where the bird is sometimes known as the half-minute bird. Jones and Hutton, however, state that the call consists of four clear whistled notes—*hoot-oot-oot-oot*, or *whoo-whoo-whoo-whoo*. Jerdon again speaks of its “ low monotonous call or whistle of two notes.”

This bird occurs at most hill stations.

---

---

## *Descriptive List of Birds*

---

### *Vultures*

Vultures are large, dull-plumaged birds which feed chiefly on carrion. Most of them have the head and neck devoid of feathers, and hooked bills adapted to tearing flesh. Their habit is to sail high in the air on outstretched wings, watching for some dead animal to devour.

146. *Otogyps calvus*: The Black or Pondicherry Vulture. • F. 1191, J. 2, +V, nearly twice the size of the kite. E. W.

Plumage black, with white thighs. White breast. Wattles and bare head and neck red.

Fairly common at most hill stations.

147. *Gyps himalayensis*: The Himalayan Griffon. F. 1193, J. 3, +V, more than twice the size of the kite. W.

A dingy, grey-brown bird, washed with buff on rump. No white in wings.

The commonest vulture of the Western Himalayan stations.

148. *Pseudogyps bengalensis*: The Indian White-backed Vulture. F. 1196, J. 5, +V, nearly a yard long. E. W.

A brownish-black bird with the lower back white. This white, which is visible when the

---

---

## *Himalayan and Kashmiri Birds*

---

bird is sitting, renders the identification of the bird easy. There is some white in the wings, visible only during flight. On the wing the white appears as a very broad band running from the body nearly to the tips of the wing, so that the wings look white with broad black edges.

This bird occurs at most Himalayan stations.

149. *Neophron percnopterus*: The Large White Scavenger Vulture. F. 1198, J. 6, +V, about the size of a kite. W. K.

Plumage sullied white, except tips of wings which are shabby black. Bare skin of head yellow. Neck sparsely covered with sullied pale yellow feathers, which do not lie flat. Legs yellow. Bill grey.

As this vulture waddles along it is a revolting-looking bird. When sailing through the air on outstretched wings, this vulture looks well, and the wings are then seen to be white margined with black. The end of the wing is rounded and not cut up into fingers like that of the wing of the white-backed vulture.

This bird is fairly common in the Kashmir valley and at most Western Himalayan stations.

150. *Gypaëtus barbatus*: The Lammergeyer

---

---

## *Descriptive List of Birds*

---

or Bearded Vulture. F. 1199, J. 7, +V, 4 feet long. W. K.

This bird is often mistaken for the Golden Eagle by people at hill stations.

Head and nape white. Rest of upper plumage grey. Lower plumage pale brown. A black band from the eye to the bill ending in the "beard," which is a tuft of black bristles growing from the lower mandible. Tail long and rounded at end; that of the scavenger vulture being diamond-shaped.

This great bird is fairly common in the Kashmir valley, Simla, Dharmsala and Mussoorie.

### *Birds of Prey*

This large family is composed of birds which bear so strong a family likeness that it is almost impossible to describe them in such a way as to enable the reader to identify them at sight. As with owls, birds of prey are easily recognised as such, but to name any particular bird soaring overhead often baffles expert ornithologists. To try to make out birds of prey by their colouring is a hopeless task. Eha remarks, "every bird of prey has its own character, some trick of flight, something in its figure

---

---

## *Himalayan and Kashmiri Birds*

---

and proportions which serves to distinguish it decisively." There is much truth in this; but it is not easy to state what this is. Mr. C. H. Donald, in Vols. XXV, XXVI, and XXXVII of the *Journal of the Bombay Natural History Society*, has given excellent descriptions of the mannerisms of the birds of prey. In what follows I have drawn freely on Mr. Donald, who probably knows more about birds of prey in India than anyone else, having made a special study of the subject.

151. *Spizaëtus nepalensis*: Hodgson's Hawk-Eagle. F. 1213, J. 37, +V. E. W. K.

A small crest. General hue pale brown, mottled with white. Three black bands separated by two white ones run down the throat. Under surface of tail pale grey. Upper surface of tail has broad black cross-bars.

The call of this bird is a shrill musical whistle.

Donald writes: "Unmistakable in flight. Wings very short and round, held very far back and the tail projecting far beyond the line of the wings. . . . Soars well and is often seen to give several short sharp flaps after a bout of circling on steady pinions. Usually found over heavy pine or oak forests."

---

---

## *Descriptive List of Birds*

---

Widely distributed, but not very common anywhere.

152. *Spilornis cheela* : The Crested Serpent-Eagle. F. 1217, J. 39, +V. E. W.

Black crest. Dark chocolate-brown plumage, with white or cream bars on wings and tail. Lower plumage spotted with large white spots.

Donald writes : " When soaring the wings are held well back, but are broad in proportion to his size, hence the tail does not project very far beyond the tertiaries (end wing feathers). This is a noisy bird and gives vent to a shrill succession of whistles when on the wing. Found on the lower hills amongst pine and oak."

This bird does not usually ascend much above 6000 feet but is occasionally seen at the various hill stations.

153. *Haliaëtus leucoryphus* : Pallas's Fishing-Eagle. F. 1223, J. 42, +V, nearly half as big again as the Kite. E. W. K.

Head, chin and throat white. A white band across the tail about 3 inches from the tip. Rest of plumage brown. The white band across the tail is conspicuous, and Jerdon styles this bird the " Ring-tailed Sea-Eagle."

---

---

## *Himalayan and Kashmiri Birds*

---

Tail does not project much beyond wings during flight. Extremely noisy.

This bird frequents rivers and marshes and is common in Kashmir on the lakes and the Jhelum.

154. *Milvus govinda*: The Common Pariah Kite. F. 1229, J. 56, V. E. W. K.

A dark-brown bird with rather a long forked tail, which serves to differentiate it from other birds of prey.

Donald writes, "Flight is easy and light, but irregular and frequently changing direction as though the bird is uncertain which line it should take. The wings are held in the same plane as the body and are frequently slightly bent as though the bird were meditating a stoop."

Very abundant at Almora, Simla, Dharm-sala and in the Kashmir valley, where it nests in chenar and walnut trees.

It utters when sailing high in the air a mournful-sounding *Chee-he-he-he-he-he*.

155. *Falco subbuteo*: The Hobby. F. 1260, J. 13, +III. E. W.

Head black with pale eyebrow and forehead. Rest of upper plumage dark slaty-grey. Under surface tinged with red on the abdomen and



---

---

## *Descriptive List of Birds*

---

thighs. Some brown streaks on the lower plumage.

Donald writes: "This beautiful little falcon is by no means rare in the Himalayas and its wonderful evolutions in the air cannot help attracting attention. Its long pointed wings make it appear bigger than it really is. . . . If watched for a few seconds it will be seen constantly to change direction and turn and twist in the air in a most amazing way in pursuit of insects on which it mostly preys. The Hobby does not usually make its appearance till late in the afternoon and may be seen circling, stooping, rising vertically, and playing extraordinary tricks in the air, some time after all diurnal birds have gone to rest."

This bird is nowhere very common.

156. *Tinnunculus alaudarius*: The Kestrel. F. 1265, J. 17, +III. E. W. K.

This is the English "Windhover."

Head, neck and tail grey. Back and wings brick-red. Lower parts cream-coloured spotted with brown. The red wings, grey body and manner in which it hovers in mid-air on rapidly-vibrating wings render the identification of this pretty little bird an easy matter. The tail is graduated, the middle feathers

---

---

## *Himalayan and Kashmiri Birds*

---

being the longest. The tip of the tail is white, with a dark broad band above the white.

Next to the Kite the Kestrel is the commonest bird of prey at the various hill stations.

### *Green Pigeons*

Green pigeons are strictly arboreal in their habits, and their plumage assimilates so well with the foliage that it is difficult to see them.

157. *Sphenocercus sphenurus*: The Kokla Green Pigeon. F. 1283, J. 778, +III. E. W. K.

Head and lower plumage yellowish-green. Upper back grey. Middle back maroon (in cock only). Rest of plumage olive green. Some black and yellow in wing. Under surface of wings and tail grey. Upper breast is tinted with pink.

The call of this pigeon is very curious. I do not find it very agreeable; but Indians cage it for the sake of its "song." The name Kokla is onomatopoeic, but does not in my opinion represent the call. "The notes," writes Blyth, "bear some resemblance to the human voice in singing and are highly musical in tone, being considerably prolonged and modulated, but always terminating abruptly, and

---

---

## *Descriptive List of Birds*

---

every time the stave is repeated exactly as before, so that it soon becomes wearisome to an European ear.”

This bird occurs in Kashmir and is common at all hill stations.

### *Blue Rock-Pigeons*

All the domestic varieties of pigeon are descended from Blue Rock-pigeons. These birds are gregarious and usually nest on rocks or buildings, hence the name.

158. *Columba intermedia*: The Indian Blue Rock-Pigeon. F. 1292, J. 788, +III. E. W. K.

This is the common pigeon of the plains. It is not nearly so abundant in Kashmir or the Himalayas as it is in the plains of India. Except at Dharmsala it does not usually occur at elevations over 5000 feet.

A slaty-blue bird with a metallic green or purple gloss on the neck. Two black bars across the wing. The tail is tipped with black. Legs reddish-pink.

### *Doves*

Doves are small pigeons, the plumage of which is of the sandy hue known as dove-colour.

---

## *Himalayan and Kashmiri Birds*

---

159. *Turtur ferrago*: The Indian Turtle-Dove. F. 1305, J. 792, III. W. K. .

This has the usual dove-coloured plumage, with a miniature black-and-grey chess-board on each side of the neck.

This is the commonest dove of the West Himalayan hill-stations.

160. *Turtur cambayensis*: The Little Brown Dove. F. 1309, J. 794, -III. W.

Plumage sand-coloured, greyish on wings and tail. Head, neck and breast suffused with dull pale red. A miniature red-and-black chess-board on each side of neck. Legs crimson-lake.

The call of this bird is a subdued *cuk-cuk-coo-coo-coo*.

This is the common dove of the plains. The only hill station at which I have seen it is Almora.

It, together with the spotted-dove (*Turtur suratensis*) and the Indian Ring-Dove (*Turtur risorius*), is common on the cart road between Rawalpindi and Kohala up to elevations of 5000 feet.

161. *Turtur risorius*: The Indian Ring-Dove. F. 1310, J. 796, +III. W. K.

A pale French-grey bird, with a half collar round the back of the neck. This collar is

---

---

## *Descriptive List of Birds*

---

black, narrowly bordered with white above and below. The call of this bird is *ku-ku-ku*. There is a considerable hiatus between the second and the third *ku*.

This is the common dove of Srinagar. At Simla it occurs up to 6000 feet, as does the Spotted Dove (*Turtur suratensis*). This last, which occurs at Dharmsala up to 5000 feet, may be recognised by the black-and-white chess-board on each side of the neck, and by the brown back having reddish spots on it. This and the other doves have a little white in the tail.

### *Pheasants*

Pheasants are well represented in the Himalayas. The monal, horned pheasants, cheer, pukras and kalij are all found in those mountains, and the monal and the pukras in Kashmir.

The monal, horned pheasants and pukras usually occur in summer at higher elevations than the hill stations. The cheer is nowhere common. The common pheasant of the various hill stations is one or other of the kalij species.

162. *Gennæus albicristatus*: The White-crested Kalij Pheasant. F. 1336, J. 810, + V. W.

---

---

## *Himalayan and Kashmiri Birds*

---

In shape more like a fowl than a pheasant.

*Cock*: Long white hairy crest. Upper plumage black glossed with blue, with white bars on lower back. Wings and fowl-like tail dark brown glossed with green.

*Hen*: Crest and upper plumage reddish-brown with fine black wavy crossbars. Middle tail feathers black, with cream wavy crossbars; the other tail feathers are black with a blue gloss.

Both sexes have a patch of bare red skin on sides of face. This bird is fond of feeding on the road and I have shot it from a motor-car while going along the road to Chakrata.

163. *Gennæus melanonotus*: The Black-backed Kalij Pheasant. F. 1338, J. 811, V. E.

The common Kalij of Darjeeling.

Like the last species, but the cock has a black crest, glossed with purple-blue, and there are no white bars on the lower back and rump.

### *Note on the other Himalayan Pheasants*

*Catreus wallichi*: The Cheer. This bird has a long tail like that of the English Pheasant. Both sexes have a long crest and a red patch of bare skin round the eye. Plumage black with

---

---

## *Descriptive List of Birds*

---

white and reddish-yellow crossbars. Found in Western Himalayas.

*Pucrasia macrolopha* : The Koklas or Pukras Pheasant.

Cock has three crests, the middle one fawn-coloured and the side ones black glossed with green. A long-tailed pheasant. Found in summer at high elevations in Kashmir and Western Himalayas.

*Lophophorus refulgens* : The Monal.

*Cock* : Brilliant metallic-purple, with green peacock-like crest. Short tail cinnamon.

*Hen* : Brown.

This bird occurs at high elevations in Kashmir and West Himalayas.

*Tragopan satyra* : The Crimson Horned Pheasant. (E. Himalayas.)

*Cock* : Black head (and breast in western race). Rest of plumage mainly crimson with white or grey spots. Tail long.

*Hen* : Plumage mainly brown.

*Tragopan melanocephalus* : The Western Horned Pheasant (W. Himalayas).

*Cock* : Black head and breast. Rest of plumage mainly crimson with white or grey spots. Tail long.

*Hen* : Plumage mainly brown.

---

---

## *Himalayan and Kashmiri Birds*

---

### *Quail*

Quail are partridges in miniature.

164. *Coturnix communis*: The Common Grey Quail. F. 1355, J. 829, +II. E. W.

A little short-tailed, short-legged bird. Upper plumage brown spotted with dark brown and buff. Lower plumage reddish-buff with dark streaks and spots.

Only passes through the Himalayas on migration in April and October. Many are shot while the spring crops are being harvested at the various hill stations.

### *Partridges*

165. *Gaccabis chucar*: The Chukor. F. 1370, J. 820, -IV. W. K.

Has red legs like the French partridge which it resembles in appearance. General colour of plumage pale reddish-brown. A black band runs across the forehead to the eyes and from there down the sides of the head across the throat to form a gorget. Some conspicuous black bars on the flanks.

Common about the West Himalayan hill stations and on the Kashmir hills.



---

---

## *Descriptive List of Birds*

---

166. *Francolinus vulgaris*: The Black Partridge. F. 1372, J. 818, +III. W. K.

*Cock*: A handsome black bird with white spots on the flanks and narrow white bars on the back. Crown and wings buff and dark brown. White patch on each side of head. Chestnut collar round neck.

*Hen*: Plumage buff and dark brown.

Call so high-pitched that some people cannot bear it. It may be syllabised *juk-juk-tee-tee-tur*.

This partridge does not ascend the Himalayas above 5000 in most parts, but it is fairly common at Simla and Mussoorie and on the Kashmir hills.

### *Water-Rails*

Water-Rails are land birds which have taken to living in marshes, and so have acquired some of the features of ducks.

167. *Gallinula chloropus*: The Moorhen. F. 1402, J. 905, +III. K.

This is the Moorhen of England.

Head and breast dark slaty-grey. Back and shoulders olive-brown. Rump brown. Wings blackish-brown, with a little white in them. Tail black. Some white bars on flanks and

---

---

## *Himalayan and Kashmiri Birds*

---

a white patch under the tail. A red shield on forehead.

Common on the Kashmir lakes.

168. *Fulica atra*: The Coot. F. 1405, J. 903, IV. E. W. K.

As this bird lives on lakes and, except in Kashmir, does not ascend the hills above 4000 feet, it is not seen at any hill station, but occurs on some of the lakes of the Kashmir valley. It used to occur on the Dal Lake, but I do not think it does now.

A black duck-like bird, with a white shield on the forehead. It swims like a duck; but when flushed from off the water, it runs along the surface, flapping its wings vigorously, before it succeeds in rising above the water. When flying it may be likened to a bottle of Bass on wings!

### *Jaçanas, Lily-trotters, or Lotus-birds*

Jaçanas are birds that live on lakes in which lotus and other water-plants grow profusely. These birds have very long toes, which enable them to run about on the broad flat floating leaves of the water-plants.

169. *Hydrophasianus chirurgus*: The Pheas-

---

---

## *Descriptive List of Birds*

---

ant-tailed Jaçana. F. 1429, J. 901. + III, with tail 10 inches long in summer. K.

Head, throat and wings white. Back of neck golden yellow. Long, thin, black pheasant-like tail. This bird swims like a duck and runs about with great address on floating water-plants. It utters a cat-like mew. On account of its pheasant-like appearance, this bird is sometimes called the water pheasant. It is a very beautiful bird. It is impossible to mistake it. It is very abundant on the Dal and the other lakes of the Kashmir valley.

### *Plovers, or Lapwings*

Plovers are short-tailed, long-legged birds that appear to be adapted to wading but which keep to dry land. They have peculiar plaintive calls and many of them perform wonderful antics in the air.

170. *Sarcogrammus indicus*: The Red-wattled Lapwing. F. 1431, J. 855, +IV. E. W. K.

This is the familiar "Did-you-do-it" of the plains.

Head, neck and upper breast black. A broad white band running from the eye down the whole length of the neck. Back and wings, bronze-brown, black and white; the white

---

---

## *Himalayan and Kashmiri Birds*

---

taking the form of a conspicuous band during flight. Lower parts white. Tail white with black band across it near the tip. Bill red, with a crimson wattle. Legs bright yellow.

This bird ascends the Himalayas up to 5000 feet and occurs in the Kashmir valley.

### *Terns*

Terns are allied to gulls. They are long-winged, slenderly-built birds, that have much white and grey in the plumage. They feed largely on fish and spend much time in flying over water in search of their quarry.

171. *Hydrochelidon hybrida*: The Whiskered Tern. F. 1496, J. 984, +II. K.

Bill red. Cap black. Upper plumage silvery-grey. Breast grey. Abdomen black.

This bird occurs abundantly on the lakes in the Kashmir valley. It makes a floating nest of water-weeds in the early summer.

It sometimes hunts for insects over fields.

### *Cormorants*

Cormorants are large black swimming birds, which feed on fish. They catch their quarry by diving. When not fishing they sometimes

---

---

## *Descriptive List of Birds*

---

squat on a post projecting out of the water and spread out their wings to dry them.

172. *Phalacrocorax carbo*: The Large Cormorant. F. 1526, J. 1005, +V. K.

Plumage all black save for a white band from eye to eye across the throat.

Found on the lakes of the Kashmir valley and on the Jhelum.

173. *Phalacrocorax fuscicollis*: The Indian Shag. F. 1527, J. 1006, +V. K.

Distinguished from the last species by having no white band across the throat and by its smaller size.

Found on the lakes of the Kashmir valley and on the Jhelum.

### *Hérons*

Hérons are wading birds with long stiletto-like bills. It is their habit to stand motionless in shallow water. When they espy a fish or a frog, the neck shoots out like a telescope and the bill is plunged into the water to secure the victim. On the wing herons are easily distinguished from other long-shanked birds by the fact that, although they, in common with other wading birds, carry the legs stiffly stretched out behind them, the head is drawn

---

---

## *Himalayan and Kashmiri Birds*

---

in, so that the neck projects like the crop of a pouter pigeon.

174. *Ardea cinerea* : The Common Heron. F. 1555, J. 923, +V, over a yard long. K.

This is the common heron of England.

Upper parts ashy-grey. Some white on the head. Black plumes hang down from the nape. Lower parts white.

Common on the margins of the lakes of the Kashmir valley.

175. *Nycticorax griseus* : The Night Heron. F. 1568, J. 937, V. K.

Head, nape, back and shoulders black. Forehead, cheeks and lower parts white. Rest of plumage grey. Some feathers of the nape are white and are lengthened to form plumes. Eyes bright red.

This bird fishes chiefly by night. About sunset it leaves its roost in some tall tree and slowly flaps its way to its fishing ground, uttering a loud *wāk*.

Common round the lakes in the Kashmir valley.

A company of these birds roosts, along with cormorants, in some tall trees dear the Nasim Bagh on the Dal Lake.

---

---

## *Descriptive List of Birds*

---

176. *Ardetta minuta* : The Little Bittern. F. 1570, J. 935, —IV. K.

Upper parts buff glossed with green. Breast buff. Some buff in the wings. Legs greenish-yellow. This little bird, which the Kashmiris call *Goi*, is very common on the Kashmir lakes. During the daytime it lies up hidden in the reeds. After sunset the little bitterns emerge and take up fishing stations in the shallow water of the lakes and waterways. Dozens of these little herons may be seen by anyone out in a *shikara* on the Dal Lake just before nightfall.

### *Ducks*

Two species of duck breed on the lakes in the vale of Kashmir ; but, owing to their eggs being collected for the Srinagar market, they are now rarely seen on the Dal Lake. They occur on the other lakes.

177. *Anas boscas* : The Mallard or Wild Duck. F. 1592, J. 958, V. K.

The Wild Duck of England.

*Male* : September to May. Head glistening emerald-green, with white collar round neck. Breast chestnut. Shoulders finely barred white and brown. Middle back brown. Lower back and middle tail feathers black glossed

---

---

## *Himalayan and Kashmiri Birds*

---

with purple or green. Middle tail feathers curl upwards. Wings brown with metallic-purple bar. Below tail velvety black.

*Male* (June to August) and *Female*: Brown with buff crescent-shaped markings. Purple wing-bar.

Legs orange-yellow in both sexes.

178. *Nyroca ferruginea*: The White-eyed Pochard. F. 1606, J. 969, —IV. K.

Head, neck and breast dull chestnut. Rest of upper plumage blackish-brown. Tail dark brown. Wings dark brown and white. Abdomen chiefly white. Hen like cock, but duller.

### *Grebes*

Grebes are small duck-like water birds of which the tail is very short. They rarely come to land, they dive for their food.

179. *Podiceps albipennis*: The Indian Little Grebe, or Dabchick. F. 1617, J. 975, +II. K.

Body rich dark brown. Chestnut on sides of neck. Looks like a tiny tailless duck. Dives every few seconds. Very common on the Kashmir lakes.



---

## INDEX

---

- Babbler, black-throated, 106  
— Nepal, 106  
— red-headed, 107  
— rusty-cheeked scimitar,  
106  
Barbet, golden-throated, 160  
— great Himalayan, 160  
Bee-eater, European, 162  
Bittern, little, 195  
Bulbul, Bengal red-vented,  
113  
— Himalayan black, 112  
— Punjab red-vented, 114  
— striated green, 113  
— white-cheeked, 114  
Bunting, Eastern meadow,  
148  
— white-capped, 147  
  
Chat, dark grey bush, 134  
— Indian bush, 133  
— pied, 133  
Cheer pheasant, 186  
Chukor, 188  
Coot, 190  
Cormorant, large, 193  
Crow, house, 94  
— jungle, 94  
Cuckoo, 166  
Cuckoo, Himalayan, 166  
— Indian, 167  
— large hawk, 168  
— pied crested, 169  
— small, 167  
  
Dabchick, Indian, 196  
Dipper, brown, 115  
Dove, little brown, 184  
— ring, 184  
— spotted, 185  
— turtle, 184  
Drongo, ashy, 116  
— black, 115  
Duck, wild, 195  
  
Eagle, crested serpent, 179  
— Hodgson's hawk, 178  
— Pallas's fishing, 179  
  
Finch, Himalayan gold, 144  
— Himalayan green, 145  
— red-crowned, 144  
— red-mantled rose, 143  
— scarlet, 143  
Firecap, 110  
Flowerpecker, fire-breasted,  
155

---

---

## *Index*

---

Flycatcher, brown, 129  
— ferruginous, 128  
— grey-headed, 130  
— Indian paradise, 132  
— rufous tailed, 130  
— sooty, 127  
— verditer, 129  
— white-browed, 128  
Forktail, Eastern, 135  
— little, 136  
— Western, 134  
  
Grebe, little, 196  
Grosbeak, black-and-yellow,  
    142  
  
Heron, common, 194  
— night, 194  
Hobby, 180  
Hoopoe, European, 164  
  
Ixulus, yellow-naped, 109  
  
Jaçana, pheasant-tailed, 196  
Jackdaw, 94  
Jay, black-throated, 96  
— Himalayan, 97  
  
Kalij Pheasant, black-backed,  
    186  
— white-crested, 185  
Kestrel, 181  
Kingcrow, 115  
Kingfisher, common, 163  
— pied crested, 162  
Kite, common pariah, 180

Koel, Indian, 169  
Kokla, green pigeon, 182  
  
Lammergeyer, 176  
Lapwing, red-wattled, 191  
Liothrix, red-billed, 109  
  
Magpie, red-billed blue, 95  
— yellow-billed blue, 95  
Mallard, 195  
Minivet, short-billed, 123  
Minla, red-tailed, 110  
Monal Pheasant, 187  
Moorhen, 189  
Myna, common, 125  
— grey-headed, 126  
— jungle, 127  
  
Niltava, large, 133  
— rufous-bellied, 132  
Nutcracker, Himalayan, 98  
— large spotted, 98  
Nuthatch, white-tailed, 115  
  
Oriole, Indian, 124  
Ouzel, grey-winged, 140  
— Tickell's, 140  
Owl, Himalayan wood, 172  
— spotted Himalayan scops,  
    173  
Owlet, collared pigmy, 174  
— large barred, 173  
  
Paroquet, large Indian, 170  
— slaty-headed, 171  
— Western blossom-headed,  
    171  
Partridge, black, 189

---

---

## *Index*

---

- Partridge, chukor, 188  
Pie, Himalayan tree, 96  
Pigeon, blue rock, 183  
— kokla green, 182  
Pipit, brown rock, 152  
— Indian, 152  
— Indian tree, 151  
— upland, 152  
Pochard, white-eyed, 196  
  
Quail, common grey, 188  
  
Redstart, blue-fronted, 137  
— plumbeous, 138  
— white-capped, 136  
Robin, magpie, 139  
— Pekin, 109  
— red-flanked bush, 138  
Roller, European, 161  
  
Shag, 193  
Shrike, cuckoo, 124  
— grey-backed, 123  
— rufous-backed, 122  
Sibia, black-headed, 108  
Skylark, 155  
Sparrow, cinnamon tree, 146  
— house, 145  
— tree, 146  
Spinetail, white-necked, 165  
Spotted-Wing, 111  
Starling, Himalayan, 125  
Sunbird, fire-tailed yellow-  
backed, 154  
— Himalayan yellow-backed,  
155  
Sunbird, Nepal yellow-  
backed, 154  
Swallow, 148  
— Sykes's striated, 149  
— wire-tailed, 149  
Swift, Alpine, 165  
— Indian, 165  
  
Tern, whiskered, 192  
Tit, brown crested, 101  
— crested black, 100  
— green-backed, 99  
— grey, 99  
— red-headed, 99  
— yellow-cheeked, 100  
Thrush, Eastern variegated  
laughing, 104  
— Himalayan streaked  
laughing, 104  
— Himalayan whistling, 107  
— Himalayan white-crested  
laughing, 101  
— red-headed laughing, 103  
— rufous chinned laughing,  
102  
— striated laughing, 105  
— Western yellow-winged  
laughing, 103  
— white-throated laughing,  
102  
Tree-creeper, Himalayan,  
117  
— Sikkim, 117  
  
Vulture, black, 175  
— Himalayan griffon, 175

---

---

## *Index*

---

Vulture, Indian white-backed, 175  
— large white scavenger, 176

Wagtail, grey, 150  
— Hodgson's pied, 150  
— Hodgson's yellow-headed, 151

Warbler, Brook's grey-headed flycatcher, 121  
— Brown hill, 121  
— Hodgson's grey-headed flycatcher, 120  
— Hume's willow, 119  
— Indian great reed, 119  
— Indian lesser white-throated, 119

White-eye, Indian, 108

Woodpecker, black-naped green, 157  
— brown - fronted pied, 158  
— lesser pied, 158  
— Northern pale-headed, 157  
— red-eared bay, 159  
— rufous - bellied pied, 157  
— West Himalayan scaly-bellied green, 156  
— Western Himalayan pied, 158

Wren, Himalayan, 118  
— scaly-breasted, 118

Wryneck, 159

BY THE SAME AUTHOR

# Indian Birds

Being a Key to the Common Birds of the Plains of India.

Crown 8vo. 6s. net.

*Spectator*.—"A charming writer whose earlier books on birds we have had the pleasure of recommending. The book is most carefully compiled and much ingenuity is displayed in framing this artificial analysis."

*Globe*.—"Mr. Dewar is not only a keen and patient observer, but he is gifted with the descriptive art in a high degree, and his vivacious style communicates the characters and habits of birds with unerring fidelity and infinite spirit."

*Daily Chronicle*.—"Here is a work worthy of all commendation to those who love birds."

*Pall Mall Gazette*.—"Mr. Dewar's volume is one of the best examples of sound information conveyed in attractive literary form."

# Birds of the Plains

With numerous Illustrations.

Demy 8vo. 10s. 6d. net.

*Globe*.—"Mr. Dewar is not only a keen and patient observer, but he is gifted with the descriptive art in a high degree, and his vivacious style communicates the characters and habits of birds with unerring fidelity and infinite spirit."

*Sportsman*.—"Mr. Dewar has a delightfully simple and quaintly humorous way of expressing himself, and his clever word pictures of bird life make charming reading."

John Lane The Bodley Head Ltd., Vigo Street, W. 1

BY THE SAME AUTHOR

## Jungle Folk

Indian Natural History Sketches.

Demy 8vo. 10s. 6d. net.

*World.*—“Full of information, it is written in a way that sounds wonderfully clever, and yet it is not over the head of the general reader. An unconventional, fresh and charming exposition.”

## Bombay Ducks

An Account of some of the everyday Birds and Beasts found in a Naturalist's El Dorado.

With numerous Illustrations from Photographs of Living Birds.

Demy 8vo. 16s. net.

*Daily News.*—“This new and sumptuous book, Mr. Dewar gives us a charming introduction to a great many interesting birds.”

## Glimpses of Indian Birds

Demy 8vo. 7s. 6d. net.

By DOUGLAS DEWAR AND FRANK FINN

## The Making of Species

With numerous Illustrations.

Demy 8vo. 7s. 6d. net.

*Nation.*—“An exceptionally interesting book. We gladly welcome two such able and independent naturalists.”

*Outlook.*—“A book of knowledge and originality.”

*Daily Telegraph.*—“Interesting and suggestive. It should receive wide attention.”

John Lane The Bodley Head Ltd., Vigo Street, W. 1

# Companions Feathered, Furred, and Scaled

By C. H. DONALD, F.Z.S., M.B.O.U.

Illustrated. Crown 8vo. 7s. net.

In this book Mr. DONALD, the well-known naturalist, describes the various wild pets acquired by himself and his wife during their jungle-camping trips. They make a varied and interesting collection, for Mr. DONALD has a catholic taste in his animal friendships, and is equally at home with his golden eagles, vultures, cormorants, weavers, and his red flying squirrel tippitty. He has illustrated the book himself from his photographs and drawings.

“His book is a bright and humorous account of the habits of his strange friends.”—*Daily News*.

“Full of entertaining stories and photographs. A capital book for animal lovers.”—*Star*.

“These chatty sketches by a keen and courageous naturalist.”  
*The Times*.

“Many of the incidents related are most amusing, and are very well told.”—*Field*.

## In Nature's Garden

By C. H. DONALD, F.Z.S.

Illustrated. Crown 8vo. 7s. 6d. net.

Vivid and intimate studies of animal life and character in Northern India, by a well-known naturalist.

JOHN LANE THE BODLEY HEAD LTD., VIGO ST., W.1

# SONGS OF THE BIRDS

By

WALTER GARSTANG, M.A., D.Sc.,

PROFESSOR OF ZOOLOGY IN THE UNIVERSITY OF LEEDS

Illustrated by J. A. SHEPHERD

Second Edition. Crown 8vo. 6s. net.

A description of the origin and development of music from the songs of birds, with reproductions of the songs set to human consonants and music.

“ Full of original remarks and good observation. Dr. Garstang has enjoyed writing his book, and has felt what he says before he has written it.”—VISCOUNTESS GREY (in a two-column review in the *Times*.)

“ A triumph of loving and careful observation. A delightful book, delightfully illustrated.”—*Morning Post*.

“ One of the most delicious books. Every lover of the country should read it. Altogether a book to love and a book to keep.”—*British Weekly*.

“ After studying this charming book the most ignorant listener of the countryside should be able to recognize the individuals in the feathered choir.—*John O'London's Weekly*.

---

JOHN LANE THE BODLEY HEAD LTD., VIGO ST., W.1



# The Civilian's South India

By "CIVILIAN"

Demy 8vo. 12s. 6d. net.

*Graphic.*—"A remarkably full and accurate impression of the country and its people."

*Truth.*—"An extremely bright and amusing book. Deliciously satirical, always humorous, never vicious, this is quite the most diverting book of its kind that has appeared for a long while."

*Court Journal.*—"The author's outlook on life in South India is so unconventional, and, with his sense of humour as well as obvious sincerity, should make many desire an early acquaintance with his fascinating work."

# The Forests of India

By E. P. STEBBING

Professor of Forestry, Edinburgh University; Author of "Stalks in the Himalayas"; "The Diary of a Sportsman Naturalist," etc.

Part I. The Early History of the Forests of India.

Part II. The Position and Treatment of the Forests of India.

Illustrated Demy 8vo. £2 2s. net. each volume.

# Topee and Turban, or here and there in India

By LIEUT.-COLONEL H. A. NEWELL, I.A.

With Illustrations from Photographs. Demy 8vo.

£1 1s. net.

*Daily News.*—"An excellent book. . . . Colonel Newell has a quick eye for things seen, and a happy interest in history that makes the book something more than a motorist guide-book. . . . Fully illustrated with photographs of great interest and variety."

*Daily Mail.*—"Everyone ought to know more about India, and a pleasant way of informing oneself is offered by Colonel Newell's 'Topee and Turban in India.'"

*Westminster Gazette.*—"In this book Colonel Newell shows that he has been a close student and observer of many things in India and Indian life."

John Lane The Bodley Head Ltd., Vigo Street, W. 1

# Rifle and Romance in the Indian Jungle

Being the Record of Thirteen Years of Indian Jungle Life.

By MAJOR A. I. GLASFURD  
(*Indian Army*)

With numerous Illustrations by the Author, and Reproductions from Photographs. Crown 8vo. 12s. 6d. net.

*Literary World.*—"To the list of books on big game shooting that can be recommended equally to the sportsman and the general reader must be added this truly fascinating work. We have read it through from cover to cover and pronounce it excellent."

# Days and Nights of Shikar

By MRS. W. W. BAILLIE

Demy 8vo. 12s. 6d. net.

Mrs. Baillie here relates the story of her wanderings in quest of big game, and her thrilling encounters with all kinds of wild beasts in the Indian Jungle.

*Evening Standard.*—"Mrs. Baillie writes in an easy, unaffected style, and for simple disregard of danger, her accounts of her escapes from danger beats anything I have ever read from a feminine pen."

# The Diary of a Sportsman Naturalist in India

By E. P. STEBBING

Profusely Illustrated from Photographs and Sketches by the Author. Demy 8vo. £1 1s. net.

*Punch.*—"Dip where you may you will find none of the anecdotes trivial or tiresome. To sportsmen this volume will be extremely welcome."

*Daily Chronicle.*—"A book with more thrills than many a novel. One is entranced with these tales of jungle adventure and jungle lore."

*New Statesman.*—"We do not remember a sporting book which gives a better idea of what Indian jungles are to the sportsman than this. Mr. Stebbing has a vast experience of different Indian jungles."

John Lane The Bodley Head Ltd., Vigo Street, W. 1

# THE BODLEY HEAD NATURAL HISTORY

By E. D. CUMING

WITH ILLUSTRATIONS

By J. A. SHEPHERD

Square 8vo ( $6\frac{3}{4} \times 5\frac{1}{4}$ ). 2s. 6d. net

VOL. I: Thrush, Blackbird, Ouzel, Redwing, Fieldfare, Wheatear, Robin, Nightingale, Winchat, Stonechat, Tits, Starling, Wren, etc.

VOL. II: Whitethroat, Crested Wren, Wood Wren, Warblers, Accentor, Dipper, Nuthatch, Tree-creeper, etc.

*Morning Post*.—"The illustrations give them a vivacity and humour which make them much more typical and suggestive than many elaborate plates we have seen."

*Outlook*.—"The descriptions of the birds and their habits are brief but very clear, and the illustrations in which characteristic attitudes and colour are indicated in impressionist studies will prove far more useful for purposes of identification than more elaborate drawings in which no movement can be seen or imagined. Very charming books."

*Guardian*.—"Here is a clever book, very unconventional in its text, and even more so in its clever illustrations, which bids fair to inaugurate a new era in popular natural history. Prettily and neatly produced."

*Tatler*.—"Such a delightful book. Bird pictures everybody will love. As an animal drawer Mr. Shepherd is incomparable."

*Graphic*.—"A great success."

*Bookman*.—"A concise and eminently well-informed work . . . delightfully readable."

---

John Lane The Bodley Head Ltd., Vigo Street, W. 1.

# BOOKS ABOUT MALAYA

BY SIR FRANK SWETTENHAM, K.C.M.G.

## BRITISH MALAYA

An Account of the Origin and Progress of British Influence in Malaya. Numerous Illustrations reproduced from Photographs.

Fourth Edition. Crown 8vo. 12s. 6d. net.

"The book is richly illustrated with photographs of great beauty. It takes its place as the standard work of reference on the ruder tribes of the peninsula."—*Daily News*.

## THE REAL MALAY

Pen Pictures.

Second Edition. Crown 8vo. 7s. net.

"Sir Frank Swettenham understands perhaps better than any other roving Englishman 'The Real Malay.'"—*Standard*.

## MALAY SKETCHES

With title-page and cover designed by  
**PATTEN WILSON**

Fourth Edition. Crown 8vo. 7s. net.

"Nothing approaching Sir Frank Swettenham's intimate knowledge and illuminative analysis has yet seen light about that fascinating country which he so well describes."—*Daily Chronicle*.

## UNADDRESSED LETTERS

With Portrait of the Author.

Seventh Edition. Crown 8vo. 7s. net.

"'Unaddressed Letters' is the revelation of a personality at once sympathetic and distinguished, imaginative, sensitive, emotional."—*Daily Chronicle*.

## ALSO AND PERHAPS

Second Edition. Illustrated. Crown 8vo. 7s. net.

"When one has reached the end of the book, one undoubtedly has a strong sense of having been in the company of a man of extraordinary wide experience and of a devouring curiosity allied with sympathy."—*Spectator*.

JOHN LANE THE BODLEY HEAD Ltd., Vigo St., W.1

