

Flora of India
Series - 2

FLORA OF SIKKIM

Volume 1
(Monocotyledons)

Editors
P.K. HAJRA
D.M. VERMA

with assistance from
S. Bandyopadhyaya



BOTANICAL SURVEY OF INDIA


Flora of India
Series - 2

FLORA OF SIKKIM

Volume 1
(Monocotyledons)

Editors
P.K. HAJRA
D.M. VERMA

with assistance from
S. Bandyopadhyay



BOTANICAL SURVEY OF INDIA

© Government of India
Date of Publication, December 1996

No part of this publication can be reproduced, stored in a retrieval system, or transmitted, in any form or means by electronic, mechanical, photocopying, recording, or otherwise, without prior written permission of the Director, Botanical Survey of India.

Cover Photo : A view of Khangchendzong from Thang sing (West Sikkim)

Price :

Published by the Director, Botanical Survey of India, P-8, Brabourne Road, Calcutta-700001 and printed at Deep Printers, 3/26, Ramesh Nagar, New Delhi-110015

Contents

Acknowledgements

Contributors

Introduction 1

Selected References 15

Enumeration of the Species 23

Index to Botanical Names 307

Acknowledgements

It gives the editors great pleasure to record their thanks to all the contributors.

They also express their thanks to Shri Utpal Chatterjee, Scientist, and all staff members of Publication Unit, Botanical Survey of India for their help.

Finally the editors acknowledge Shri M.L. Jain of Deep Printers for printing of this volume.

Contributors

BASU, P. Botanical Survey of India, P-8, Brabourne Road, Calcutta - 700 001.

KRISHNA, B. Shiv Niketan, M.P. Sinha Road, Kadamkuam, Patna - 800 003.

KUMAR, S. Botanical Survey of India, Northern Circle, 192, Kaulagarh Road, Dehra Dun - 248 195.

MANDAL, N.R. Botanical Survey of India, Central National Herbarium, P.O. Botanic Garden, Howrah - 711 103.

MITRA, B. Botanical Survey of India, Central Botanical Laboratory, P.O. Botanic Garden, Howrah - 711 103.

RAJU, D.C.S. "Lumbini", 7th Road, Banjara Hills, Hyderabad - 500 034.

SINGH, PARAMJIT. Botanical Survey of India, Sikkim, Himalayan Circle, Sikkim.

SINGH, S. Remote Sensing Institute, Kalidas Road, Dehra Dun.

SRIVASTAVA, R.C. Botanical Survey of India, Central Circle, 10 Chatham Line, Allahabad - 211 002.



Hedychium spicatum Buch. - Ham. ex J.E. Smith



Pandanus nepalensis St. John



Maimaichu Lake (East Sikkim District)



Arisaema griffithii Schott.

INTRODUCTION

(R.C. Srivastava)

The State of Sikkim situated on the flanks of the Eastern Himalayas between $27^{\circ}10'$ - $28^{\circ}05'$ N and $88^{\circ}30'$ - 89° E is bounded by Nepal in the West, Bhutan in the South-East, Tibet in the North and North-East and the Darjeeling district of West Bengal in the South.

The topography of this pretty Himalayan State is varied. The altitudes range from 244 m to over 8598 m. Most of the 7300 sq km area is interlaced with jungle-clad ridges and deep ravines created by, and through which, the raging torrents of the mountain rivers speed and emerald valleys alternating with their terraced hill-sides and dense forests ; a variety of plant beauties along with water-falls, rivulets, lakes and snow-capped mountains girdling the state in a protective embrace, embellish the land.

The climatic, edaphic, altitudinal and biotic variations with their complex inter-relationship and species composition have resulted in different kinds of vegetation cover in the area. The forests cover *ca* 42.8% (3124 sq. km) of the total geographic area, of which 2260 sq km are protected and reserve forests. The state is very rich in biological diversity. It is estimated that *ca* 5000 species of flowering plants, *ca* 350 pteridophytes and *ca* 15 species of Gymnosperms occur in this region. In addition to these a large number of non-flowering plants such as liver-worts, lichens, mosses, algae and fungi also occur.

CLIMATE : The abrupt variations in the altitude (elevations) have created diverse climatic conditions. The climate is warm and humid during summer and monsoon season (June - Oct.) and moderately cold during winter (Dec. - Feb.) at lower elevations. The winter months become more severe as one goes up. Places like Lachen, Lachung and Dzongri areas remain almost snow covered during most of the winter months. The perennial snow line however, begins from *ca* 4880 m only.

Most of the areas of the state are very wet during June to September, except the areas in North which are shielded from the monsoon clouds by the high embattlements of the Himalayas, or those areas where the monsoon loses its intensity. Average annual rainfall is *ca* 4000 mm and

average minimum/maximum temperature ranges between 4.2° - 17.1° C/13.9° - 23.9° C. Rainfall is usually very heavy during May - Sept. being maximum in July.

Relative humidity is 70 - 80% throughout the year and at most of the places in Sikkim.

PEOPLE : Sikkim's population as per the 1991 census is 4,03,612 with a density of 57 per sq. km. However, the density varies district wise due to varied climatic conditions and topography. East district has the highest of 157, while it is 133 for South District, 83 for West District and only 7 for North District. Gangtok being the Capital is the most populated place in the State of which *ca* 83% is rural and consists mainly of Nepalese, Bhutia, Lepcha, Limbus (very few). In the township, service - group and business people have come from the plains. But of these, Nepalese who themselves have a conglomeration of different ethnic types are at present largest in number. But the Lepchas (Rongkup-Children of Rong, Rongpa - ravine dwellers) are the earliest settlers of Sikkim. They have an extremely well stocked vocabulary on plants. (Rip-flower; Kung-tree). Even today the old and rural folk can distinguish most of the species at a glance and tell their local uses (mostly medicinal/edible). Sir J.D. Hooker the world renowned explorer of the Sikkim Himalaya also mentioned in glorious terms about the Lepcha's wonderful knowledge of the Sikkim plants.

The state has been divided into four districts viz. East, North, South and West.

BOTANICAL EXPLORATIONS

Sikkim's botanical diversity has attracted a large number of plant collectors from different parts of the world since Griffith's visit in 1843. The famous botanist J.D. Hooker visited Sikkim during 1848-49 and the account published by him in 'Himalayan Journals' is still one of the most comprehensive description of the botanical splendour of this region.

Subsequently distinguished botanists such as Sir G. King, C.B. Clarke, G.H. Cave, W.W. Smith and J.M. Cowan visited the area during later part of the 19th century and early 20th century. In the post 1940 period, comparatively little collections were made except for K.P. Biswas, R.S. Rao, B.D. Sharma, B. Ghosh Hara *et al.* and A.R.K. Sastry & P.K. Hajra. But the exploration work gained momentum after the establishment of the Sikkim Himalayan Circle of the Botanical Survey of India at Gangtok in December 1979. In the post 1979 period, P.K. Hajra, P. Chakraborty, B. Krishna, A.K. Verma, D.C.S. Raju,

R.C. Srivastava, S. Kumar and N.R. Mandal, P. Basu, B. Mittal, S. Singh, M. Sanjappa, M. Ahmedullah, V. Sampat Kumar, K.D. Kumar, A.K. Raut, L.B. Chowdhury collected plants from different areas of the state including the areas like Sakyong Valley (Lepchas land). The Herbarium of this Circle until now houses *ca* 14000 specimens.

VEGETATION

Broadly Sikkim can be divided into three distinct botanical zones viz. Tropical, Temperate and Alpine, depending on the elevation and characteristics of the vegetation. The foothills of Sikkim are covered with forests consisting of the species of *Shorea*, *Adina*, *Dalbergia*, *Dillenia*, *Artocarpus*, *Ficus*, *Bauhinia*, *Litsea*, *Lagerstroemia*, *Terminalia* etc. In the lower hill ranges (600-1500m) the forests chiefly include the species of *Schima*, *Syzgium*, *Duabanga*, *Engelhardtia*, *Ficus*, *Castanopsis*, *Pandanus*, *Cyathia*, *Michelia*, *Quercus*, *Saurauia*, *Photinia*, *Juglans*, *Leucoceptrum* etc.

The temperate vegetation comprises of forests consisting of the species of *Alnus*, *Acer*, *Betula*, *Magnolia*, *Rhododendron*, *Larix*, *Berbeirs*, *Salix*, *Cotoneaster*, *Vaccinium*, *Daphne*, *Sorbus*, *Rubus* and the herbaceous species of *Aconitum*, *Anemone*, *Potentilla* etc. *Abies densa*, *Larix griffithiana*, *Tsuga dumosa*, *Picea spinulosa* and *Taxus wallichiana* represent the coniferous belt in the altitudinal range of 2700-3900m. The Rhoddendron-Conifer forests comprising of the several species of *Rhododendron*, *Daphne*, *Betula* etc. mark the timberline in the altitude of *ca* 4000m.

The alpine vegetation which occurs above 4500m and up to a limit of *ca* 5500m is confined to moorlands of coarse meadows with many stunted and dwarf shrubby speices of *Rhododendron*, prostrate plants of *Juniperus squammata* , and cushion-like herbaceous species of *Arenaria*, *Androsace*, *Aconitum*, *Cassiope*, *Saxifraga*, *Primula*, *Pinguicula*, *Sedum*, *Rheum*, *Saussurea*, *Gentiana*, *Kobresia*, *Carex* etc.

The vegetation in the area may be classified into the following categories :

i) *Low Hill Forests* (tropical to sub-tropical type, up to 900m) : From the base (244m) to 900m, the sub-mountainous tracts are inhabited by dense broad-leaved semi-evergreen forests with trees attaining 24-36 m height. The rainfall is heavy, even up to 500cm annually. Canes yams and other climbers are common. Epiphytes like orchids, aroids etc. are abundantly represented. The undergrowth is luxuriant and varied. *Shorea robusta* (sal) is very common, covering large tracts especially

along the Teesta and Rangit rivers. *Schima wallichii*, *Bauhinia purpurea*, *Cedrela toona*, *Stereospermum tetragonum*, *Bombax ceiba*, *Dillenia pentagyna*, *Lagerstroemia parviflora*, *Sterculia villosa*, *Terminalia myriocarpa*, *T. tomentosa* and *Albizia* spp. are prominent components of the forests. Other trees occasionally met with *Garuga pinnata*, *Cedrela microcarp*, *Amoora wallichii*, *A. rohituka*, *Chukrasi tabularis*, *Evodia mellaefola*, *Ailanthus grandis*, *Duabanga grandiflora*, *Tetrameles nudiflora*, *Celtis tetrandra*, *Castanopsis indica*, *Syzygium formosa*, and *Michelia champaca* together with laurel like *Phoebe lanceolata*, *P. hainesiana*, *P. attenuata* *Litsea polyantha* and *Cinnamomum tamala*. Several species of *Artocarpus* such as *A. integrifolia*, *A. chaplasha* together with *Bischofia javanica* also occur. Large cultivated trees of *Ficus elastica* are often seen along banks of the river Teesta, along side Pakyong road, and near Dickchu. Occasionally interspersed in the forest are *Ficus semicordata* and *Pandanus nepalensis*. Tree-ferns are not uncommon in the low-land forests of Sikkim.

ii) *Middle Hill Forests* (subtropical type, 750-1500 m) : These are formed largely of evergreen species which are dominant. Deciduous trees may also occur. The trees are usually 20-30 m in height. Epiphytes and climbers occur in large numbers. The undergrowth is not usually dense and consists of numerous herbaceous and shrubby species. *Castanopsis tribuloides*, *C. indica*, *Schima wallichii* and *Phoebe hainesiana* are the commonest tree species between 750 and 1200m. *Michelia champaca* and *Stereospermum tetragonum* may also be occasionally met with. Other prominent components of the forests of this region are : *Drimycarpus racemosus*, *Juglans regia*, *Engelhardtia spicata*, *Spondias lutea*, *Exbucklandia populnea*, *Michelia cathcartii*, *Talauma hodgsonii*, *Saurauia nepalensis*, *Ficus oligodon*, *F. semicordata*, *Betula alnoides*, *Alnus nepalensis*, *Terminalia* spp., *Macaranga* sp., *Litsea polyantha*, *Phoebe lanceolata*, *P. attenuata*, and members of the family Meliaceae . Large evergeen trees of *Quercus glauca*, *Q. spicata*, *Q. serrata*, and *Q. griffithii* grow in dense formations between 1200 and 1600 m or above. Bamboos may also be found near habitations.

It will not be out of place to mention that *Cryptomeria japonica* is being extensively cultivated . It covers large areas and forms dense forests between 1200 and 2400 m altitude. It thrives best in this climate but due to rapid growth the wood remains soft and thus loses much of its commercial values. The ground vegetation underneath these forests is scanty due to lack of sun light and unsuitable substratum. Only a few ferns grow on the fringes of such forests.

iii) *Upper Hill forests* (Warm or wet temperate type 1500 -2700 m) : The forests are evergreen with medium-sized trees, rarely over

24 m height. There are a number of deciduous tree species but these form only a small proportion. Oaks and laurels form large patches in otherwise mixed forests. The oaks have branched crowns and are abundantly covered with mosses and other epiphytes. The forests extremely thick and the requisite amount of moisture available for the ground growing as well as epiphytic vegetation. The shelter of trees provided shade and prevents rapid air movements to a considerable extent. This results in a prolific growth of small herbs, shrubs and ferns on the forest floor. Woody climbers are frequent but not conspicuous.

Several altitudinal zones may be distinguished by the preponderance of certain such as laurels between 1800 and 2100 m, *Quercus lamellosa* between 2100 and 2400 m and *Q. pachyphylla* between 2400 and 2700m, though freely overlapping. *Michelia cathcartii*, *Mangolia campbelli*, *Machilus edulis*, *Quercus fenestrata* and *Castanopsis hystrix* are quite common in all the forests up to 2100m. Between 2100 and 2400 m of altitudes *Quercus lamellosa*, *Castanopsis tribuloides*, *Acer campbellii*, *Michelia excelsa* and *M. cathcartii* are dominant. *Quercus lineata* *Betula alnoides* and *Symplocos theaeifolia* are also frequently met with the former being quite prominent. The oaks constitute the greater part of the trees of top canopy and Lauraceae is usually relegated to the second storey, though numerically predominant. Laurels like *Machilus grammiana* *M. gamblei* *M. edulis* *Litsea sericea*, *elongata* *L. kingii*, *L. zeylanica* and *Cinnamomum obtusifolium* are not uncommon. *Alnus nepalensis* grows mainly along water courses and is the chief colonizer of new landslip areas. *Michelia excelsa* is well known timber tree of this zone.

Still higher up, between 2400m and 2700m or so *Quercus lamellosa* *Q. pachyphylla*, *Castanopsis tribuloides* *Acer campbellii* *Magnolia campbelli* *Symploco theaeifolia* and *Taxus wallichiana* are the prominent elements of the forests. Above 2700m in this zone *Quercus pachyphylla* occurs in pure formations. Under the shade of these forest *Rhododendron griffithianum* finds a favourable place. Dwarf bamboos, *Arundinaria* spp. are not uncommon as undergrowth at higher altitudes.

iv) *Rhododendron-Conifer Zone* (cold temperate or sub-alpine, 2700m-3600m) : The forests of this zone are also evergreen, mainly composed of Rhododendrons and conifers. Quite often *Quercus pachyphylla* and *Q. lineata* formations extend above 2,700m altitude and *Acer combellu*, *A. caudatum* *Betula utilis* an *Magnolia campbelli* may also be met with though very infrequently. As one proceeds higher up, there is a gradual replacement of oak trees by *Rhododendron arboreum* *R. campanulatum* and *R. grande* and other species of the genus. *Betula utilis* is occasionally found in the high level *Rhododendron*

forest at the head of Lachen valley near or above Samdong (3300 m). *Taxus wallichiana* grows in the forests as one proceeds above Lachung. Interspersed with these are patches of *Tsuga wallichiuma* and *Abies densa*. At about 2700-3000 m, in Lachen valley, *Tsuga dumosa* grows in abundance and is the dominant tree. It also grows at Chhoka in West Sikkim. *Picea spinulosa* grows abundantly on all the hills around Lachen intermixed with *Tsuga dumosa* but does not reach the heights above 3000 m. The bamboo, *Arundinaria aristata* forms dense undergrowth in silver fir forests, especially where fire has destroyed the tree canopy. *Abies densa* also occurs in almost pure formations between Karponang and Chhangu (East Sikkim), and Simdong to Thangu (North Sikkim) extending upto 3600 m or a little above. Few trees of *Salix wallichiana*, are also seen growing near Thangu along streams.

Rhododendron arboreum forms scrub on steeper slopes at about 3000 m. Above the tall tree line, the vegetation is a sort of mosaic of *Rhododendron campanulatum*, *R. wightii*, *R. thomsonii*, *R. cinnabarinum* and *R. decipens*. Lacaita scrub on slopes near Chhangu (3900 m) and near Thangu (3900 m). *Rhododendron anthopogon*, *R. setosum* and *R. barbatum* may also be occasionally met with. Grasslands are frequent at 2700 m altitude and above. *Arisaema* spp. may be found in open places. Various species of *Aconitum* grow abundantly on the forest floor underneath Rhododendrons at high altitudes especially around Thangu.

v) *Alpine Scrub and Grasslands* (3600-4300 m and above) : At the heights above 3600 m where the tree line ends, *Juniperous pseudosabina*, and *J. recurva* grow in bushy formations in North and East Sikkim especially on the exposed sunny hill slopes around Thangu (4200 m) and Chhangu (4300 m). *Ephedra gerardiana* covers vast areas tops of hills around Thangu (ca 4200). The species has not been seen flourishing anywhere else. The beautiful yellow-flowered *Rhododendron lepidotum* is another plant of high altitudes (3600 m or above) and grows in exposed rock crevices, hardly attaining 30 cm. In the open meadows on gentle mountain slopes a few species each of the genera like *Ranunculus*, *Anemone*, *Delphinium*, *Rhus*, *Potentilla*, *Primula*, *Fragaria*, *Cassiope*, *Allium* etc. are seen.

PHYTogeOGRAPHICAL ASPECTS

Owing to the varied climatic and ecological conditions in its entire length and breadth, the Sikkim Himalayan region offers very suitable spots into which a variety of floristic elements have migrated from several near and far-off lands. Sino-Japanese elements (viz. species of *Quercus*, *Schima* etc.) are quite common in this region. The floristic

elements of Western China, which are distributed all along the Sikkim Himalayas are *Aletris pauciflora*, *Anemone rupicola*, *A. vitifolia* and others. *Primula sikkimensis* and *Magnolia campbellii* are also the species of Western China which extend from Yunnan in to the Eastern Himalaya. The European and Mediterranean elements are represented by the species of *Ranunculus*, *Gentiana*, *Swertia*, *Anemone*, *Tamarix*, *Allium*, *Artemisia*, etc. The American elements in Sikkim flora are exhibited generally by weeds of agricultural lands, open forest edges and waste places e.g. *Eupatorium adenophorum*, *E. odoratum*, *Parthenium hysterophorus*, *Mikania cordata* and *Lantana camara*. African elements such as the species of *Flacourtiella*, *Grewia*, *Holarrhena*, *Alstonia* are represented in this region. The Siberian elements are exhibited in many predominantly temperate genera like *Potentilla*, *Pedicularis* and *Lonicera*. Tibetan elements like *Hippobae*, *Prezwalskia* etc. are also found. An example of considerable phytogeographical significance is the occurrence of *Glaux maritima* (Primulaceae) in Leh area of Kashimr and Llonhak valley of Sikkim. This plant is distributed along the costal and inland salt marshes of northen temperate and arctic regions. In addition to such examples, there are very interesting areas that can be called 'Isolation belts' that have led to the isolation of certian species. *Meconopsis bella*, *Cathcartia lyrata*, *Senecio chola*, *Sausurea laneana*, *Geranium* spp., *Primula elwesiana*, *Primula wattii* and *Swertia burkilliana* serve as such examples in the Sikkim Himalayas.

Studies made by Janaki Ammal in genera viz. *Magnolia*, *Camellia*, *Lonicera*, *Rhododendron* and *Viburnum* represented by many species in this region revealed high polyploidy which led her to consider this region (Eastern Himalayas) as a region of active speciation. The occurrence of many families and genera of primitive flowering plants such as *Magnolia*, *Manglietia*, *Euptelea*, *Tetracentron*, *Pycnarrhena*, *Haematocarups*, *Aspidocarya*, *Holboellia*, *Exbucklandia*, *Houttuynia*, *Myrica*, *Alnus* and *Betula* has led Takatajan to consider this region (Eastern Himalaya) the cradle of flowering plants.

PLANT RESOURCES

Germ Plasm

In Sikkim, the tribal populations have primitive but important germplasm of several crops. Still there is a tendency to preserve the local land races and varieties of crop plants which are the products of many years of natural selection and contain genetic treasures having natural resistance to pests and diseases and adaptability to stress conditions. Noteworthy among these is the prevailing enormous genetic diversity in Iskush, Maize, Wheat, Ginger, Cardamom, Mango, Banana, Orange,

Saccharum etc. The region is particularly important for crop plants such as brassicas, rice, cucumber, banana, mango, cardamom, *Dioscorea*, *Alocasia*, *Colocasia*, *Amorphophalus*, and horticultural significance like Orchids, Rhododendrons, Primulas, Pedicularis etc. Apart from these, several other elements in the flora which may not have an obvious economic value at present as the same has not been perceived as yet but in future may prove to be of much economic value in view of new vistas in plant research and changing patterns of our needs.

It is the sum total of such remarkable diversity which has made this state to be a 'Gene-bank' for a number of food crops, forest trees, medicinal plants, aromatic and highly ornamental plants etc.

Endangered plants

The flora of Sikkim is at present under great pressure due to biotic factors like various developmental projects viz. many Hydel power projects, road construction activities, heavy deforestation by burning and tree-felling for preparation of agricultural fields, intensive grazing, tourist-bungalows etc. These activities have destroyed many rich diversity centres on one side and on the other side have initiated several new land slides zones due to which a large number of precious rare plants are getting lost within a very short spell of time. These include : *Acer hookeri* var. *majus*, *Pimpinella tongloensis*, *P. wallichii*, *Pternopetalum radiatum*, *Lactuca cooperi*, *Arenerai thangoensis*, *Coelogyne treutleri*, *Cymbidium eburneum*, *C. hookerianum*, *C. whiteae*, *elegans*, *Cypripedium himalaicum*, *Didiciea cunninghamii*, *Diplomeris hirsuta*, *Paphipedium venustum*, *Zeuxine pulchra*, *Aconitum ferox*, *Cotoneaster simonsii*, *Picrorrhiza kurrooa*, *Acronema pseudotonera*, *Angelica bulbigena*, *Ceropegia hookeri*, *C. lucida*, *Codonopsis affinis*, *Rhopalocnemis phalloides*, *Carex kingiana*, *Lloydia himalensis*, *Aphyllorchis parviflora*, *Calanthe alpina*, *Ophiorrhiza lucida*, *Nardostachys grandiflora*, *Dennstaedtia elwesii*, *Mecodium levingei*, *Panax pseudoginseng*, *Calamus inermis*, *Livistonia jenkinsiana*, *Begonia rupicola*, *B. satrapis*, *B. sanctata*, *Lagerstroemia minuticarpa*, *Vanda spectabilis*, *Cyclogramma squanaestipes*, *Oreopteris elwesii*, *Christiopteris tripuspis*, *Rhynchospora sikkimensis*, etc.

Endemic plants

Endemics constitute another important element of biologically interesting plants. Being confined to narrow and restricted ecological niches, they are more prone to extinction due to adverse biotic or natural

factors. Sikkim as such has comparatively less number of endemics (which are confined to present political boundaries of Sikkim) because many of them range from Nepal to Bhutan. It is estimated that the total number of endemics truly confined to present Sikkim may be *ca* 2 percent of the total number in the flora - elements. These include *Agrostis neodebilis*, *Calamagrostis tripilifera* var. *tripilifera*, *C. tripilifera* var. *cumminsit*, *Catabrosa aquatica*, *Cyathopus sikkimensis*, *Drepanostachyum intermedium*, *Poa gammieana*, *Trisetum sikkimense*, *Carex kingiana*, *Rhynchospora sikkimensis*, *Coelogyne treutleri*, *Anaphalis cavei*, *A. hookeri*, *A. subumbellata*, *Artemisia thellungiana*, *Blumea sikkimensis*, *Cremanthodium decaisnei* f. *clarkei*, *C. palmatum* ssp. *benthamii*, *Crepis atropappa*, *Gentiana glabriuscula*, *G. prainii*, *G. pluviarum*, *G. recurvata*, *Inula macrosperma*, *Jaeschkea microsperma*, *J. smithii*, *Ligularia dur*, *L. hookeri* ssp. *clarkei*, *L. pachycarpa*, *Saussurea forrestii*, *S. lineana*, *S. nimborum*, *S. obscura*, *S. pantlingiana*, *Swertia ramosa*, *S. rex*, etc. besides many more.

To conserve the depleting resources of this region, several attempts are being made. Kanchanjanga National Park, Fambongla Wildlife sanctuary and Kabi sacred groove are good examples where several endangered taxa are conserved. There are some botanical gardens viz. Jawaharlal Botanic Garden, Saramsta Botanic Garden and Orchid sanctuary, Rhododendron sanctuary near Yumthang, etc., wherein exist and insist conservation of many species is being attempted. Different sacred grooves situated in various parts of the state are playing a very positive role. But at present the main emphasis is to relocate or recollect the threatened taxa and attempts should be made for their *in-situ* conservation. However, in some cases the help of advance technologies viz. Tissue culture technique etc. can be used with advantage. In addition to this, the areas of diversity centre like Pangolekha range, Tendong ridge, Sakyong valley, Dongbong valley etc. should be brought under full protection.

Medicinal plants

Sikkim Himalayan region is the abode of a large variety of medicinal plants. *Prezwalskia tangutica*, *Nardostachys jatamansi*, *Picrorrhiza kurrooa*, *Aconitum luridum*, *Podophyllum hexandrum*, *Dactylorrhiza hatagirea*, *Taxus wallichiana*, *Ephedra gerardiana* and *Lycopodium clavatum* constitute the most important medicinal plants of the alpine zone. Several medicinal plants like *Dichroa febrifuga*, *Houttuynia cordata*, *Artemisia vulgaris*, *Rubia cordifolia*, *Panax pseudoginseng*, *Dioscorea deltoidea*, *Digitalis purpurea*, *Bergenia ciliata* are quite common in temperate and sub-temperate zones.

Tropical zone is also quite rich in the medicinal flora wherein plants viz. *Costus speciosus*, *Vitex negundo*, *Solanum viarum*, *Cissampelos pariera*, *Woodfordia fruticosa*, *Oroxylum indicum*, *Alstonia scholaris*, *Abroma augusta* and to some extent *Rauvolfia serpentina*, *Terminalia chebula*, *Hollarhena pubescens* etc. grow in good number.

Species of Horticultural importance

As the Flora of Sikkim includes plants of tropical, temperate and alpine characters in great diversity, a large number of plants found wild in this state (or the Eastern Himalayas as such), are of great horticultural importance. Many of them have been introduced into the European gardens. These include the species like : *Hedychium gardnerianum*, *Luculia gratissima*, *Allium wallichii*, *Acer oblongum*, *Anemone vitifolia*, *Arisaema griffithii*, *Berberis aristata*, *B. asiatica*, *B. sikkimensis*, *Bergenia ciliata*, *Betula utilis*, *Boenninghausenia albiflora*, *Cautleya gracilis*, *C. spicata*, *Cymbidium cyperifolium*, *Daphne bholua*, *Juniperus recurva*, *Mahonia acanthifolia*, *Meconopsis bella*, *Meillia thrysiflora*, *Paris polyphylla*, *Pleione praecox*, *Polygonatum oppositifolium*, *Prunus cerasoides*, *Rhododendron anthopogon*, *R. arboreum*, *R. barbatum*, *R. hodgsonii*, *R. nivale*, *Rosa macrophylla*, *R. sericea*, *Vandopsis undulata*, *Viburnum cordifolium*, *Primula* spp. and many others. But still several species of *Corydalis*, *Gentiana* and *Primula* being of interest and beauty, await introduction.

Food plants

There are over 350 species of flowering plants and Pteridophytes found wild in diverse localities of Sikkim which may serve as emergency life-saving food-products. These include : flowers of *Urtica dioica*, *U. parviflora* (Sisnoo-Nep.) ; rhizomes of *Dioscorea* species (Tarul-Nep.) ; fruits of *Aesandra butyracea* (Chiwari-Nep, Yelkung-Lep.) ; *Calamus erectus* (Betfal-Nep., Rue-Lep.); *Elaeagnus conferta* (Malindo-Nep.); *Heracleum lanatum* (Chimpin-Nep.) ; *Podophyllum hexandrum* (Papri-Nep.); *Machilus edulis* (Lepcha-Kawla, Phunchey-Nep., Phom-Kung-Lep.) ; *Melia eubia* (Lapsi-Nep., Silot-Kung-Lep.) ; *Morus australis* (Sanukimbu-Nep., Mekrake Kung-Lep.), *Terminalia chebula* (Hara-Nep., selim pot-Lep.), *Zanthoxylum acanthopodium* (Bogay Timur-Nep., Nangryupot-Lep.) ; leaves of *Phytolacca acinosa* (Jaringo-Nep.) ; young shoots of *Rheum nobile* (Tohuka-Nep.) *Diplazium esculentum*, sheath from young shoots of *Polygonum molle* and *Ficus virens*; tubers of *Satyrium ciliatum* (Khiru-Tibetan) ; flower buds of *Bauhinia purpurea* ; inflorescence of *Tupistra natans* (Nakima-Nep.). Leaves of *Camellia*

kissii and bark of *Betula* spp. are used for making tea, white millet. *Rhododendron* (Flowers), Junipers, corn, East Himalayan cherry etc. are brewed as local wine which are named after the source plant viz. Guras wine, Cherry brandy, Juniper wine etc.

Fire-wood yielding species

Due to the total dependence of the rural folk on fire wood for cooking and warming of the houses, any woody plant is cut and used as a fire wood ; yet some important firewood yielding species include : *Alnus nepalensis* (Ujis-Nep.), *Betula cylindrostachys* (Saur-Nep.) *Castanopsis indica*, *Engelhardtia spicata* (Mauwa-Nep.), *Toona ciliata* (Tooni-Nep.), *Cryptomeria japonica* (Dhupi-Nep.), *Acer campbellii* (Kapasey-Nep.), *Schima wallichii* (Chilaune-Nep.), etc. At higher elevations even the plants like *Abies densa*, *Tsuga dumosa*, *Picea smithiana*, *Rhododendron* spp. *Larix griffithiana* are also used.

Timber yielding species

Most important timber yielding species of the State include *Shorea robusta*, *Tectona grandis*, *Juglans regia*, *Castanopsis indica*, *Quercus lamellosa*, *Toona ciliata*, *Cryptomeria japonica* and *Terminalia myriocarpa*.

Dye yielding plants

Most important plants used on commercial scale include : roots of *Rumex nepalensis* (Halhale-Nep.), stem of *Ruia manjith* (Majitho-Nep.) ; bark of *Mahonia nepalensis* (Kerapsing-Bhutia) ; fruits of *Dichroa febrifuga* (Basak-Nep., Gebokhamok-Lep.); leaves of *Symplocos glomerata* (Sungi-Nep.) ; bark of *Juglans regia* (Okhar-Nep.) etc. In addition to these several other dye yielding plants viz. *Byxa orellena*, *Mallotus philippensis*, several spp. of *Indigofera* etc. occur wild in the state.

Fodder yielding species

Although a large number of species are employed for feeding the livestock, yet the preferred ones include *Ficus auriculata*, *F. oligodon*, *F. hirta*, *F. hispida*, *F. virens*, *F. clavata*, *F. nemoralis*, *Sauraia nepalensis*, *Morus alba*, *Brasssiopsis mitis*, *Artocarpus lakoocha*, *Grewia oppositifolia*, *Litsea polyantha* etc.

Botanical curiosities

Several examples of biologically very interesting plants in Sikkim Flora include the insectivorous plants like *Drosera burmanii*, *Pingicula alpina*, *Utricularia* spp. (*Lentibulariaceae*), beautiful parasites like *Aeginitia indica* (*Orobanchaceae*) etc. Several plants, which grow especially in the high alpine meadows, survive extreme adverse ecological conditions by special adaptations; e.g. woolly species of *Saussurea* (*S. gassypiphora*) or cushion like or bushy habit (species of *Arenaria*, *Festuca*, *Juniperus*, *Gaultheria*, *Saxifraga* etc.). Some of the very ancient living angiosperms viz. *Magnolia pterocarpa* (*Magnoliaceae*), *Tetracentron sinense* var. *himalayana* (*Tetracentraceae*), *Talauma hodgsonii* (*Magnoliaceae*) occur in this state. Several medicinal plants like *Podophyllum hexandrum*, *Panax pseudoginseng*, *Dactylorrhiza hatagirea*, *Picrorrhiza kurrooa*, *Nardostachys grandiflora*, *Berberis aristata*, *B. asiatica*, *Valeriana wallichii*, *Ephedra gerardiana* var. *sikkimensis*, *Taxus wallichiana*, *Mandragora caulescens*, *Aconitum ferox*, *A. spicatum*, *Dioscorea deltoidea*, *Rauvolfia serpentina* etc. which created sensational news are natural components of Sikkim Flora.

A large number of epiphytes such as foliose lichen; a variety of mosses and ferns cover available space on the bark of the roots suspended in the air with the ability to absorb moisture from the mist and the clouds. Some species of *Rhododendron*, *Impatiens* (1 sp.) *Aeschranthes*, *Polygonatum*, *Cautleya*, *Hedychium*, *Gonathanthus* and many orchids and aroids are epiphytic in these forests.

Sacred groves

Sacred groves are playing important role in 'in-situ' conservation of the indigenous elements of the Flora and Fauna of Sikkim. Most of these are the place of worship and have Monastery at the summit. A few like Kabi-sacred grove being the place of union of Lepchas and Bhutias, are of historical importance. Many of them are declared as 'reaseerve forests' and are well protected from the biotic pressure. Khetchipuri Sacred-grove in West district, Mt. Mainam, Rabong, Tendong Sacred groves in South district, Churten Sacred grove in East district and Kabi Sacred-grove in North district are a few good examples where a variety of precious elements of Flora and Fauna are well protected. *Cyathea spinulosa* and *C. gigantea* - the tree-ferns which are under threat due to demand of their trunks in orchid-culture and trade, are well protected in Churten Sacred-grove. *Camellia kissi* - the wild-tea is common in the protected woods around Khetchipheri lake. 'Red-Panda' (*Ailurus fulgens*) - the State-Animal of Sikkim-finds the dense

thickets of Kabi Sacred-grove quite suitable for its inhabitation. *Dendrobium nobile* the State-Flower of Sikkim is also quite common here.

Conservation aspects

Vavilov (1951) was very correct in saying that “many of the areas of centre of origin of crop plants are the areas where the tribal people live, and sooner or later the Governments of these regions, are bound to uplift them socially and economically. Once these communities are in different socio-economic milieu, the working of thousands of years of natural selection will be lost because the people would like to go in for high out-put agriculture.” This statement is very true in case of Sikkim.

During the recent past a great deal of damage has been done to the richness of the biodiversity of Sikkim Himalayas. The forests have been denuded by the haphazard felling of trees for timber, through unmanaged grazing, road-construction activities etc. Rural folk of Sikkim still depend totally on firewood for cooking and also for warming their houses. Frequent landslides resulting from deforestation are swallowing several habitats of many biologically and economically important plants even before they or their utility is known to the society.

Not only deforestation, but even afforestation programmes are also responsible to a greater extent for the loss of biodiversity of Sikkim Flora. Plantation of exotic trees like *Cryptomeria japonica*, *Pinus patula* etc. may have added to the glamorous green-cover but the fact is that because of its impact on soil, several precious elements of the ground flora are getting lost. Actually the indigenous species like *Alnus nepalensis*, *Duabanga grandiflora*, *Terminalia myriocarpa* etc. should be included in afforestation programme. Now a days great emphasis is being paid on promotion of Tourism in the State which is showing a disruptive influence on local traditional and social mores by introducing new and not necessary desirable life-styles, and has destructive effect on biodiversity as well.

Because of large scale cultivation of large-cardamom which is a major revenue yielding crop of the State, several precious elements of ground Flora (e.g. *Paphiopedilum venustum*-the lady's slipper orchid) are getting lost.

Another major threat to the floristic diversity is the mushroom growth of small and large scale Hydel-power projects. These are causing an irreparable damage to Flora of the State.

Being the border line state defence set-ups in Sikkim cannot be avoided. But it has definitely resulted in a large scale deforestation and soil erosion because the impact of army activities on the fragile ecosystem and geological aspects have not been taken into consideration.

Although several botanical expeditions have been made to Sikkim since 1843, yet even today any serious botanical exploration yields a number of novelties such as new taxa, new records, and many interesting plants, some of which are of great economic and botanical importance. Therefore, our present knowledge about the plant resources of Sikkim is not the last word on the real situation in this region. Still much is to be done and much is to be known about the hidden 'Green-Gold' treasure of the State.

The strategy of utilization of plant resources of the Sikkim Himalayas is to be drawn after taking into consideration all the factors and to ensure that the large variety of flora of this region not only continues to survive but maintains conditions for the evolution of the flora i.e. for speciation or origin of new taxa.

SELECTED REFERENCES

- AHLOOWALIA, B.S. AND N.L. DHAWAN (1972) A synopsis in maize from Sikkim. *Indian J. Genet.* 32(2) : 299 - 233.
- AIRYSHAW, H.K. (1932) *Leycesteria gracilis* (Kurz) Airy - Shaw *Hook. Icon. Pl.* 32. 3166.
- ANDERSON, T. (1964) *Cultivation of quiniferous cinchona in British Sikkim*. Edinburg.
- ANDERSON, T. (1868) An enumeration of the palms of Sikkim. *J. Linn. Soc.* 11 : 4 - 14.
- ANDERSON, T. (1871) An enumeration of the palms of Sikkim - Tibet Frontier. *Kew Bull.* 297 - 315.
- ANONYMUS (1853) Rhododendrons of Sikkim Himalaya. *Hook. J. Bot.* 152 - 154.
- ANONYMOUS (1893) Botanical explorations in Sikkim - Tibet Frontier. *Kew Bull.* 297 - 315.
- ANONYMOUS (1963) *Spring Flora of Sikkim Himalaya* by Japanese members of the Indo - Japanese botanical expedition to Sikkim and Darjeeling in 1960. Tokyo.
- BANERJI, A. (1948) A track in Sikkim. *Ind. For.* 75 : 228 - 230.
- BASU, P. (1990) Cobra Lilies (*Arisaema Mart.*) in Sikkim. *J. Hill. Res.* 3 : 135 - 145.
- BASU, P. (1990) Pipeworts (Eriocaulaceae) of Sikkim. *J. Econ. Tax. Bot.*
- BASU, P. & B. MITRA (1991) A note on the less known plants *Bischofia javanica* (Bischofiaceae) in Sikkim. *J. Econ. Tax. Bot.* 15 (3) : 703 - 704.
- BASU, P. & B. MITRA (1992) Less known Noteworthy plants of Sikkim Himalaya. *J. Econ. Tax. Bot.* 16 (1) : 209 - 212.
- BHANJA, K.C. (1948) *Mystic Tibet and the Himalayan Darjeeling*. Oxford Book and Stationary (One Chapter on Sikkim).
- BISWAS, K. (1956) Observation on the vegetation of Eastern Sikkim. *Yearb. R. Asiat. Soc. Beng.* 18 : 146.
- BISWAS, K. (1956) *Common medicinal plants of Darjeeling and Sikkim Himalayas*. Calcutta.

BISWAS, K. (1966) *Plants of Darjeeling and Sikkim Himalayas*, Pt. I. Calcutta.

BISWAS, K. AND H.P.V. TOWNEDN (1936) Common plants of Northern Sikkim. *Proc. 23rd Indian Sci. Congr. Abst. Soc. (Bot.) IV* : 7 - 8.

BRANDIS, P. (1906) The spruce of Sikkim and Chumbi Valley, *Ind. For.* 32 : 579 - 581.

BROWN, P. (1934) *Tours to Sikkim*. Revised 2nd ed. by J. Towned. Calcutta.

BRUHL, P. (1926) *A guide to the orchids of Sikkim*, (being a guide to the identification of those species of orchids found between the Terai and the Northern frontier of independent Sikkim including the Chumbi valley and British Bhutan). Calcutta.

BUCHENAU, F.R. (1867) Zwei neue *Juncus* Arten Aus dem Sikkim Himalayas, Gesammelt von, J.D Hooker & Thomson, Bot. Ztg. : 145 - 158.

BURKILL, I.H. (1907) Alpine notes from Sikkim. *Kew Bull.* 92 - 94.

BURKILL, I.H. (1908) Notes on the pollination of flowers in India. Note No. 5, some autumn observations in the Sikkim Himalaya. *J. Asiat. Soc. Beng. (n.s.)* 4 (4) : 179 - 195.

BURKILL, I.H. (1916) A note on the Terai forests between the Gandak and Teesta. *J. Asiat. Soc. Beng. (n.s.)* 12 : 267 - 272.

CAMPBELL, A. (1849) A journal of a trip to Sikkim in Dec. 1848. *J. Asiat. Soc. Beng.* 18 : 482 - 541.

CAMPBELL, A. (1852) Diary of a journey through Sikkim to the Frontiers of Thibet. *J. Asiat. Soc. Beng.* 21 : 407 - 428.

CAVE, G.H. AND W.W. SMITH (1911) The vegetation of the Zemu and Lonakh Valleys of Sikkim. *Rec. Bot. Surv. India* 4(1) : 141 - 260.

CAVE, G.H. AND W.W. SMITH (1911) A note on the Himalayan species of *Alangium*. *Rec. Bot. Surv. India* 6 (4) : 93 - 98.

CHING, R.C. (1931) Genus *Vittaria* of China and Sikkim - Himalaya (The studies of Chinese ferns -VI), *Sinensis* 1 : 175 - 192.

CHING, R.C. (1938) A revision of the Chinese and Sikkim Himalayan *Dryopteris* with reference to some species from

neighbouring regions. *Bull. Fan. Inst. Biol. Peking* 6 : 237 - 352. 196 ; 8 (6) : 363 - 507.

CHOPRA, G.L. (1931) The Distribution of Lichen in Darjeeling and Sikkim. *Proc. 18th Indian Sci. Congr. Abs. (Bot.)* v : 264.

CLARKE, C.B. (1877) Botanic notes from Darjeeling to Tonglo. *J. Linn. Soc. Bot.* 15 : 116 - 159.

CLARKE, C.B. (1886) Botanic notes from Drajeeling to Tonglo and Sundakphoo. *J. Linn. Soc. Bot.* 21 : 384 - 291.

DEB, B.D. (1957) Endemism and outside influence on the flora of Manipur. *Proc. 4th Indian Sci. Congr. Abst. (Sect. Bot.)* VI : 283 - 284. (Phytogeographical aspect of Sikkim - Himalayan plants).

DUNCAN, P.C. (1935) Sikkim Rhododendrons. *Himalayan J.* 7 : 111 - 112.

FISCHER, C.E.C. AND FRODERSTROM (1912) New plants from Tibet and Sikkim. *Kew Bull.* 135 - 140.

GAMBLE, J.S. (1917) The Himalayan species of *Skimmia*. *Kew Bull.* 301. - 303.

GAMMIE, G.A. (1893) Note on Sikkim tree ferns. *J. Linn. Soc. (Bot.)* 29 : 482 - 485.

GAMMIE, G.A. (1893) Botanical exploration in Sikkim - Tibet frontier, *Kew Bull.* 297 - 315.

GAMMIE, G.A. (1894) Report of a botanical tour in Sikkim. *Rec. Bot. Surv. India* 1 (2) : 1 - 24. 1894 ; *J. Bombay Nat. Hist. Soc.* 9(2) : 197 - 216 (Oringinal paper was published in *Rec. B.S.I.*, reprinted in *J. Bombay Nat. Hist. Soc.*).

GAMMIE, G.A. (1894) Vegetation of temperate and Alpine Sikkim. *Gazeteer of Sikkim* 95 -111.

GAMMIE, J.G. (1981) Vegetation in Risby's *Gazetter of Sikkim* : 80 - 94.

GHOSH, B.N. (1951 - 57) A catalogue of plants of Sikkim Himalaya. *J. BengNat. Hist. Soc.* 25 : 22 - 36. 1951 ; 26 : 127 - 141. 1953 ; 186 - 195. 1954 ; 27 : 74 - 82. 1954 ; 104 - 122. 1955 ; 160 - 186. 1955 ; 28 : 67 - 77. 1955 ; 135 - 137. 1955 ; 29 : 43 - 51. 1957 ; 97 - 106. 1957.

GHOSH, B.N. (1953) Orchids of Sikkim. *J. Beng. Nat. Hist. Soc.* 26 : 33 - 37.

GUPTA, B.K. (1994) Higher plants of Indian subcontinent, Vol. III. 161 - 166 ; 201 - 210.

HAJRA, P.K. & P. CHAKRABORTY (1981) A survey of wild plants sold in the Lal Market of Gangtok. *Ind. J. For.* 4(3) : 217 - 220.

HAJRA, P.K. AND B.K. DAS (1982). Vegetation of Gangtok with special reference to exotic plants. *Indian Forester* 554 - 566.

HARA, H. (1965) *Spring Flora of Sikkim Himalaya*, Osaka.

HERZOG, TH. (1939) Zwei Bryophytensammlungen aus dem Sikkim Himalaya. *Ann. Bryol., Hague* 12 : 71.

HODGSON, B.H. (1948) Route from Kathmandu, capital of Nepal to Darjeeling and Sikkim, interspersed with remarks on the people and country. *J. Asiat. Soc. Beng.* 17(2) : 634 - 646.

HOOKER, J.D. (1849 - 1851) *The Rhododendrons of Sikkim Himalaya*, 2nd ed. London.

HOOKER, J.D. (1849) Notes, chiefly botanical, made during an excursion from Darjeeling to Tonglo, a lofty mountain on the confines of Sikkim and Nepal. *J. Asiat. Soc. Beng.* 18 : 419 - 446.

HOOKER, J.D. (1849 - 1850) Botanical mission to India, Calcutta to Darjeeling in Sikkim - Himalaya. *Hook. J. Bot.* 1 : 1 - 14, 41 - 56, 81 - 89, 113 - 120, 129 - 136, 161 - 175, 226 - 233, 274 - 282, 301 - 320, 331 - 336, 361 - 370. 1894 ; 2 : 11 - 23, 52 - 59, 88 - 91, 112 - 118, 115 - 151, 161 - 173, 213 - 218, 224 - 249. 1850.

HOOKER, J.D. (1851) On the physical geography of Sikkim Himalaya. *Hook. J. Bot.* 3 : 21 - 31.

HOOKER, J.D. (1852 - 1854) On the climate and vegetation of the temperate and cold regions of East Nepal and the Sikkim - Himalayan mountain. *J. Roy. Hort. Soc.* 7(1) : 69 - 131, 1852 ; *J. Agri. Hort. Soc. India* 8 : 35 - 65, 73 - 95. 1854 (Original paper was published in *J. Roy. Hort. Soc.* reprinted in *J. Ari. Hort. Soc. India*).

HOOKER, J.D. (1854) *Himalayan journals*, or notes of a naturalist in Bengal, the Sikkim and Nepal Himalayas, the Khasia mountains etc. Vol. 1 -2, London.

HOOKER, J.D. (1875 - 97) Flora of British India vols. I - VII London.

HUMBOLDT, B.A. (1851) Physical geography of Sikkim Himalaya. *Hook. J. Bot.* 3 : 21 - 31.

KANAI, H. (1963) Phytogeographical observations on the Japano - Himalayan elements. *J. Fac. Sci. Tokyo Univ. III. (Bot.)* 8(8) : 305 - 339.

KING, G. AND R. PANTLING (1985) On some new orchids from Sikkim. *J. Asiat. Soc. Beng.* 64. Pt. 2 (3) : 329 - 344.

KING, G. & R. PANTLING (1895) A second series of new orchids from Sikkim. *J. Asiat. Soc. Beng.* 65. Pt. 2 (2) : 118 - 134.

KING, G. & R. PANTING (1989) The corycids of the Sikkim. Himalaya. *Ann. Roy. Bot. Gard. (en)* and *put. Calc.* 8(1) : 1 - 342 ; 8 (2 - 4) : 1 - 448.

KRISHNA, B. & B.K. DAS (1983) Fibre yielding plants of Sikkim. *J. Econ. Tax. Bot.* 4 : 129 - 132.

KRISHNA, B. & B.K. DAS (1984) Some new plant records for Sikkim Himalaya. *Geobios New Report* 3 (2) : 145 - 146.

KRISHNA, B. & B.K. DAS (1984) Additions to the Pteridophytic Flora of Sikkim Himalaya. *Bull. Bot. Surv. India* 26 (3 - 4) : 20 - 221.

KRISHNA, B., B. MITRA & D.C.S. RAJU (1988) Yams (*Dioscorea* L.) in Sikkim. *J. Hill. Res.* 1 : 67 - 72.

KRISHNA, B. (1990) Fuelwood species of Sikkim Himalayas. *Proc. Nat. Symp. Fuelwood* 1896 ; 100 - 102 (Publ. 1990).

KRISHNA, B. & S. SINGH (1985) Ethnobotany of Sikkim *J. Econ. Tax. Bot. Vol. & Page lacking*

KUMAR, S. & D.C.S. RAJU (1991) Large cardamom and its wild relatives in Sikkim Himalayas. *J. Hill. Res.* 2 : 102 - 107.

KUMAR, S. & D.C.S. RAJU (1991) Ginger and its allies in Sikkim Himalaya in B.K. GUPTA, *Higher plants of Indian sub - continent* Vol. II : 249 - 225.

KUMAR, S. (1991) Turmeric (*Curcuma longa* L.) & related taxa in Sikkim Himalaya. *J. Econ. Tax. Bot.* 15 (3) : 721 - 724.

KUMAR, S. & P. SINGH (1991) A synoptic account of Grasses Flora of Sikkim - Panicoideae. *Ind. J. For.* (In press).

KUMAR, S. & P. SINGH, (1991) Nomenclatural notes on taxa of Bambusoideae of Sikkim. *J. Ind. Bot. Soc.* 70 : 423 - 424.

LACAITA, C.C. (1915 - 1917) Plants collected in Sikkim including the Kalimpong District, April 8th to May 9th. *J. Linn. Soc. Bot.* 43 : 457 - 492.

MAJUMDAR, N.C. (1970) Notes on *Steilaria saxatilis* Buch. - Ham ex D.Don, *Stellaria vestita* Kurz and *Sellaria sikkimensis* Hook. f. *J. Bombay Nat. Hist. Soc.* 67 (i) : 26 - 29.

MATTHEW, K.M. (1970) A bibliography of the botany of Sikkim. *Bull. Bot. Soc. Beng.* 24 (1 - 2) : 57 - 59.

MEHRA, P.N. (1932) *Ceropteris calomelanos* L. in Sikkim. *J. Ind. Bot. Soc.* 11 (4) : 340 - 341.

MEHRA, P.N. & S.S. BIR (1964) Pteridophytic flora of Darjelling and Sikkim Himalayas. *Res. Bull. (n.s.) Panj. Univ.* 15 (1 - 2) : 69 - 181.

MITRA, J.N. (1985) *Flowering plants of Eastern India*. Calcutta.

MITRA, B. (1989) Screw-Pine (*Pandanus nepalensis* st John) in Sikkim Himalaya *J. Hill Research* 2 : 66 - 67.

MUELLER, J. (1895) *Licheness sikkimensis & Reveenois montibus Stevens* in Sikkim, *Indiae Orientalis. Bull. Herb. Boissier.* 3 : 194 - 195.

OSMASTON, G. C. (1935) An expedition into Sikkim. *Indian For.* 61 : 424 - 434, 487 - 499.

PRADHAN, B.B. (1940) *Kikiyu grass*. Gangtok.

PRADHAN, K.C. (1962) *Flora of Lachen*. London.

PRADHAN, K.C. (1963) Distribution and ecological habitat of Sikkim Himalayan orchids. *Orchidconf.* 61 - 66.

PRADHAN, U.C., K.C. PRADHAN & S.T. LACHUNGPA (1991) *Rhododendrons of Sikkim Himalaya*, Kalimpong.

RAJU, D.C.S. & R.C. SRIVASTAVA (1990) *Satyrium ciliatum* Lindl. - a less - known edible orchid of Sikkim Himalaya. *Proc. Nat. Acad. Sci.*

RAJU, D.C.S. & R.C. SRIVASTAVA (1993) Pokeweed family (Phytolaccaeae) in Sikkim. *J. Himal. Bot.*

RAJU, D.C.S. & R.C. SRIVASTAVA (1990) Lalchandan family (Daphniphyllaceae) in Sikkim. *J. Himal. Bot.* (13 & 14) : 12 - 14.

RAJU, D.C.S., B. KRISHNA & S.T. LACHUNGPA (1987) Deorali Orchid Sanctuary, Gangtok. *Bull. Bot Surv. India* 29 : 253 - 271.

RAJU, D.C.S., B. KRISHNA & S.T. LACHUNGPA (1990) Sharmsha Orchid Sanctruary, Ranipool, Sikkim, *Ibid.* 32 : 121 - 128.

- RAO, R.S. (1963) A botanical tour in the Sikkim state. Eastern Himalaya. *Bull. Bot. Surv. India* 5(2) : 165 - 205.
- RISLEY, H.H. (1894) *Gazetteer of Sikkim*, Calcutta 77 - 112 p.p.
- ROY CHOWDHURY, K.C. (1951) Sikkim, the country and its forests. *Indi. For.* 77 (ii) : 676 - 683.
- SCOTT, J. (1874) On the tree ferns of British Sikkim, with descriptions of three new species and a few supplemental remarks on their relations to Palms and Cycads. *Trans. Linn. Soc. London* 30 : 1 - 44.
- SEN, G.C. (1963) The epiphytic plants of Darjeeling Hills, other than orchids. *Bull. Bot. Surv. India* 5 (2) : 111 - 115.
- SEN, S. (1957) A visit to Changu. *J.Beng. Nat Nist. Soc.* 29 (1 - 2) : 1 - 5.
- SHARMA, B.D. AND. B. GHOSH (1970) Contribution to the flora of Sikkim - Himalayas. *Bull. Bot. Soc. Beng.* 24 (1 - 2) : 45 - 55.
- SHARMA, B.D, N.P. BALAKRISHNAN, R.R. Rao & P. K. Hajra (1993). Flora of India, Vol. I., B.S.I. Calcutta.
- SHARMA B.D, N.P. BALAKRISHNAN & M. SANJAPPA (1993) Flora of India, Vol II, B.S.I. Calcutta.
- SHARMA, B.D, M. SANJAPPA & N. P. BALAKRISHNAN (1993) Flora of India, Vol. III, B.S.I., Culcutta.
- SHEBBEAR, E.O. (1934) The conifers of the Sikkim Himalaya and adjoining contry. *Ind. For.* 60 (10) : 710 - 713.
- SHERHILL. J.L. (1892) Journal of trip undertaken to explore the glaciers of the Kanchanjungha group in the Sikkim Himalaya in November 1891. *J. Asiat. Soc. Beng.* 31 (5) : 457 - 479.
- SMITH, H. (1970) New or little known Himalayan species of *Swertia* and *Veratrilla* (Gentianaceae), *Bull. Brit. Mus. (Nat. Hist.) Bot.* 4 (6) : 239 - 258.
- SMITH, H.W. (1913) The Alpine and Sub - alpine vegetation of South - Eastern Sikkim. *Rec. Bot. Surv. India* 4(7) : 323 -431.
- SMITH, W.W. AND G.H. CAVE (1911). The vegetation of Zemu and Llonakh valleys of Sikkim. *Rec. Bot. Surv. India* 4 (1) : 141 - 260.
- SMITH., H.W. AND G.H. CAVE (1913). A note on the Himalayan species of *Daphne*. *Rec. Bot. Surv. India* 6(2) : 45 - 54.

SRIVASTAVA, R.C. (1990) A note on the arrow poison used by Lepchas of Dzongu in Sikkim and its antidote. *Nat. Acad. Sci. Letters* 13 (ii) : 399 - 400.

SRIVASTAVA, R.C. (1991) *Gnephallium affine* D.Don-a less known useful herbs of Sikkim, *Ibid* 14 : 9.

SRIVASTAVA, R.C. (1992) Maple family (Aceraceae) in Sikkim. *Proc. Nat. Acad. Sci.* 62 (B) : 385 - 392.

SRIVASTAVA, R.C. (1993) Gymnosperms of Sikkim, *Asia J. Life. Sci.* 2 (1) :

SRIVASTAVA, R.C. (1993) High - altitude medicinal plants of Sikkim. *Jour. Res. Edu. Ind. Med.* 1993 (Jan. - March) 5 - 14.

SRIVASTAVA, R.C. (1994) Diversity of Flora of Eastern Himalaya and its conservation, in Dwivedi & Pandey (ed.) *Biotechnology in India*, India.

SRIVASTAVA, R.C. AND A.K. MAINERA (1994) *Dactylorhiza hatagirea* (D.Don) Soo - a less known medicinal plant of Sikkim. *Nat. Acad. Sci. Letters*.

THAPA, J.K. (1966) Primitive maize with the Lepchas. *Bull. Tibetol.* 3 (1) : 29 - 32.

WHITE, J.C. (1909) *Sikkim and Bhutan ; Twenty one years on the North - East, Frontier, 1887 - 1908.*

PLAN OF ENUMERATION

The families are arranged in the sequence of Bentham & Hooker's *Genera Plantarum* (1862 - 63) which was adopted in J.D. Hooker's *Flora of British India* (1872 - 97), but their delimitations have sometimes been amended in accordance with later concepts. For example *Scitamineae* of *Flora of British India* (FBI) is here divided into *Zingiberaceae*, *Cannaceae*, *Marantaceae* and *Musaceae*. Genera and species are listed alphabetically. A good number of exotic taxa have been included, especially those which have become or are becoming naturalised or have become prominent feature of the general vegetation. For each species, the accepted correct name with its place and date of publication, basionym, if any, and a few synonyms relevant to nomenclature or to earlier important regional publications are given. The local names are given in capital letters followed by the name of the languages in brackets. The local language are abbreviated as Lep. (Lepcha), Nep. (Nepali), and Bhut. (Bhutia), Tib. (Tibetan). Other abbreviations used in literature citations are listed separately.

Only brief distinguishing characters are mentioned. The information about the distribution within Sikkim is provided. The localities of recorded occurrence are followed by the altitudes of occurrence. The period of flowering/fruiting is also provided in most of the cases. Additional information regarding economic importance or any other peculiarity is sometimes given under notes.

BURMANNIACEAE

(R.C. Srivastava)

BURMANNIA L.

Burmannia coelestis D. Don, Prodr. Fl. Nep. 44. 1825; Hook. f., F.B.I. 5: 665. 1888 ; Hajra in Fascl. Fl. India 19 : 9. 1988.

Herbs, up to 20 cm tall. Leaves linear-lanceolate, acute or acuminate. Flowers blue or purplish with yellow lobes.

Sikkim, s. l. ; up to 1500 m.

Fl. & Fr. : June - Nov. In marshy habitats.

B. disticha L., Sp. Pl. 1 : 287. 1753; Hook. f., F.B.I. 5 : 644. 1888 ; Hajra, l. c. 11.

Herbs, up to 50 cm tall. Leaves linear or lanceolate, 2 - 12 x 0.5 - 1.5 cm, acute or acuminate at apices. Flowers blue or purple with yellow tipped greenish lobes.

Sikkim, s.l. ; 1000 - 1500 m.

Fl. & Fr. : June - Sept. In paddy fields.

B. nepalensis (Miers) Hook.f., F.B.I. 5 : 666. 1888 ; Hajra, l.c. 12. *Gonyanthes nepalensis* Miers in Trans. Linn. Soc. London 18 : 537. t. 38. f. 1. 1841.

Herbs, up to 15 cm tall. Leaves reduced to scales, appressed to stem, elliptic. Flowers white with yellow tinge.

Sikkim, s.l. ; 1000 - 1500 m.

Fl. : July - Sept. In moist habitats.

ORCHIDACEAE

(R.C. Srivastava)

ACAMPE Lindl.

Acampe ochracea (Lindl.) Hochr. in Bull. New York Bot. Gard. 6 : 270. 1910. *Saccolabium ochraceum* Lindl. in Edw., Bot. Reg. Misc. 2. 1842 ; Hook. f., F.B.I. 6 : 62. 1890.

Epiphytes. Leaves oblong. Flowers yellow, ca 10 mm across, borne on leaf-opposed 2 - 6-flowered panicles.

In valleys.

Fl. : Nov. - Dec.

A. papillosa (Lindl.) Lindl., Fol. Orchid. 2. 1853. *Saccolabium papillosum* Lindl. in Bot. Reg. t. 1552. 1841 ; Hook. f., F.B.I. 6 : 63. 1890.

Epiphytes. Leaves coriaceous, curved, conduplicate, apically truncate or obliquely notched. Flowers yellow with brown base, ca 8 mm across.

Found at lower elevations.

Fl. : Oct. - Jan.

Notes : Roots tonic, said to have cooling effect, useful in rheumatism.

A. rigida (Buch. - Ham. ex J.E. Sm.) Hunt in Kew Bull. 24 : 98. 1970. *Aerides rigida* Buch. - Ham. ex J.E. Sm. in Rees, Cyclop. 39, Aerides n. 12. 1819. *Saccolabium longifolium* (Lindl.) Hook. f., F.B.I. 6 : 62. 1890 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 220. t. 292. 1898. *Acampe longifolia* (Lindl.) Lindl., Fol. Orch. Acampe 1. 1853. *Vanda longifolia* Lindl., Gen. & Sp. Orchid. 215. 1833.

Epiphytes. Leaves oblong, coriaceous. Inflorescence leaf opposed, much shorter than the leaves.

Found in tropical valleys ; up to 1500 m.

Fl. : June.

ACANTHEPHIPPIUM Bl.

Acanthephippium striatum Lindl. in Bot. Reg. 241. 45. 1838 ; Hook. f., F.B.I. 5 : 816. 1890.

Terrestrial herbs with ovoid pseudobulbs. Leaves subcoriaceous, plicate, elliptic. Flowers dull white, suffused and blotched with yellow, the apices of sepals and petals purplish.

Sikkim, s.l. 600 - 1200 m.

Fl. & *Fr.* : April - May.

A. sylhetense Lindl., Gen. & Sp. Orchid. 117. 1833 ; Hook. f., F.B.I. 5 : 815. 1890.

Terrestrial herbs with elongate - obpyriform pseudobulbs. Leaves elliptic. Flowers pale pink with bold red lines ; lip yellow tinged.

Sikkim, s. l. ; 540 - 900 m.

Fl. : July - Aug.

ACRIOPSIS Reinw. ex Bl.

Acriopsis javanica Reinw. ex Bl., Bijdr. 377. 1825 ; Hook f., F.B.I. 6 : 79. 1890. *Acriopsis harai* Tuyama in J. Jap. Bot. 39 : 129. 1964 ; Pradhan, Ind. Orch. 2 : 717. 1979.

Sympodial epiphytes. Rhizomes creeping. Pseudobulbs crowded. Flowers 2 - 40 mm apart, 8 - 14 mm across, lip 3 - lobed.

In the vicinity of Gangtok ; ca 1000 m.

Fl. : April.

ACROCHAENE Lindl.

Acrochaene punctata Lindl., Fol. Orchid. 1. 1853 ; Hook. f., F. B.I. 5 : 783. 1890.

Epiphytic herbs with ovoid pseudobulbs. Leaf solitary, oblong, petiolate. Flowers pale yellowish green, dotted and streaked with purplish brown.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : Oct. - Nov.

AERIDES Lour.

Aerides multiflora Roxb., Pl. Corom. 3 : 68. t. 271. 1820 ; Hook. f., F.B.I. 6 : 44. 1890.

Epiphytic herbs with stout stem. Leaves narrowly oblong. Flowers white, marked with pink or purple.

In tropical valleys ; up to 1100 m.

Fl. : April.

A. odorata Lour., Fl. Cochinch., 2 : 528. 1790 ; Hook. f., F. B.I. 6 : 47. 1890.

Epiphytic herbs. Stems stout. Leaves flat, keeled oblong, unequally lobed at apex. Flowers white, spotted with pink.

Sikkim s. l. ; ca 1200 m.

Fl. : May - June.

AGROSTOPHYLLUM Bl.

Agrostophyllum brevipes King & Pantl. in Ann. Roy. Bot. Gard. Calcutta. 8 : 156. t. 213. 1898.

Epiphytic herbs. Rhizomes narrow, covered with spreading subulate scales. Leaves linear-oblong, membranous. Flowers white, column and anther red.

Sikkim, s.l. ; ca 600 m.

Fl. : July.

A. callosum Reichb. f. in Sem. Fl. Vit. 296. 1868 ; Hook. f., F.B.I. 5 : 824. 1890.

Epiphytic herbs. Rhizomes stout, branched, clothed with triangular scales. Leaves sessile, linear-oblong. Flowers reddish, borne in heads.

Sikkim, s.l. ; ca 1800 m.

Fl. : June - July.

A. khasianum Griff. in Calc. J. Nat. Hist. 4 : 378. t. 19. 1844 ; Hook. f., F.B.I. 5 : 824. 1890.

Epiphytic herbs with tufted double fusiform compressed, sheathed stem. Leaves narrowly oblong, subcoriaceous. Flowers white with pale yellow patches.

In tropical valleys.

Fl. : Aug.

A. myrianthum King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 155. t. 211. 1898.

Epiphytic herbs. Rhizomes stout, clothed with adpressed lanceolate scales. Leaves linear-oblong, sub-coriaceous. Flowers yellow.

Teesta valley ; ca 3600 m.

Fl. : Aug.

AMITOSTIGMA Schltr.

Amitostigma puberula (King & Pantl.) Schltr. in Fedde Repert. 4 : 92. 1919. *Orchis puberula* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 304. 1898.

Terrestrial herbs, less than 15 cm tall. Leaf solitary, linear. Flowers white.

Lachung valley ; ca 3000 m.

Fl. : July.

ANOECTOCHILUS Bl.

Anoectochilus crispus Lindl. in J. Linn. Soc. 1 : 180. 1857. *Odontochilus crispus* (Lindl.) Hook. f., F.B.I. 6 : 99. 1890.

Terrestrial herbs, up to 20 cm tall. Leaves ovate, acute. Sepals green ; petals and lip white with yellow tinges on sac.

Mahalderam Peak (?) ; ca 1800 m.

Fl. : Sept.

A. elwesii (Clarke ex Hook. f.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 296. t. 394. 1898. *Odontochilus elwesii* Clarke ex Hook. f., F.B.I. 6 : 100. 1890.

Terrestrial herbs, up to 25 cm tall. Leaves ovate. Sepals green with white mid-ribs and brown tips. Petals white with brown tips. Lip white, teeth of claw brown, basal sac green.

Chungthang, Tendong ; 900 - 1800 m.

Fl. & Fr. July-Aug.

A. grandiflorus Lindl. in J. Linn. Soc. 1 : 179. 1857. *Odontochilus grandiflorus* (Lindl.) Benth. in Gen. Pl. 2 : 598. 1883 ; Hook. f., F.B.I. 6 : 100. 1890.

Terrestrial herbs, up to 45 cm tall. Leaves elliptic. Flowers green, mottled with white and tipped with pink.

Tendong ; 1200 - 1800 m.

Fl. : July.

A. lanceolatus Lindl., Gen. & Sp. Orchid. 499. 1840. *Odontochilus lanceolatus* (Lindl.) Bl., Orchid. Arch. Ind. 81 1858 ; Hook. f., F.B. I. 6 : 101. 1890.

Terrestrial herbs, up to 30 cm tall. Leaves ovate-lanceolate. Flowers pale green ; dorsal sepal brown-tinged ; lip bright yellow : anthers pink.

Yoksum ; 1500 - 2100 m.

Fl. : July - Sept.

A. roxburghii (Wall.) Lindl. in Royle, Illustr. Bot. Himal. 368. 1839 & Gen. & Sp. Orchid. 499. 1840 ; Hook. f., F.B.I. 6 : 95. 1890. *Chrysobaphus roxburghii* Wall., Tent. Fl. Nep. 37, t. 27. 1826.

Plants including inflorescence less than 30 cm tall. Leaves ovate or ovate-oblong, velvety, purplish red with golden reticulations. Flowers white.

In tropical valleys.

Fl. : Nov.

A. sikkimensis King & Pantl. in J. Asiat. Soc. Beng. 65 : 124. 1896.

Plants less than 23 cm tall. Leaves elliptic ovate, velvety, dark red, with golden yellow veins on upper surface, dull red beneath. Flowers olive green and white.

Sikkim, s.l.; 900 - 1500 m.

Fl. : Sept.

ANTHOGONIUM Wall. ex Lindl.

Anthogonium gracile Wall. ex Lindl., Gen. & Sp. Orchid. 426. 1840 ; Hook. f., F.B.I. 5 : 822. 1890.

Terrestrial herbs with small ovoid pseudobulbs. Leaves 2-5, linear, plicate, sessile, sheathing. Flowers white or rose coloured ; lip white with dark purple spots ; anthers bright yellow.

On way to Lachen ; 1200 - 2100 m.

Fl. : Sept.

APHYLLORCHIS Bl.

Aphyllorchis alpina King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 26. t. 347. 1898.

Terrestrial herbs up to 75 cm tall. Stems sheathed. Flowers yellowish red.

Singale - La, Mon - Lepcha (?) ; ca 4200 m.

Fl. : July - Aug.

A. montana (Thw.) Reichb. f. in Linnaea 41 : 57. 1876 ; Hook. f., F.B.I. 6 : 116. 1890. *Apaturia montana* Thw., Enum. Zeyl. 301. 1861. *Aphyllorchis prainii* Hook f., F.B.I. 6 : 117. 1890.

Terrestrial herbs, 60 cm tall. Whole plant pale brown except yellow pollinia.

Sikkim, s.l. ; ca 600 m.

Fl. Aug.

A. pantlingii W.W. Smith in Rec. B.S.I. 4 : 243. 1911. *A. parviflora* King & Pantl. in J. Asiat. Soc. Beng. 65 (2) : 128. 1896 *et* in Ann. Roy. Bot. Gard. Calc. 8 : 262. t. 348. 1898.

Terrestrial herbs, ca 35 cm tall. Almost all parts pale brown except white anthers and yellow pollinia.

Lachung ; ca 3000 m.

Fl. : July.

APOSTASIA Bl.

Apostasia wallichii R. Br. in Wall., Pl. Asiat. Rar. 1 : 75. t. 84. (*A. odorata*) 1830 ; Hook. f. F.B.I., 6 : 175. 1890.

Terrestrial herbs, 30-60 cm tall. Leaves linear, lanceolate. Flowers yellow.

Sikkim, s. l.; (cf. Hara *et al.* 1978).

APPENDICULA B1.

Appendicula cornuta Bl., Bijdr. 7 : 302. 1825. *A. bifaria* Lindl. in Hook. f., F.B.I. 6 : 84. 1890.

Herbs with tufted, slender, leafy, often compressed stem. Leaves many, lanceolate. Flowers white, flushed with yellow.

In tropical valleys.

Fl. : Aug.

ARACHNIS B1.

Arachnis cathcartii (Lindl.) J.J. Smith in Naturw. Tijdrachr. Nederl. Ind. 72 : 75. 1912. *Vanda cathcartii* Lindl., Fol. Orchid. Vanda 8 : 1853. *Arachnanthe cathcartii* (Lindl.) Benth. in Benth. & Hook. f., Gen. Pl. 3 : 573. 1883 ; Hook. f., F.B.I. 6 : 27. 1890.

Epiphytic herbs. Stems stout, pendulous. Leaves sessile, oblong, unequally and obliquely bifid at apex. Flowers brown-yellow on upper surface.

Sikkim, s.l. ; 900 - 1800 m.

Fl. : March - April.

ARUNDINA Bl.

Arundina graminifolia (D. Don) Hochr. in Bull. New York Bot. Gard. 6 : 270. 1910. *Bletia graminifolia* D. Don., Prodr. Fl. Nep. 29. 1829. *Arundina chinensis* Bl., Bijdr. 8 : 502. 1825 ; Hook. f., F.B.I. 5: 857. 1890. *A. bambusifolia* Lindl., Gen. & Sp. Orchid. 125. 1830 ; Hook. f., l.c.

Terrestrial herbs. Stems reed-like, up to 2 m tall. Leaves narrowly oblong-lanceolate. Flowers pinkishpurple.

Gangtok, Jorethang-Namchi, Teesta valley. Rangeet valley ; ca 1750 m.

Fl. & Fr. : April - Sept.

ASCOCENTRUM Schltr.

Ascocentrum ampullaceum (Roxb.) Schltr. in Fedde Repert. 1 : 975. 1913. *Aerides ampullaceum* Roxb., Fl. Ind. 3 : 476. 1832.

Saccolabium ampullaceum (Roxb.) Lindl., Sert. Orchid. t. 17. 1838 ; Hook. f., F.B.I. 6 : 64. 1890.

Epiphytic herbs without pseudobulbs. Leaves narrowly oblong, keeled, coriaceous, mottled with brown. Flowers pink, except deep purple pollinia.

Sikkim, s.l. ; 300 - 1000 m.

Fl. : March - April.

BIERMANNIA King & Pantl.

Biermannia bimaculata (King & Pantl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 200. t. 267. 1898. *Sarcochilus bimaculatus* King & Pantl. in J. Asiat. Soc. Beng. 44 : 340. 1875.

Epiphytes. Leaves linear - oblong. Flowers white, with 2 brown blotches on the calli or lip.

Teesta valley ; ca 450 m.

Fl. : July.

BRACHYCORYTHIS Lindl.

Brachycorythis obcordata (Buch. - Ham. ex D. Don) Summerh. in Kew Bull. 243. 1955. *Orchis obcordata* Buch. - Ham. ex D. Don, Prodr. Fl. Nep. 23 : 1825, non Willem. 1796. *Habendria galeandra* auct. non Benth : Hook. f., F.B.I. 6 : 163. 1890.

Terrestrial herbs, with globose or oblong tubers. Leaves oblong. Flowers pale purple.

Sikkim, s.l. ; 1000 - 2000 m.

BULBOPHYLLUM Du. Petit Thouars.

Bulbophyllum affine Lindl., Gen. & Sp. Orchid. 48. 1830 ; Hook. f., F.B.I. 5 : 756. 1890 ; King & Pantl., Ann. Roy. Bot. Gard. Calc. 8 : 67. t. 91. 1898.

Epiphytes. Pseudobulbs cylindric, ca 7.5 mm apart. Leaves narrowly oblong, obtuse at apex. Flowers pale green with purple nerves.

Teesta valley, Zemu, Tadong - Rumtek ; 600 - 1000 m.

Fl. : June - Sept.

B. andersonii (Hook. f.) J.J. Smith in Bull. Jard. Bot. Buitenz. 2, 8 : 22. 1912. *Cirrhopetalum andersonii* Hook. f., F.B.I. 5 : 777. 1890.

Epiphytes. Pseudobulbs elliptic-ovoid, fibrous at base. Leaf oblong, subacute. Flowers pink with minute dark spots.

Teesta valley ; ca 450 m.

Fl. : Oct.

B. bisetum Lindl. in Ann. Nat. Hist. 10 : 186. 1842 ; Hook. f., F.B.I. 5 : 767. 1890.

Epiphytes. Pseudobulbs ovoid - conic, with a fibrous sheath at base. Leaves narrowly oblong - lanceolate, acuminate and obliquely bifid at apex. Flowers purplish.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. - Sept.

Notes : Flowers are remarkable for the spurs which run parallel to the dorsal sepal and the petals.

B. bootanense (Griff.) Par. & Reichb.f. in Trans. Linn. Soc. 30 : 153. t. 32. f. 2. 1874. *Cirrhopetalum bootanense* Griff., Notul. 3 : 297. 1851. & Ic. Pl. Asiat. t. 299. 1851 ; Hook. f., F.B.I. 5 : 775. 1890.

Epiphytes. Pseudobulbs narrowly ovoid. Leaves fleshy, narrowly elliptic, obtuse or subacute. Flowers red with dark red spots.

Sikkim. s.l. ; tropical valleys.

Fl. : April.

B. careyanum (Hook.) Spr., Syst. Veg. 3 : 732. 1826 ; Hook f., F.B.I. 5 : 760. 1890. *Anisopetalon careyanum* Hook., Fl. Exot. 2 : t. 149. 1825.

Epiphytes. Pseudobulbs ovoid-subsessile. Flowers brownish.

Sikkim, s.l. ; 450 - 1200 m.

Fl. Oct. - Dec.

B. caudatum Lindl., Gen. & Sp. Orchid. 56. 1830 ; Hook. f., F.B.I. 5 : 759. 1890.

Epiphytes. Pseudobulbs ovoid, compressed, furrowed. Leaves ovate-lanceolate, acute, coriaceous. Flowers white.

Gangtok ; ca 1800 m.

Fl. : April - July.

B. caulinorum Hook. f. in Hook., Ic. Pl. t. 2036. 1890 & F.B.I. 5 : 758. 1890.

Epiphytes. Pseudobulbs cylindric-elliptic. Leaves oblong, obtuse. Flowers green, flushed with brown.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : July.

B. clarkeanum King & Pantl. in J. Asiat. Soc. Beng. 64 : 333. 1895.

Epiphytes. Pseudobulbs cylindric or ovoid. Leaves narrowly elliptic-oblong, notched at apices. Flowers white with orange-red tips.

Sikkim, s.l. ; 450 m.

Fl. : June.

B. cornu-cervi King & Pantl. in J. Asiat. Soc. Beng. 64 : 332. 1895.

Epiphytes without rhizomes. Pseudobulbs minute, 2 - 3 mm, crowded, globular. Leaves ovate - elliptic, sessile or subsessile. Flowers green with reddish brown margins ; lip yellow inside, reddish brown outside.

Sikkim, s.l. ; ca 759 m.

Fl. : July.

B. cylindraceum Lindl., Gen. & Sp. Orchid. 53 : 1830 ; King & Pantl., l.c. 70 : t. 96. 1898.

Epiphytes without pseudobulbs. Leaves oblong-elliptic, acute. Flowers blackish purple.

Sikkim, s.l. ; 1600 - 2400 m.

Fl. : Oct. - Jan.

B. ebulbum King & Pantl. in J. Asiat. Soc. Beng. 64 : 334. 1895.

Epiphytes without pseudobulbs. Leaves solitary, oblong-lanceolate. Flowers pale green.

Sikkim, s.l.

B. eublepharum Reichb. f. in Walp. Ann. 6 : 252. 1861 ; Hook. f., F.B.I. 5 : 746. 1890.

Epiphytes. Rhizomes lacking. Pseudobulbs cylindric, closely placed. Leaf oblong. Flowers green.

Sikkim, s.l. ; 1800 - 2100 m.

Fl. : Aug.

B. elatum (Hook. f.). J.J. Sm. in Bull. Jard. Bot. Buitenz. 3, 3 : 23. 1912. *Cirrhopetalum elatum* Hook. f., F.B.I. 5 : 775. 1890.

Epiphytic caespitose herbs. Pseudobulbs cylindric. Leaves narrowly oblong, obtuse. Flowers dull yellow spotted.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : May.

B. gamblei (Hook. f.) Hook. f. in Hook. Ic. Pl. 3. t. 2039. 1890 & F.B.I. 6 : 188. 1890. *B. leptanthum* Hook. f. var. *gamblii* Hook. f., l.c. 5 : 759.

Epiphytes. Pseudobulbs narrowly obpyriform or subcylindric. Leaves oblong-lanceolate, apically notched.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : July - Aug.

B. griffithii (Lindl.) Reichb. f. in Walp. Ann. 6 : 247. 1881 ; Hook. f., F.B.I. 756. 1890. *Sacropodium griffithii* Lindl., Fol. Orchid 6. 1855.

Epiphytes. Pseudobulbs caespitose, narrowly ovoid. Leaves sessile, narrowly oblong. Inflorescence 1 - flowered. Flowers yellowish with red - brown spots.

Lachung ; ca 2700 m.

Fl. : Aug.

B. guttulatum Wall. ex Hook. f., F.B.I. 5 : 776. 1890. *Cirrhopetalum guttulatum* Hook. f., l.c. *B. umbellatum* in Edw., Bot. Reg. 30. t. 44. 1844, non Lindl. 1830.

Epiphytes. Pseudobulbs caespitose, crowded, ovoid. Leaves elliptic-oblong, subacute. Flowers pale yellow, purple spotted.

Sikkim s.l. ; 600 - 1500 m.

Fl. : July - Sept.

B. gymnopodus Hook. f. in Hook. f., Ic. Pl. 3, t. 2040. 1890 & F.B.I. 5 : 764. 1890.

Epiphytes. Pseudobulbs ovoid, sheathed at base. Leaves narrowly oblong, notched at apices. Flowers white.

Sikkim, s.l.

Fl. & Fr. : Nov. - Dec.

B. helenae (O. Ktze.) J.J. Sm. in Bull. Bot. Gard. Buitenz. 2, 8 : 24. 1912. *Phyllum helenae* O Ktze., Rev. Gen. Pl. 2 : 676 1891. *Cirrhopetalum cornutum* Lindl. in Edw., Bot. Reg. 24 : Misc. 75. 1838 ; Hook. f., F.B.I. 5 : 774. 1890.

Epiphytes. Pseudobulbs narrowly ovoid, compressed. Leaf narrowly oblong, obliquely notched. Flowers reddish brown.

Sikkim, s.l.

Fl. : May - June.

B. hirtum (J. E. Sm.) Lindl., Gen. & Sp. Orchid. 51 : 1830 ; Hook. f., F.B.I. 5 : 672. 1890. *Stelis hirta* J.E. Sm. in Res. Cyclop. 34. 1816.

Epiphytes. Pseudobulbs closely placed, ovoid - oblong, terete with a scarious terminal collar, oblong - oblanceolate, flaccid, absent during flowering time. Flowers yellow, green outside, white inside.

Sikkim, s. l. ; 900 - 1800 m.

Fl. : Oct. - Jan.

B. hymenanthum Hook. f. in Hook., Ic. Pl. 3 : t. 2046. 1890 ; Hook. f., F.B.I. 5 : 767. 1890.

Epiphytic herbs with thread like rhizomes. Pseudobulbs lacking. Leaves sessile, ovate - elliptic. Flowers yellowish, streaked with purple.

Lachung valley, Yoksum ; 1700 - 2400 m.

Fl. : May.

B. leopardianum (Wall.) Lindl., Gen. & Sp. Orchid. 48. 1830 ; Hook f., F.B.I. 7 : 756. 1890. *Dendrobium leopardianum* Wall., Tent. Fl. Nepal 39 : t. 1826.

Epiphytes. Pseudobulbs obpyriform, crowded. Leaves elliptic-oblong. Flowers whitish or pale ochraceous, spotted with crimson.

Sikkim, s.l. ; 1700 - 2200 m.

Fl. : Oct. - Nov.

B. leptanthum Hook. f., F.B.I. 5 : 759. 1890.

Epiphytes. Pseudobulbs cylindric. Leaves narrowly oblong. Flowers yellowish green.

Sikkim, s.l. ; 900 m.

Fl. : July.

B. listeri King & Pantl. in J. Asiat. Soc. Beng. 64 : 34. 1895.

Epiphytes without rhizomes. Pseudobulbs narrowly ovoid, compressed. Leaves linear-oblong, tapering at both ends.

Ratong Chu ; 900 - 2100 m.

Fl. : May - Sept.

B. odoratissimum (J.E. Sm.) Lindl., Gen. & Sp. Orchid. 55. 1830 ; Hook. f., F.B.I. 5 : 758. 1890. *Stelis odoratissima* J.E. Sm. in Rees, Cyclop. 34 : Stelis no. 12. 1816.

Epiphytes. Pseudobulbs subcylindric. Leaves oblong-lanceolate, slightly notched at apex. Flowers white with yellow tinge.

Ratong Chu ; ca 100 m.

Fl. : May.

B. ornatissimum (Reichb. f.) J.J. Sm. in Bull. Jard. Bot. Buitenz. 2, 8 : 26. 1912. *Cirrhopetalum ornatissimum* Reichb. f. in Gard. Chron. 2 : 424. 1882 ; Hook. f., F.B.I. 5 : 773. 1890.

Epiphytes. Pseudobulbs narrowly ellipsoid. Leaves oblong-obtuse and slightly notched at apices. Flowers yellowish green with purple markings.

Sikkim, s.l.

Fl. : Oct. Nov.

B. pantlingii Lucksom in J. Bombay Nat. Hist. Soc. 90 (3) : 551. 1993. *B. flavidum (flavida)* Lucksom *Ibid* 90 (1) : 71. 1993, non Lindl. 1840.

Epilithic herbs. Pseudobulbs ovoid, broadly ridged. Leaves 2, oblong-ob lanceolate, flaccid. Inflorescence solitary, from the base of pseudobulbs. Flowers orange yellow.

Phyangla R.F., Lachung valley R.F. ; 1000 - 2135 m.

Fl. & Fr. : July - Dec.

B. parvulum Hook. f., F.B.I. 5 : 778. 1890.

Epiphytes. Pseudobulbs ovoid. Leaves narrowly elliptic, subacute, sessile. Flowers drooping.

Sikkim, s.l. ; 2100 - 2700 m.

Fl. : Aug.

B. piluliferum King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 76. t. 104. 1898.

Epiphytes. Pseudobulbs caespitose, globose or depressed globose. Leaf fleshy, linear-oblong, obtuse. Flowers pale reddish brown with streaks of darker shade.

Sikkim, s.l. ; ca 300 m.

Fl. : May.

B. polyrhizum Lindl., Gen. & Sp. Orchid. 53. 1830 ; Hook. f., F.B.I. 5 : 767. 1890.

Epiphytes. Pseudobulbs globular or ovoid. Leaves narrowly oblong, sessile. Flowers yellowish.

Teesta valley ; ca 450 m.

Fl. : April.

B. protractum Hook. f., F.B.I. 5 : 758. 1890.

Epiphytes. Pseudobulbs cylindric. Leaves oblong, acute. Flowers pale yellow ; sepals orange in upper half.

Sikkim, s.l. ; 450 - 600 m.

Fl. : July.

B. reptans (Lindl.) Lindl., Gen. & Sp. Orchid. 51. 1830 ; Hook. f., F.B.I. 5 : 768. 1890. *Tribrachia reptans* Lindl., Col. Bot. t. 41. a. 1825.

var. *reptans*

Epiphytes with thin, wiry branching rhizomes. Pseudobulbs obpyriform. Leaves linear-oblong. Flowers yellowish with purple spots.

Sikkim, s.l. ; 400 - 2400 m.

Fl. : Oct. - Nov.

var. **sub-racemosa** Hook.f., F.B.I. 6: 769. 1890.

Sepals buff coloured, flushed with brown and spickled slightly with red ; petals coloured; lip yellow at tip.

Sikkim, s.l.

Fl.: Oct. - Dec.

B. retusiusculum Reichb. *f.* in Gard. Chron. 1182. 1869.
Cirrhopetalum wallichii Lindl., Gen. & Sp. Orchid. 59. 1830; Hook. f.,
F.B.I. 5: 776. 1890.

Epiphytes. Pseudobulbs ovoid. Leaves narrowly oblong, subacute, obliquely notched. Flowers deep violet, with darker spots.

Sikkim, s.l.; 1800 - 2250 m.

Fl.: July - Aug.

B. rigidum King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 69.
t. 94. 1898.

Epiphytes. Pseudobulbs lacking. Leaves rising direct from rhizome, each with a whorl of fibres at base. Flowers brownish yellow flushed with green, nerves pusplish.

Sikkim, s.l. ; 1800 m.

Fl. : May - June.

B. rolfei (O.Ktze.) Seid. in Dansk. Bot. Ark. 33: 149. 1979.
Phyllum rolfei O. Ktze., Rev. Gen. Pl. 677. 1891. *Cirrhopetalum parvulum* Hook. f., F.B.I. 5: 778. 1890.

Eptphytes. Pseudobulbs ovoid. Leaf narrowly elliptic, sub-acute. Flowers drooping.

Sikkim, s.l. ; 2100- 2700 m.

Fl.: Aug.

B. sarcophyllum (King & Pantl.) J.J. Sm. in Bull. Jard. Bot. Buitenz. 2, 8: 27. 1912. *Cirrhopetalum sarcophyllum* King & Pantl. in J. Asiat. Soc. Beng. 64 : 335. 1895.

Epiphytes. Pseudobulbs discoid. Leaves pendulous, fleshy, oblong, subfalcate, acute at apices. Flowers reddish brown, much mottled ; lip purple with dark purple blotches.

Sikkim, s.l. ; 900 - 1500 m.

Fl. : June - Sept.

B. scabratum Reichb. *f.* in Walp. Ann. 6: 259. 1861 ; Seid. in Dansk. Bot. Akk. 29: 67. t. 24. 1973. *Cirrhopetalum caespitosum* Wall.

ex Lindl. in Edw. Bot. Reg. 24: Misc. 35. 1830 ; Hook. f., F.B.I. 5 : 775 1890. *B. confertum* Hook.f., F.B.I. l.c. 757.

Epiphytic, caespitose herbs. Pseudobulbs much crowded, ovoid. Leaves linear-lanceolate, acute. Flowers ochraceous yellow.

Sikkim, s.l. ; 1200 m.

Fl. : April.

B. secundum Hook. f., F.B.I. 5: 764. 1890.

Epiphytes. Pseudobulbs crowded, turbinate, *ca* 7 mm in diam. Leaves narrowly oblong. Flowers pale purple with dark nerves.

Chungthang, Tendong ; 1800 - 2100 m.

Fl. : June - July.

B. sikkimense (King & Pantl.) J.J. Sm. in Bull. Jard. Bot. Buitenz. 2, 8: 12. 1912. *Cirrhopetalum sikkimensis* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8: 90. t. 125. 1898.

Epiphytes. Pseuobulbs ovoid-globose. Leaves oblanceolate, obtuse, obliquely notched, fleshy. Dorsal sepal greenish with few brownish red markings, lateral sepals ochraceous, minutely spotted with reddish brown. Petals ochraceous with three broad reddish brown nerves and purplish marginal cilia.

Sikkim, s.l.

Fl. : May.

B. striatum (Griff.) Reichb. f. in Walp. Ann. 6: 257. 1861. ; Hook. f., F.B.I. 5: 775. 1890. *Dendrobium striatum* Griff., Notul. 3: 318. 1851.

Epiphytes. Pseudobulbs obpyriform. Leaf elliptic, retuse at apex, coriaceous. Flowers yellowish green with purple strips.

Sikkim, s.l. ; 2100 m.

Fl. : Oct.

B. thomsonii Hook.f., F.B.I. 5 : 764. 1890.

Epiphytes. Pseudobulbs subdiscoid. Leaf narrowly oblong, acute. Flowers yellowish white.

Sikkim, s.l. ; 1200 m.

Fl. : Sept.

B. triste Reichb. f. in Walp. Ann. 6: 253. 1861; Hook.f., F.B.I. 5: 768. 1890. *B. micranthum* Hook. f., l.c.; S. Misra in Bull. Bot. Surv. India 22: 152. (1980) 1982. *B. alopecurum* Reichb. f. Gard. Chron. 2: 70. 1880; Hook.f., l.c. 770.

Epiphytes. Pseudobulbs turbinate with umbonate apices, closely placed or 1.2 - 2.5 cm apart; leafless during flowering. Leaves 2, from each pseudobulb, narrowly oblong, acute at apex. Flowers purplish, except for a yellow line on the edges of the lip and ovary.

Teesta valley; ca 450 m.

Fl. : March.

B. viridiflorum (Hook.f.) Schltr. in Orchids 4: 108. 1910.
Cirrhopetalum viridiflorum Hook. f., F.B.I. 5 : 779. 1890.

Epiphytic, caespitose herbs. Pseudobulbs ovoid. Leaf in pairs on young pseudobulbs, narrowly oblong or oblong - lanceolate, Flowers yellowish green turning to pale brown.

Sikkim, s.l ; 1500 - 2100 m.

Fl. : Oct - Nov.

B. umbellatum Lindl., Gen. & Sp. Orchid. 56. 1830. *Cirrhopetalum maculosum* Lindl., Edw. Bot. Reg. 27. Misc. 81. 1841 : Hook. f., F.B.I. 5 : 776. 1890.

Epiphytes. Pseudobulbs ovoid, only when dried. Leaf narrowly oblong, obtuse and notched at apex. Flowers cream coloured.

Tendong ; ca 1500 m.

Fl. : May.

B. wallichii Reichb.f. in Walp. Ann. 6: 259. 1861 ; Seid. in Dansk. Bot. Ark. 29: 234. t. 127. 1973. *Cirrhopetalum refractum* sensu Hook. f., F.B.I. 5: 779. 1890, non (Reichb. f.) Zoll. 1847.

Epiphytic, caespitose herbs. Pseudobulbs concial, wrinkled during flowering, crowded. Leaves 2, narrowly oblong, acute. Flowers pale green, turning pale brown.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : Oct. - Nov.

B. yoksunense J.E. Sm. in Bull. Bot. Jard. Buitenz. 2, 8 : 29. 1912. *Cirrhopetalum brevipes* Hook.f., F.B.I. 5 : 777. 1890, non *B. brevipes* Ridl. 1903.

Epiphytes. Pseudobulbs oblong-cylindric. Leaves elliptic-oblong, coriaceous. Flowers rose coloured ; lips purple.

Sikkim, s.l. ; 1800 - 2400 m.

Fl. : Sept. - Oct.

B. xlophyllum Par. & Reichb. f. in Trans. Linn. Soc. Bot. 151. 1847 ; Hook. f., F.B.I. 5 : 766. 1890. **B. gracilipes** King & Pantl. in J. Asiat. Soc. Beng. 65 : 119. 1896.

Epiphytes without pseudobulb. Leaves solitary, oblanceolate or elliptic-oblanceolate, obtuse, minutely bifid at apex. Flowers dull purple.

Teesta valley ; ca 450 m.

Fl. : Sept. - Oct.

CALANTHE R. Br.

Calanthe alismaefolia Lindl., Fol. Orchid. 8. 1854 ; Hook. f., F.B.I. 5 : 849. 1890.

Terrestrial herbs. Flowers white ; lip with a pale violet blotch

Teesta valley ; 600 - 900 m.

Fl. : May - June.

C. alpina Hook. f. ex Lindl., Fol. Orchid. 4. 1854 ; Hook. f., F.B.I. 5 : 850. 1890.

Terrestrial herbs. Leaves oblanceolate. Flowers white, tipped with green ; lip dull red with yellowish base.

Lachen, Zemu ; 1800 - 3200 m.

Fl. : July - Aug.

C. angusta Lindl., Fol. Orchid. 7. 1854 ; Hook. f. F.B.I. 5 : 849. 1890.

Terrestrial herbs. Leaves narrow, elliptic. Flowers white.

Sikkim, s.l. ; ca 1050 m.

Fl. : April.

C. biloba Lindl., Fol. Orchid. 3. 1854 ; Hook. f., F.B.I. 5 : 848. 1890.

Terrestrial herbs. Leaves broadly elliptic. Flowers ochraceous yellow, mottled with purplish brown ; lip pale violet.

Sikkim, s.l. ; 1050 - 2100 m.

Fl. : Sept. - Oct.

C. brevicornu Lindl., Gen. & Sp. Orchid. 251. 1833 ; Hook. f., F.B.I. 5 : 854. 1890.

Terrestrial herbs. Leaves 3, subequal, oblong-elliptic. Flowers dull purple to orange with white lines ; lamellae of lip with yellow steaks.

Sedonchen-Kibek ; 800 - 2600 m.

Fl. : May - June.

C. chloroleuca Lindl., Fol. Orchid. 10. 1854 ; Hook. f., F.B.I. 5 : 852. 1890.

Terrestrial herbs. Leaves 2, elliptic-lanceolate. Flowers yellowish green ; lip white with an elongated yellow patch in middle.

Sikkim, s.l. ; 1800 - 2400 m.

Fl. : April - May.

C. densiflora Lindl., Gen. & Sp. Orchid. 250. 1833 ; Hook. f., F.B.I. 5 : 855. 1890.

Terrestrial herbs. Leaves narrowly elliptic. Flowers yellow.

Sikkim, s.l. ; 900 m.

Fl. : Oct.

C. gracilis Lindl., Gen. & Sp. Orchid. 251. 1833 ; Hook. f., F.B.I. 5 : 855. 1890.

Terrestrial herbs. Leaves 5-8, elliptic-oblong. Flowers white ; lip with yellow band in centre.

Sikkim, s.l. ; ca 1800 m.

Fl. : Oct.

C. herbacea Lindl., Fol. Orchid. 9. 1854 ; Hook. f., F.B.I. 5 : 852. 1890.

Terrestrial herbs. Leaves elliptic. Flowers green or yellowish.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : June.

C. mascua (D. Don) Lindl., Gen. & Sp. Orchid. 249. 1833 ; Hook. f., F.B.I. 5 : 850 : 1890. *Bletta mascua* D. Don, Prodr. Fl. Nepal. 30. 1825.

var. *mascua*

Terrestrial herbs. Leaves elliptic. Flowers violet ; lip yellow.

Tingbong, Gangtok ; 2000 - 2800 m.

var. *fulgens* (Lindl.) Hook. f., F.B.I. 5 : 851 1890. *C. fulgens* Lindl., Fol. Orchid. 10. 1854.

Sikkim, s. l. (cf. Hooker 1890)

C. puberula Lindl., Gen. & Sp. Orchid. 252. 1833 ; Hook. f., F.B.I. 5 : 848. 1890.

Terrestrial herbs. Leaves 5 - 7, unequal, elliptic-oblong. Flowers pale lavender ; lip darker, whitish at base.

Sikkim, s.l. ; 1800 - 2100 m.

/

Fl. : July - Aug.

C. tricarinata Lindl., Gen. & Sp. Orchid. 252. 1833 ; Hook. f., F.B.I. 5 : 847. 1890.

Terrestrial herbs with small broadly ovoid pseudobulbs. Flowers green with white edges.

Lachen valley ; 1500 - 3200 m.

C. treutleri Hook. f., F.B.I. 5 : 848. 1890.

Herbs. Leaves sessile. Flowers solitary, ca 3.5 cm across.

Sikkim Himalaya (*Treutler* K.). Represented by the type specimen only.

C. trulliformis King & Pantl. in J. Asiat. Soc. Beng. 64 : 337. t. 2. 1895.

Terrestrial herbs. Leaves sessile, linear-lanceolate. Flowers brown ; lip white.

Sikkim, s.l. ; 1800 - 2600 m.

Fl. : July.

C. uncata Lindl., Fol. Orchid. 6. 1854 ; Hook. f., F.B.I. 5 : 853. 1890.

Terrestrial herbs. Leaves broadly elliptic. Scape longer than leaves.

Sikkim, s.l. ; Himalaya (*Griffith's Collector*).

C. whiteana King & Pantl. in J. Asiat. Soc. Beng. 65 : 121. 1895.
Terrestrial herbs. Leaves linear-lanceolate. Flowers yellowish.

Chungthang ; ca 1800 m.

Fl. : May.

CEPHALANTHERA L.C. Rich.

Cephalanthera longifolia (L.) Fritsch in Ost. Bot. Z. 38 : 81. 1888. *Serapta helleborine* L. var. *longifolia* L., Sp. Pl. 950. 1753. *Cephalanthera ensifolia* L.C. Rich. in Mem. Mus. Hist. Nat. Paris. 4 : 60. 1818 ; Hook. f., F.B.I. 5 : 125. 1890.

Terrestrial, aphyllous herbs, 15 - 45 cm long. Roots tuberous. Flowers white, lip with a few yellow markings.

Near Gnathang ; 3600 m.

Fl. : July - Aug.

CEPHALANTHEROPSIS Guill.

Cephalantheropsis gracilis (Lindl.) Hu in Quart. J. Taiwan Mus. 25 : 213. 1972. *Calanthe gracilis* Lindl., Gen. & Sp. Orchid. 251. 1833 ; Hook. f., F.B.I. 5 : 855. 1890.

Terrestrial herbs. Leaves 5-8, elliptic or elliptic-oblong. Flowers white ; lip with a yellow band along centre.

Sikkim, s.l. ; ca 1800m.

Fl. : Oct.

CERATOSTYLIS Bl.

Ceratostylis subulata Bl., Bijdr. 7 : 206. 1825. *Appendicula teres* Griff., Notul. 3 : 359. 1851. *Ceratostylis teres* (Griff.) Reichb. f. in Bonap. 2 : 89. 1854.

Epiphytic herbs. Stems crowded. Pseudobulbs lacking. Flowers white ; lip yellow at end.

Teesta valley ; ca 300 m.

Fl. : May - June.

CHEIROSTYLIS Bl.

Cheirostylis griffithii Lindl. in J. Linn. Soc. 1 : 188. 1857 ; Hook. f., F.B.I. 6 : 105. 1890.

Terrestrial herbs. Stems decumbent and succulent at base. Leaves ovate, membranous. Flowers white.

Near Saramsa ; ca 1100 m.

Fl. : June.

CHILOSCHISTA Lindl.

Chiloschista lunifera (Reichb. f.) J.J. Sm. ex Holttum in Kew Bull. 14 : 273. 1960. *Thrixspermum luniferum* Reichb. f. in Gard. Chron. 786. 1868. *Sarcochilus luniferus* (Reichb. f.) Benth. ex Hook. f. in Curtis Bot. Mag. 115 : 7044. 1898 ; Hook. f., F.B.I. 6 : 37. 1890.

Stemless, leafless herb. Inflorescence 7 - 37 cm long, lax racemes. Flowers yellow with brown spots.

In tropical valley.

Fl. : May.

CHRYSOGLOSSUM Bl.

Chrysoglossum erraticum Hook. f., F.B.I. 5 : 784. 1890.

Terrestrial herbs. Pseudo bulbs ovoid. Flowers green, minutely spotted with brown ; lip white with a yellow spot at base and reddish spots on edges.

Sikkim, s.l. ; 900 - 1500 m.

Fl. : June.

CLEISOCENTRON Brühl

Cleisocentron trichromum (Reichb. f.) Brühl, Guide Orchid. Sikkim 137. 1926. *Saccolabium trichromum* Reichb. f. in Hamb. Gaertn. Zeit. 51. 1859 ; Hook. f., F.B.I. 6 : 65. 1890.

Pendulous herbs, 30 - 130 cm long. Leaves narrow-oblong. Inflorescence supra-axillary, much shorter than leaves, one sided, few flowered. Flowers straw coloured with a pink band in centre.

Sikkim, s.l. ; 300 - 750 m.

CLEISOSTOMA Bl.

Cleisostoma appendiculatum (Lindl.) Benth. & Hook. f. ex Jackson, Index Kew. 1 : 555. 1895. *Aerides appendiculatum* Lindl., Gen. & Sp. Orchid. 242. 1833. *Sarcanthus appendiculatus* (Lindl.) Par. in Mason, Burma, its people & products 2, bot. 181. 1883 ; Hook. f., F.B.I. 6 : 67. 1890.

Erect, epiphytic herbs, 20 - 40 cm tall. Leaves terete, spirally arranged. Flowers purplish.

In tropical valleys.

Fl. : Aug.

C. aspersum (Reichb. f.) Garay in Bot. Mus. Leafl., Harv. Univ. 23(4) : 169. 1972. *Sarcanthus aspersus* Reichb. f. in Hamb. Gartenz. 21 : 297. 1865 ; Hook. f., F.B.I. 6 : 70. 1890. *C. bicuspisidatum* Hook. f., l.c. 75. *Stereochilus bicuspisidatum* (Hook. f.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 236. t. 314. 1898. *Sarcanthus bicuspisidatus* (Hook. f.) J.J. Sm. in Nat. Tijds. Med. Ind. 72 : 84. 1912.

Epiphytic herbs. Leaves linear-oblong. Flowers yellowish, marked with pink along the midribs.

Sikkim, s.l. ; 600 m.

Fl. : July - Aug.

C. filiforme (Lindl.) Garay in Bot. Mus. Leafl., Harv. Univ. 23 (4) : 171. 1972. *Sarcanthus filiformis* Lindl., Bot. Reg. 28 : Misc. 61. 1842 ; Hook f., F.B.I. 6 : 66. 1890, *non* Wight 1851.

Pendulous epiphytic herbs, 25 - 180 cm long. Leaves on one side. Flowers brown, edges and midribs yellow ; lip white flushed with pink on apical lobe.

Tropical valleys.

Fl. : Aug. - Sept.

C. racemiferum (Lindl.) Garay in Bot. Mus. Leafl., Harv. Univ. 23 (4) : 173. 1972. *Saccolabium racemiferum* Lindl., Gen. & Sp. Orchid. 224. 1833. *Sarcanthus pallidus* Lindl. in Edw. Bot. Reg. Misc. 78. 1840 ; Hook f., F.B.I. 6 : 68. 1890.

Erect epiphytic herbs. Leaves strap like. Flowers light brown, edged with yellow ; lip white fleshy.

Sikkim, s.l. ; 600 - 1200 m.

Fl. : Aug.

C. subulatum Bl., Bijdr. 226. 1825. *Sarcanthus secundus* Griff., Notul. 3 : 362. 1851 ; Hook. f., F.B.I. 6 : 67. 1890. *S. subulatus* (Bl.) Reichb. f. in Bonpland. 5 : 41. 1857.

Pendulous epiphytic herbs, 15-30 cm long. Leaves linear-lanceolate. Flowers yellow with intramarginal brown bands.

In valleys.

C. striatum (Reichb. f.) Garay in Bot. Mus. Leafl., Harv. Univ. 23(4) : 175. 1972. *Echinoglossum striatum* Reichb. f. in Gard. Chron. (n.s.) 12 : 390. 1879 ; Hook. f., F.B.I. 6 : 76. 1890. *Cleisostoma brevipes* Hook. f., F.B.I. 6 : 73. 1890.

Pendulous epiphytic herbs. Leaves lanceolate. Spikes extra axillary, much shorter than leaves. Flowers yellow with brown bands ; lip pale buff, apical lobe white.

Tropical valleys.

Fl. : Aug.

COELOGYNE Lindl.

Coelogyne corymbosa Lindl., Fol. Orch. Coelog. 7. 1854 ; Hook. f., F.B.I. 5 : 831. 1890 ; King & Pantl., l.c. 8 : 134. t. 185. 1898. Das & Jain, Fascl. Fl. India 5 : 9. 1980.

Epiphytic herbs. Pseudobulbs ovoid or obovoid. Leaves lanceolate. Flowers white.

Dzongu, 1500 - 2400 m.

Fl. & *Fr.* : April - May.

C. cristata Lindl., Coll. Bot. Sub. t. 33 (*Textus*) 1821 ; Hook. f., F.B.I. 5 : 829. 1890 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 133. t. 184. 1898 ; Das & Jain, l.c. 9.

Epiphytic herbs. Pseudobulbs ovoid-oblong. Leaves subsessile, linear-oblong or linear-lanceolate. Flowers white.

Sikkim, s.l. ; 1500 - 2400 m.

Fl. & *Fr.* : April - June.

C. fimbriata Lindl., Bot. Reg. 11. t. 868. 1825 ; Hook. f., F.B.I. 5 : 836. 1890 ; Das & Jain, l.c. 10.

Epiphytes. Pseudobulbs oblong to orbicular, cylindric or fusiform. Flowers yellow.

Sikkim, s.l.

Fl. & Fr. : Aug. - Dec.

C. flaccida Lindl., Gen. & Sp. Orchid. 39. 1830 ; Hook. f., F.B.I. 5 : 829. 1890 ; King & Pantl., l.c. 133. t. 183 ; Das & Jain, l.c. 10.

Epiphytes. Pseudobulbs ridged, clustered. Flowers creamy white.

Sikkim, s.l. ; 750 - 1200 m.

Fl. : March - May.

C. fuscescens Lindl., Gen. & Sp. Orchid 41. 1830 ; Hook. f., l.c. 833 ; King & Pantl., l.c. 132. t. 181 ; Das & Jain, l.c. 13.

Epiphytes. Pseudobulbs cylindric. Leaves elliptic. Flowers pale brown.

Sikkim, s.l. ; 600 - 1800 m.

Fl. : Oct. - Nov.

C. longipes Lindl., Fol. Orch. Coelog. 10. 1854 ; Hook. f., F.B.I. 5 : 839. 1890, *non* King & Pantl. 1898 ; Das & Jain, l.c. 16.

Epiphytes. Pseudobulbs cylindric or ovoid. Flowers yellowish brown.

Sikkim, s.l. ; 1800 - 2100 m.

Fl. : May - June.

C. nitida (Wall. ex D. Don) Lindl., Gen. & Sp. Orchid. 40. 1830. *Cymbidium nitidum* Wall. ex D. Don, Prodr. Fl. Nep. 35. 1825, *non* Roxb. *C. ochracea* Lindl., Bot. Reg. t. 69. 1846 ; Hook. f., l.c. 831 ; King & Pantl., l.c. 132. t. 182 ; Das & Jain, l.c. 19.

Epiphytes. Pseudobulbs cylindric. Leaves lanceolate. Flowers white.

Yoksam, Dikchu, Gangtok, Karponang, Kabi, Ratong Chu, Tingling bridge ; 1200 - 2100 m.

Fl. : May - June.

C. occultata Hook. f., F.B.I. 5 : 831. 1890 ; King & Pantl. l.c. 135. t. 186. 1898.

Epiphytic herbs. Pseudobulbs ellipsoid or bipyramidal. Flowers white with some yellow streaks and a yellow blotch.

Sikkim, s.l. ; 600 - 1800 m.

Fl. : Sept. - Nov.

C. ovalis Lindl., Bot. Reg. 24 : 91. Misc. 191. 1838 ; Hook. f., F.B.I. 5 : 836. 1890 ; King & Pantl., l.c. 135. t. 187. 1898 ; Das & Jain., l.c. 23.

Epiphytes. Pseudobulbs narrowly ovoid to fusiform. Flowers pale brown.

Sikkim, s.l. ; 600 - 1800 m.

Fl. : Sept. - Nov.

C. prolifera Lindl., Gen. & Sp. Orchid. 40. 1830. *C. flavidia* Wall. ex. Lindl., Fol. Orch. Coelog. 10 : 1854 ; Hook. f., F.B.I. 5 : 838. 1890 ; Das & Jain, l.c. 23-24.

Epiphytes. Pseudobulbs oblong-ovoid. Leaves elliptic-oblong. Flowers yellow-brown.

Chungthang, Kaydoom (Lachung valley) Lachen ; 1800 - 2100 m.

Fl. : May - July.

C. punctulata Lindl., Coll. Bot. 7. t. 33. 1821 ; Das & Jain, l.c. 24.

Epiphytic herbs. Pseudobulbs ovoid. Flowers white.

Gangtok ; up to 1800 m.

Fl. : May - July.

C. raizadae Jain & Das in Proc. Ind. Acad. Sci. 87 B(5) : 119. 1978. *C. longipes* auct. non Lindl. 1854 ; King & Pantl., l.c. 136. t. 189. 1898.

Epiphytes. Leaves elliptic-lanceolate. Flowers pale green, flushed with yellow.

Raydong, 1800 - 2100 m.

Fl. : May.

C. schultesii Jain & Das in Proc. Ind. Acad. Sci. 87 B(5) : 119. 1978.

Epiphytes. Pseudobulbs ovoid to ellipsoid. Flowers yellowish brown.

Sikkim (*cf.* Das & Jain 1980).

Fl. : March - June.

C. stricta Schtr. in Fedde Repert. Beit. 184. 1919. *C. elata* Lindl., Gen. & Sp. Orchid. 40. 1830 ; Hook. f., l.c. 838 ; King & Pantl. l.c. 137. 1898.

Epiphytes. Pseudobulbs oblong-ovoid. Flowers white with ochre-yellowish tint.

Gangtok ; 1200 - 1800 m.

Fl. : May - June.

C. treutleri Hook. f., F.B.I. 5 : 837. 1890 ; Das & Jain, l.c. 30.

Epiphytes. Pseudobulbs ovoid. Flowers slightly produced.

Sikkim, *s.l.* Represented by the type specimen only.

C. uniflora Lindl., Gen. & Sp. Orchid. 42. 1830 ; Hook. f., F.B.I. 5 : 842. 1890 ; King & Pantl., l.c. 138. 1898 ; Das & Jain, l.c. 30.

Epiphytes. Leaves narrow-lanceolate. Flowers brownish.

Sikkim, *s.l.* ; up to 1200 m.

Fl. : April - May.

CORYBAS Salisb.

Corybas himalaicus (King & Pantl.) Schltr. in Fedde Repert. 19 : 19. 1924. *Corysanthes himalaica* King & Pantl. in J. Asiat. Soc. Beng. 65 : 128. 1896 ; Brühl, l.c. 158.

Terrestrial herbs, 5 - 7 cm tall. Flowers solitary, *ca* 1.5 cm long ; petals lacking ; lip and dorsal sepal transparent with purple markings.

Lachen valley at Lemteng ; *ca* 3000 m.

Notes : It grows on moist vertical rocks.

CORYMBORKIS Thou.

Corymborkis veratrifolia (Reinw.) Bl., Coll. Orch. Arch. Ind. 125. tt, 42. & 43. 1859. *Hysteria veratrifolia* Reinw. in Bot. Zeit. 2 : 5. 1825. *Corymbis veratrifolia* (Reinw.) Bl., l.c. H 42, E. 43, f. 1 ; Hook. f., F.B.I. 6 : 91. 1890 ; Brühl., l.c. 160.

Herbs, 90 - 120 cm tall. Leaves elliptic-lanceolate. Flowers white, fragrant.

Sikkim, s.l. ; 450 - 750 m.

CREMASTRA Lindl.

Cremastra appendiculata (D. Don) Makino in Bot. Mag. Tokyo 18 : 24. 1904. *Cymbidium appendiculatum* D. Don, Prodr. Fl. Nepal. 36. 1825. *Cremastra wallichiana* Lindl., Gen. & Sp. Orchid. 173. 1833 ; Hook. f., F.B.I. 6 : 16. 1890 ; Brühl, l.c. 112.

Herbs with large tubers covered with nodules. Leaves elliptic. Racemes conical. Flowers pink, crowded, drooping.

Sikkim, s.l. ; 1800 - 2400 m.

Fl. : May.

CRYPTOCHILUS Wall.

Cryptochilus lutea Lindl. in J. Linn. Soc. 3 : 21. 1859 ; Hook. f., F.B.I. 5 : 827. 1890.

Epiphytes with cylindric pseudobulbs. Flowers ca 6 mm long, nodding, yellow.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : May - June.

C. sanguineus Wall., Tent. Fl. Nepal. 36. t. 26. 1824 ; Hook. f., F.B.I. 5 : 827. 1890.

Epiphytes with oblong-ellipsoid or ovoid pseudobulbs. Flowers scarlet.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : June - Aug.

CYMBIDIUM Swartz

Cymbidium aloifolium (L.) Swartz in Nov. Act. Upsal. 2, 6 : 73. 1799 ; Hook. f., F.B.I. 6 : 10. 1890. *Epidendrum aloifolium* L., Sp. Pl. 953. 1753.

Epiphytic herbs. Leaves loriform, up to 90 x 5 cm. Flowers brownish red.

Singtam ; 300 - 1000 m.

Fl. : April - May.

Cult. in Deorali (Gangtok).

C. bicolor ssp. **obtusum** Du Puy & Cribb, The Genus *Cymbidium* 70.1988. *C. pendulum* (Roxb.) Swartz in Nov. Act. Upsal. 2, 6 : 73. 1799 ; Hook. f., F.B.I., 6 : 10. 1890. *Epidendrum pendulum* Roxb., Pl. Corom. 35. 1795.

Terrestrial herbs. Leaves 30-90 x 1.6-5 cm. Flowers dark purple-brown with yellowish margin.

Sikkim, s.l. ; ca 300 m.

Fl. : May.

C. cochleare Lindl. in J. Linn. Soc. 3 : 28. 1859. *Cyperorchis cochleare* (Lindl.) Benth. in J. Linn. Soc. 18 : 318. 1882 ; Hook. f., F.B.I. 6 : 15. 1889.

Terrestrial herbs. Flowers greenish brown.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : Oct. - Nov.

C. cyperifolium Lindl., Gen. & Sp. Orchid. 163. 1833 ; Hook. f., F.B.I. 6 : 13. 1890.

Terrestrial herbs. Flowers green ; lip spotted with red.

Chungthang ; ca 1500 m.

Fl. : Nov.

C. dayanum Reichb. f. in Gard. Chron. 710. 1869. *C. eburneum* Lindl. var. *dayana* (Reichb. f.) Hook. f., l.c. 6 : 12. 1890. *C. simonsianum* King & Pantl. in J. Asiat. Soc. Beng. 64 : 239. 1895.

Terrestrial herbs. Flowers white with crimson lines.

Teesta valley ; ca 3000 m.

Fl. : Aug.

C. devonianum Paxt. in Mag. Bot. 10. 97. cum. Ic. 1843 ; Hook. f., F.B.I. 6 : 10. 1890.

Terrestrial herbs. Flowers olive green, spotted with crimson.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : May - June.

C. eburneum Lindl. in Edw. Bot. Reg. n. s. 23. t. 67. 1847 ; Hook. f., F.B.I. 6 : 11. 1890.

Terrestrial herbs. Flowers white.

Sikkim, s.l. ; 300 - 900 m.

Fl. : March - April.

C. elegans Lindl., Gen. & Sp. Orchid. 161. 1833. *Cyperorchis elegans* (Lindl.) Bl., Rumph. 4 : 47. 1836 ; Fl. Jav. 4 : 93. t. 48c. 1858 ; Hook. f., F.B.I. 6 : 14. 1890.

Herbs. Leaves up to 60 × 2.5 cm. Racemes elongate ; flowers white.

Gangtok, Yoksam - Bakhim, Ratong - Chu Tingling Bridge ; 1500 - 1800 m.

Fl. : Oct - Nov.

C. ensifolium (L.) Swartz in Nov. Act. Upsal. 6 : 77. 1799 ; Hook. f., F.B.I. 6 : 13. 1890. *Epidendrum ensifolium* L., sp. pl. 90. 1753. *C. murnronianum* King & Pantl. in J. Asiat. Soc. Beng. 64 : 238. t. 2. 1895.

Terrestrial herbs. Flowers straw coloured, purple dotted.

Teesta valley ; ca 450 m.

Fl. : May.

Notes : Grows in dry knolls.

C. gammieanum King & Pantl. in J. Asiat. Soc. Beng. 64 : 330. 1895.

Terrestrial herbs. Flowers dull yellow, suffused with red flowers, flushed with brown lines.

Sikkim, s.l. ; 200 - 1200 m.

Fl. : Sept. - Oct.

C. hookerianum Reichb.f. in Gard. Chron. 1866. *C. grandiflorum* Griff., Notul. 3 : 321. 1851, *non* Swartz 1799 ; Hook. f., F.B.I. 6 : 12. 1890.

Terrestrial herbs. Flowers bright olive green, suffused with yellow.

Gangtok, Deorali ; 1500 - 2100 m.

Fl. : Feb.

Notes : Lip and column become red after fertilization.

C. lancifolium Hook., Exot. Fl. t. 51. 1823 ; Hook. f., F.B.I. 6 : 9. 1890.

Terrestrial herbs. Flowers greenish white with red stripes.

Gangtok (Deorali) ; up to 1800 m.

Fl. : June. Cult.

C. longifolium D. Don, Prodr. Fl. Nepal. 36. 1825 ; Hook. f., F.B.I. 6 : 13. 1890. *Cyperorchis longifolia* (D. Don) Schltr. in Seeds report 20 : 108. 1924.

Terrestrial herbs. Leaves 60 - 90 x 1.6 cm. Flowers greenish white with purple streaks.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : Sept. - Oct.

Notes : It can be distinguished from *C. giganteum* Wall. by its narrower leaves, narrower and differently coloured sepals and the lip which differs in colour and shape as well.

C. mastersii Griff. ex Lindl. in Bot. Reg. t. 50. 1845. *Cyperorchis mastersii* (Griff. ex Lindl.) Benth. in J. Linn. Soc. 18 : 318. 1881 ; Hook. f., F.B.I. 6 : 15. 1890.

Terrestrial herbs. Leaves up to 50 x 2 cm. Racemes short, decurved at tip. Flowers white, flushed with pink.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : Oct. - Nov.

C. sikkimense Hook. f., F.B.I. 6 : 9. 1890.

Epiphytic herbs. Racemes 10 - 20 cm long, somewhat decurved. Flowers white.

Lachen valley ; ca 1880 m.

Fl. : May.

C. tridioides D. Don, Prodr. Fl. Nepal. 36. 1825. *C. giganteum* Lindl., Gen. & Sp. Orchid. 163. 1833. *non* Swartz 1800 ; Hook. f., F.B.I. 6 : 12. 1890.

Terrestrial herbs. Flowers light yellow-green, suffused with red.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : Oct.- Nov.

Notes : Allied to *C. grandiflorum* but differs in size and colour of flowers and flowering time.

C. whiteae King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 193. t. 21. 258. 1898.

Terrestrial herbs. Flowers yellowish green, dull red-spotted.

Gangtok ; up to 1750 m.

Fl. : Nov.

CYPRIPEDIUM L.

Cypripedium elegans Reichb. f. in Flora 69 : 561. 1886 ; Hook. f., F.B.I. 6 : 169. 170. 1890.

Herbs, up to 10 cm tall. Leaves 2, opposite, orbicular. Flowers brownish with dark brown lines.

Lachen valley ; ca 3000 m.

Fl. : June - Aug.

C. himalaicum Rolfe *apud*. Hemsl. in J. Linn. Soc. 29 : 317. 1893.

Herbs. Leaves 3-4, broadly elliptic to lanceolate. Flowers brownish ; lip pendant, forming a broadly ovoid purple tinged pouch.

Thangu, Lachen ; 3000 - 4500 m.

Fl. : June - July.

C. macranthon Swartz in Act. Holm. 251. 1800 ; Hook. f., F.B.I. 6 : 170. 1890. *C. tibeticum* King *ex* Hemsl. in J. Linn. Soc. 29 : 320. 1892.

DENDROBIUM O. Swartz

Dendrobium aduncum Wall. in Bot. Reg. 1842. Misc. 62. 1846. t. 15. 1846 ; Hook. f. F.B.I. 5 : 730. 1890.

Epiphytic herbs. Leaves linear-lanceolate. Flowers pale purple, translucent, dark purple.

In Trop. valleys ; 600 m :

Fl. : June - Aug.

D. aggregatum var. **Jenkinsii** King & Pantl. in Ann. Roy Bot. Gard. Calc. Epiphytic herbs. Leaves solitary, oblong.

Fl. : March - May.

D. amoenum Wall. in Lindl., Gen. & Sp. Orchid, 78. (excl. syn *D. aphyllum*) 1830 ; Hook. f., F.B.I. 5 : 738. 1890.

Epiphytic herbs. Leaves linear-lanceolate, acuminate. Flowers white, with purple spots.

In trop valleys ; upto 1750 m ;

Fl. : June ; Common.

D. anceps Sw. in Act. Holm. 246. 1800 ; Hook. f., F.B.I. 5 : 724. 1890.

Stems compressed. Leaves lanceolate or ovate-lanceolate acute. Flowers singly borne on sides of stem, yellowish green.

Melli ; ca 350 m ;

Fl. : April - May.

D. aphyllum (Roxb.) C.E.C. Fischer in Gamble, Fl. Pres. Madras 1416. 1928. *D. pierardi* Roxb. in Hook., Exotic Flora t. 9. 1822 ; Hook. f., l.c. 738. *Limodorum aphyllum* Roxb., Pl. Corom. t. 41. 1795.

Epiphytic herbs. Leaves sessile, lanceolate-acuminate. Flowers rose coloured or pale purple ; lip yellow with radiating lines of purple at base.

In tropical valleys, Gangtok ; up to 1600 m.

Fl. : April - May.

D. bicameratum Lindl. in Edw. Bot. Reg. 59. 1839. Misc. 85 ; Hook. f., F.B.I. 5 : 729. 1890. *D. bolboflorum* Falc. ex Hook. f., l.c.

Epiphytic herbs. Leaves linear-lanceolate. Flowers yellow with red strips.

Deorali ; 1200 - 1800 m.

Fl. : Aug. - Sept.

D. candidum Wall. ex Lindl., Edw. Bot. Reg. 54. 1838 ; Hook. f., F.B.I. 5 : 735. 1890.

Epiphytic herbs. Leaves lanceolate. Flowers white with a yellow spot near the base.

In Lachen valley ; 1800 - 2100 m.

Fl. : May - June.

D. cathcartii Hook. *f.*, F.B.I. 5 : 727. 1890.

Epiphytic herbs. Leaves linear-lanceolate, unequally 2-fid. Flowers golden yellow ; lip marked with purple.

In tropical valleys ; ascending to 750 m.

Fl. : April - Sept.

D. chrysanthum Wall. *ex* Lindl., Gen. & Sp. Orchid. 80. 1830 & in Edw. Bot. Reg. 15. t. 1299. 1830 ; Hook. *f.*, F.B.I. 5 : 747. 1890.

Epiphytic herbs. Leaves lanceolate, acuminate. Flowers yellow ; lip with 2 dark, suborbicular brownish purple spots.

Sikkim, s.l. ; 900 - 1800 m.

Fl. : July - Oct.

D. crepidatum Lindl. in Paxt., Fl. Gard. 1. 63. f. 45. 1850 ; Hook. *f.*, F.B.I. 5 : 740. 1890.

Epiphytic herbs. Leaves linear-lanceolate, acute. Flowers pink or white ; lip with a large spot.

In tropical valleys, up to 1050 m.

Fl. : April - May. Cult. at Gangtok (Deorali).

D. cumulatum Lindl. in Gard. Chron. 1855. Hook. *f.*, F.B.I. 5 : 731. 1890.

Epiphytic herbs. Leaves oblong-lanceolate, acuminate. Flowers pale purple.

In tropical valleys ; up to 900 m.

Fl. : June - July.

Cult. at Deorali.

D. denneanum A.F.G. Kerr in J.Stam. Soc. Nat. Hist. Suppl. 9 : 229. 1933. *D. clavatum* Wall. *ex* Lindl. in Lindl. & Paxt., Fl. Gard. 2 : 104. t. 189. 1852, *non* Roxb. 1832 ; Hook. *f.*, F.B.I. 5 : 746. 1890.

Epiphytic herbs. Leaves linear-oblong, obtuse. Flowers dark yellow, lip with a large purple blotch near centre, edges pale yellow.

Tadong ; ca 900 m.

Fl. : May - June.

D. densiflorum Wall. ex Lindl. in Wall., Pl. As. Rar. 1 : 34. t. 40. 1830 ; Hook. f., F.B.I. 5 : 748. 1890.

Epiphytic herbs. Leaves subterminal, acute. Flowers yellow ; lip orange-yellow, paler towards margins.

Pamianchi, Saramsa-Gangtok ; up to 1800 m.

Fl. : April - May.

Notes : Unexpanded racemes have a cone-like appearance which is not found in any other species.

D. denudans D. Don, Prodr. Fl. Nep. 34. 1825 ; Hook. f., F.B.I. 5 : 715. 1890.

Epiphytic herbs. Leaves linear, oblong. Flowers pale green or whitish ; lip with a band of radiating red lines around its crisped or serrate edges ; foot or column red.

Sikkim, s.l. ; 900 - 1500 m. Common.

Fl. : Sept. - Oct.

D. eriaeaeformis Griff., Notul. III. 316. 1851 ; Hook. f., F.B.I. 5 : 715. 1890.

Epiphytic herbs. Leaves linear-lanceolate. Flowers green ; lip spotted with dull purple.

Sikkim, s.l. ; Deorali 900 - 1800 m.

Fl. : Sept. - Oct.

D. falconeri Hook. f. in Bot. Mag. 5 : 4944. 1856 ; Hook. f., F.B.I. 5 : 742. 1890.

Epiphytic herbs. Flowers white suffused with amethyst.

Deorali (Cult.) ; ca 1500 m.

Fl. : Feb. - May.

D. farmeri Paxt., Mag. Bot. 15 : 241. 1849 ; Hook. f., F.B.I. 5 : 750. 1890.

Epiphytes. Leaves 2 - 4, elliptic or oblong. Flowers pale mauve, changing to pure white after opening ; lip deep yellow, pale on margins.

Sikkim, s.l. ; 300 - 700 m.

Fl. : April - May.

D. formosum Roxb. (Wall. Cat. 1898) Fl. Ind. 3. 485. 1832 ; Hook. f., F.B.I. 5 : 721. 1890.

Epiphytic herbs. Leaves oblong. Flowers white ; lip yellow.

Sikkim, s.l. ; 350 - 1500 m.

Fl. : May. Cult. at Deorali.

D. gibsoni Lindl. in Paxt. Bot. Mag. 5 : 169. 1838 ; Hook. f., F.B.I. 5 : 746. 1890.

Epiphytic herbs. Leaves lanceolate, acuminate. Flowers saffron - yellow to orange.

Deorali ; 750 - 1200 m.

Fl. : July - Aug.

D. heterocarpum Wall. in Lindl., Gen. & Sp. Orchid. 78. 1830 ; Hook. f., F.B.I. 5 : 737. 1890.

Epiphytic herbs. Leaves oblong-lanceolate. Flowers faintly fragrant, white or pale ochraceous, often tinged with green ; lip yellow or white.

In tropical valleys ; 900 m.

Fl. : April.

D. hookerianum Lindl., Gen. & Sp. Orchid. 78. 1830 ; Hook. f., F.B.I. 5 : 745. 1890.

Epiphytes. Leaves oblong-lanceolate. Flowers golden yellow with 2 oblique dark purple patches on lip.

Sikkim, s.l. ; 1000 - 1500 m.

Fl. & Fr. : Sept.

D. longicornu Lindl. in Bot. Reg. 16 : t. 1315. 1830 ; Hook. f., F.B.I. 5 : 720. 1890.

Epiphytes. Leaves linear-lanceolate, unequally obtusely bifid. Flowers white.

Sikkim, s.l. ; 1200 - 2400 m.

Fl. : Sept. - Nov.

D. nobile Lindl., Gen. & Sp. Orchid. 24. 1830 ; Hook. f., F.B.I. 5 : 740. 1890.

Epiphytic herbs. Leaves oblong, obliquely notched. Flowers purple-white.

Kabi, Phodong, Gangtok, Pamianchi ; up to 1800 m.

Fl. : April - May.

D. ochreatum Lindl. in Bot. Reg. t. 1765. 1835 ; Hook. f., F.B.I. 5 : 747 - 748, 1890.

Epiphytic herbs. Leaves ovate-lanceolate, acuminate. Flowers golden yellow with a large maroon blotch.

Deorali (cf. Raju *et al* 1987-89).

Fl. : Feb. - April.

D. parishii Reichb. f. in Bot. Zeit. 21 : 237. 1863 ; Hook f., F.B.I. 5 : 740. 1890.

Epiphytic herbs. Leaves oblong-lanceolate. Flowers purplish.

Deorali (Cult.) ; 700 - 1300 m.

Fl. : June.

D. pauciflorum King & Pantl. in J. Asiat. Soc. Beng. 64(2) ; 332 t. 2. 1895.

Epiphytes. Flowers golden yellow ; sepals and petals broadly margined with purple ; apical lobe of lip red spotted.

Teesta valley ; 600 - 1200 m.

Fl. : June.

D. porphyrochilum Lindl. in J. Linn. Soc. III. 18. 1859 ; Hook. f., F.B. I. 5 : 716. 1890. *D. caespitosum* King & Pantl. in J. Asiat. Soc. Beng. 64 : 332. 1895.

Epiphytic herbs. Leaves obtuse or obliquely notched. Flowers yellow.

Dentam-Pamianchi ; 1500 - 2100 m.

Fl. : April - May.

D. primulinum Lindl. in Gard. Chron. 400. 1858 ; Hook. f., F.B.I. 5 : 735. 1890.

Epiphytes. Leaves broadly oblong. Flowers whitish pink.

Deorali (Cult.) ; up to 1200 m.

Fl. : April.

D. peguanum Lindl. in J. Proc. Linn. Soc. 3 : 19. 1859. *D. pygmaeum* Lindl., Gen. & Sp. Orchid. 85. 1830, *non* Sm. ex Rees 1808 ; Hook. f., l.c. 717. 1890.

Epiphytic herbs. Leaves 2 - 3, linear. Flowers white or purplish.

Sikkim, s.l. ; ca 300 m.

Fl. : Dec. - March.

D. ramosum Lindl., Gen. & Sp. Orchid. 82. 1830 ; Hook. f., F.B.I. 5 : 736. 1890.

Epiphytic herbs. Leaves lanceolate, acuminate. Flowers primrose yellow ; lip with numerous radiating brownish purple streaks.

Sikkim, s.l. ; 1200 - 1500 m.

Fl. : April - May.

D. stuposum Lindl. in Bot. Reg. 52. 1838 ; Hook. f., F.B.I. 5 : 737. 1890.

Epiphytic herbs. Leaves linear-oblong. Flowers white with a blotch of dull yellow on the lip.

Teesta valley ; ca 3600 m.

Fl. : June.

D. sulcatum Lindl. in Bot. Reg. n.s. 24. t. 65. 1838 ; Hook. f., F.B.I. 5 : 749. 1890.

Epiphytic herbs. Leaves subterminal, oblong, acuminate. Flowers dingy yellow with lines of a darker shade ; lip orange yellow with 2 patches of brown nerves near base.

Common in tropical valleys.

Fl. : April - May.

D. terminale Par. & Reichb. f. in Trans. Linn. Soc. 30 : 149. 1859 ; Hook. f., l.c. 5 : 725. 1890.

Epiphytic herbs. Leaves ovate-oblong, acute. Flowers whitish or pale pink.

Tropical valleys ; up to 1500 m.

Fl. : Sept. - Oct.

D. transprens Lindl., Gen. & Sp. Orchid. 79. 1830 ; Hook. f., l.c. 5 : 738. 1890.

Epiphytes. Leaves linear-lanceolate. Flowers white, sometimes suffused with pink ; lip with a large purple patch in middle and purple lines towards base ; column greenish yellow with a purple patch at lower end ; stigma red-lined.

Commonly found at foothills extending up to 900 m.

Fl. : May.

D. wardianum Warner, Sel. Orchid. 1, t. 19. 1862 ; Hook. f., F.B.I. 5 : 742. 1890.

Epiphytes. Leaves linear-lanceolate. Flowers reddish white ; lip ochraceous with maroon blotches.

Deorali (*cf.* Raju *et al.* 1987 - 89) ; 750 - 1200 m.

Fl. & *Fr.* : Feb. - March.

D. williamsoni Day & Reichb.f. in Gard. Chron. 78 : 1869 ; Hook. f., F.B.I. 5 : 721. 1890.

Epiphytes. Leaves oblong or lanceolate. Flowers yellowish white ; lip yellow, often dashed with red within.

Deorali (Cult.) ; 750 m.

Fl. : Feb. - March.

DIDICIEA King & Prain

Didiciea cunniighnami King & Pantl. in J. Asiat. Soc. Beng. 64 (2) : 119. 1895.

Terrestrial herbs. Pseudobulbs with a single ovate leaf arising from sides. Spikes lax - flowered, 2.5 - 3.8 cm long.

Lachen valley ; *ca* 3600 m.

Fl. : July.

DIDYMOPLEXIS Griff.

Didymoplexis pallens Griff. in Calc. J. Nat. Hist. 4 : 383. t. 17. 1843 ; Hook. f., F.B.I. 6 : 122. 1890.

Terrestrial aphyllous herbs. Flowers ochraceous white, tinged with brown.

In valleys.

Fl. : May - June.

DIGLYPHOSA Bl.

Diglyphosa macrophylla (King & Pantl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 98. 1898. *Chrysoglossum macrophyllum* King & Pantl. in J. Asiat. Soc. Beng. 64. 335. 1895.

Terrestrial herbs. Pseudobulbs subcylindric, broadly elliptic. Leaves elliptic, plicate. Flowers golden brown ; rachis purple.

Sikkim (?) ; *ca* 1200 m.

Fl. : May.

DIPHYLAX Hook. f.

Diphylax urceolata (Clarke) Hook. f. in Hook. Ic. Pl. 19. t. 1865. 1889. *Habenaria urceolata* Clarke in J. Linn. Soc. 25 : 73, t. 30. 1889 ; Hook. f., F.B.I. 6 : 165. 1890.

Terrestrial herbs, 5 - 15 cm tall. Leaves usually solitary, narrowly elliptic-oblong. Flowers white.

Yakla ; 3000 m.

DIPLOMERIS D. Don

Diplomeris hirsuta (Lindl.) Lindl., Gen. & Sp. Orchid. 330. 1835 ; Hook. f., F.B.I. 5 : 167. 1890. *Diplochilos hirsutus* Lindl. in Edw. Bot. Reg. 18. sub. t. 1499. 1832. 'Snow Orchid' (Eng.).

Small terrestrial herbs with globose tubers. Leaf solitary, sometimes with a much smaller one inserted at the base, linear-oblong, hirsute. Flowers white.

Near Phodong (on hill side walls).

Fl. : July - Aug., *Fr.* : Sept. - Oct.

DIPLOPRORA Hook. *f.*

Diploprora championii (Lindl.) Hook. *f.*, F.B.I. 6 : 26. 1890.
Cottonia championii Lindl. in Hook., J. Bot. 7 : 35. 1855.

Pendulous herbs. Leaves flaccidly fleshy, oblong-lanceolate. Flowers yellow ; lip white ; column white marked with cinnamon brown.

Sikkim ; ca 600 m.

Fl. : July - Aug.

EPIGENEIUM Gagnep.

Epigeneium amplum (Lindl.) Summerh. in Kew Bull. 1957 ; 260. 1957. *Dendrobium amplum* Lindl. in Wall., Pl. As. Rar. 1 : 25, t. 29. 1830 ; Hook. *f.*, F.B.I. 5 : 711. 1890 ; Brühl, l.c. 66.

Epiphytic herbs with pendulous rhizomes. Leaves 2, oblong or elliptic-oblong. Flowers pale greenish brown.

Sikkim, *s.l.* ; 1200 - 1800 m.

Fl. : Nov.

E. fuscescens (Griff.) Summerh. in Kew Bull. 1957 ; 264. 1957. *Dendrobium fuscescens* Griff., Notul. 3 : 308. 1851 ; Hook. *f.*, F.B.I. 5 : 712. 1890.

Epiphytes with pendulous rhizomes. Leaves 2, narrowly oblong-lanceolate. Flowers dull greyish brown.

Sikkim Himalaya ; 1500 - 2100 m.

Fl. : Oct.

E. rotundatum (Lindl.) Summerh. in Kew Bull. 1957. *Sacropodium rotundatum* Lindl., Fol. Orchid Sacropodium 2. 1853. *Dendrobium rotundatum* (Lindl.) Hook. *f.*, F.B.I. 5 : 712. 1890.

Epiphytes with pendulous rhizomes. Leaves 2, elliptic-oblong, notched at apex. Flowers brown.

Sikkim, *s.l.* ; 1800 - 2100 m.

Fl. : April.

EPIPACTIS Zinn

Epipactis helleborine (L.) Crantz, Strip. Austr. ed. 2 : 467. 1769. *emend.* Watson & Coulter in Asa Gray, Manual ed. 6 : 504. 1890.

Serapias helleborine L., Sp. Pl. 949. 1753. *Epipactis latifolia* Allioni, Fl. Ped. Mont. 2 : 151. 1785 ; Hook. f., F.B.I. 6 : 125. 1890.

Terrestrial herbs, up to 90 cm tall. Leaves broadly elliptic. Flowers green.

Gnathang ; ca 3300 m.

Fl. : July.

E. veratrifolia Boiss., Diagn. Pl. Orient. Nov. 1. 131. 11. 1854. *Epipactis consimilis* Wall. ex Hook. f., F.B.I. 6 : 126. 1890, non D. Don 1825.

Terrestrial herbs, up to 60 cm tall. Leaves lanceolate. Flowers lavender coloured.

Teesta valley ; ca 300 m.

Fl. : Feb. - March.

In damp soil.

EPIPOGIUM Gmelin ex Borkhausen

Epipogium aphyllum (F. W. Schmidt) Swartz, Summ. Veg. Scand. 32. 1814 ; Hook. f., l.c. 6 : 124. 1890. *Orchis aphylla* F.W. Schmidt in Mayer, Samml. Phys. Aufs. 240. 1791.

Terrestrial leafless yellowish herbs, up to 20 cm tall. Roots coralloid. Flowers white ; lip with red streaks on spur.

Lachen valley ; ca 3600 m.

E. roseum (D. Don) Lindl. in J. Linn. Soc. 1 : 177. 1857. *Limodorum roseum* D. Don, Prodr. Fl. Nep. 30. 1825. *Epipogium nutans* Reichb. f. in Bonpland. 36. 1857 ; Hook. f., F.B.I. 6 : 124. 1890.

Terrestrial herbs, up to 40 cm tall. Roots tuberous. Stem yellow. Flowers white ; lip with red-brown spots on upper surface.

In hot valleys ; up to 1200 m.

Fl. : April - June.

ERIA Lindl.

Eria acervata Lindl. in Paxt., Fl. Gard. 1 : 170. 1850 ; Hook. f., F.B.I. 5 : 796. 1890.

Epiphytic herbs. Pseudobulbs crowded, narrowly ellipsoid. Leaves 3 - 5, oblong-ob lanceolate. Flowers white, with yellow tinge on lip.

Deorali (Cult.) ; 550 - 750 m.

Fl. : June - July.

E. alba Lindl., Gen. & Sp. Orchid. 67. 1830 ; Hook. f., F.B.I. 5: 795. 1890.

Epiphytic herbs. Flowers yellowish white.

Deorali (Cult.) ; up to 1800 m.

Fl. : July - Aug.

E. amica Reichb. f., Gen. Orchid. 2 : 162. t. 168. f. 6 - 9. 1870 ; Hook. f., F.B.I. 5 : 800. 1890. *E. andersonii* Hook. f., l.c. 795. *E. confusa* Hook. f. in Hook., Ic. Pl. t. 1850, 1889 & l.c. 796.

Epiphytic herbs. Pseudobulbs clavate. Leaves 3 or more, oblong-lanceolate. Flowers dull yellow with reddish brown nerves.

Teesta valley ; ca 900 m.

Fl. : April.

E. bambusifolia Lindl. in J. Linn. Soc. 3 : 61. 1859 ; Hook. f., F.B.I. 5: 800. 1890.

Epiphytes, up to 90 cm long. Leaves many, elliptic-oblong. Flowers pale brown with darker lines.

Rissisoom ; ca 1800 m.

Fl. : Oct. - Dec.

E. biflora Griff., Notul. 3 : 302. 1851 ; Hook. f., l.c. 5 : 800. 1890.

Epiphytic herbs. Pseudobulbs tufted, stem like. Leaves 3 - 5, narrowly elliptic. Flowers pale yellow.

Deorali (Cult.) ; 350 - 1200 m.

Fl. : Sept.

E. bractescens Lindl. in Edw. Bot. Reg. 27. 1841 & Misc. 18 : 30, t. 29. 1844 ; Hook. f., F.B.I. 5 : 796. 1890.

Epiphytic herbs. Pseudobulbs narrowly cylindric. Leaves (2-) 3 - 6, elliptic-oblong. Flowers pale yellow.

Deorali (Cult.) ; 300 - 1000 m.

Fl. : April.

E. carinata Gibbs. in Calc. J. Nat. Hist. 5 : 365. 1865 ; Hook. f., F.B.I. 5 : 796. 1890.

Epiphytic herbs. Pseudobulbs oblong-ovoid, furrowed on maturity. Leaves elliptic - oblong. Flowers yellowish green and crimson veined.

Deorali (Cult.) ; 500 - 1200 m.

Fl. : Nov. - Jan.

E. clausa King & Pantl. in J. Asiat. Soc. Beng. 65 : 121. 1896.

Epiphytes. Pseudobulbs ovoid - ellipsoid. Leaves 2 - 3, narrowly lanceolate. Flowers green with brown stripes.

Deorali (Cult.) ; 1000 - 1500 m.

Fl. : Feb. - March.

E. coronaria (Lindl.) Reichb. f. in Walp. Ann. 272. 1861. *Codogyne coronaria* Lindl. in Bot. Reg. Misc. 83. 1841. *Trichosma suavis* Lindl. in Bot. Reg. 28. t. 21. 1842 ; Hook. f., F.B.I. 5 : 827. 1890.

Epiphytic herbs. Pseudobulbs tufted, thin, cylindric. Leaves 2, terminal, elliptic to ovate - lanceolate. Flowers white.

Deorali (Cult.) ; 1500 - 2000 m.

Fl. : Nov.

E. excavata Lindl., Gen. & Sp. Orchid. 67. 1830 ; Hook. f., F.B.I. 5 : 795. 1890.

Epiphytic herbs. Pseudobulbs oblong. Leaves 6, elliptic-lanceolate. Flowers dull white, with apical lobe of lip yellow.

Deorali (Cult.) ; 1200 - 2000 m.

Fl. : June.

E. ferruginea Lindl. in Edw. Bot. Reg. n.s.25 : t. 35. 1839 ; Hook. f., F.B.I. 5 : 804. 1890.

Epiphytic herbs. Leaves lanceolate, coriaceous. Flowers white.

Deorali (Gangtok). Cult. (*cf.* Raju *et al.* 1987)

E. graminifolia Lindl. in J. Proc. Linn. Soc. 3 : 54. 1859 ; Hook. f., F.B.I. 5 : 794. 1890.

Epiphytes. Pseudobulbs narrowly cylindric. Leaves 2 - 6, linear-lanceolate. Flowers white.

Sikkim, s.l. ; 1500 - 1800 m.

E. javanica Bl., Rumph. 2 : 23. 1836. *E. fragrans* Reichb. f. in Bot. Zeit. 22 : 415. 1864 ; Hook. f., F.B.I. 5 : 794. 1890.

Epiphytes. Pseudobulbs ellipsoid. Leaves in pairs, elliptic-oblong. Flowers white with some red spots in the apex of the column and side-lobes of lip.

Sikkim, s.l.

Fl. : July - Aug.

E. laniceps Reichb. f. in Hamb. Gratenzeit. 19 : 10. 1863. *E. elongata* Lindl. in J. Linn. Soc. 3 : 49. 1858 ; Hook. f., F.B.I. 5 : 801. 1890.

Epiphytes. Leaves 3 - 5, narrowly oblong. Flowers green ; lip purple in lower half.

Sikkim, s.l.

Fl. : April.

E. muscicola (Lindl.) Lindl. in J. Proc. Linn. Soc. 3 : 47. 1859 ; Hook. f., F.B.I. 5 : 789. 1890. *Dendrobium muscicola* Lindl., Gen. & Sp. Orchid. 75. 1830.

Epiphytic herbs, less than 5 cm tall. Pseudobulbs caespitose, depressed-globose. Leaves 2 - 4, oblanceolate. Flowers pale green.

Sikkim, s.l.

Fl. : April.

E. paniculata Lindl. in Wall., Pl. Asiat. Rar. 1 : 32. t. 36. 1830 ; Hook. f., F.B.I. 5 : 789. 1890.

Pendulous epiphytes. Stems caespitose, 30 - 60 cm long. Leaves many, fleshy, linear. Flowers yellowish green.

Sikkim, s.l.

Fl. : April - May.

E. pannnea Lindl. in Bot. Reg. 64. Misc. 79. 1842 ; Hook. f., F.B.I. 5 : 804. 1890.

Epiphytic wooly herbs. Leaves 1 - 4, cylindric. Sepals and petals white wooly outside ; lip brown.

Tropical valleys.

Fl. : May.

E. pubescens (Hook.) Lindl. in Edw. Bot. Reg. 11 : t. 904. 1825.
Dendrobium pubescens Hook., Exot. Fl. 2 : t. 124. 1824. *E. flava* Lindl.,
 Gen. & Sp. Orchid. 65. 1830 ; Hook. f., F.B.I. 5 : 801. 1890.

Epiphytes. Pseudobulbs furrowed, sheathed at base. Leaves 3 - 5, narrowly oblong. Flowers green, lip purple in basal portion.

In warm valleys.

Fl. : April.

E. pumila Lindl., Gen. & Sp. Orchid. 68. 1830 ; Hook. f., F.B.I. 5 : 791. 1890.

Epiphytes. Pseudobulbs crowded, subcylindric. Leaves 3 - 4, linear-lanceolate. Flowers white, flushed with pink.

In tropical valleys.

E. pusilla (Griff.) Lindl. in J. Linn. Soc. 3 : 48. 1859 ; Hook. f., l.c. 787. *Conchidium pusillum* Griff., Notul. 321. 1851.

Epiphytic herbs with solitary small flowers.

Deorali (Cult.) ; 1300 - 1800 m.

Fl. : May.

E. stricta Lindl., Coll. Bot. t. 41 B. 1825 ; Hook. f., F.B.I. 5 : 791. 1890.

Epiphytes. Pseudobulbs crowded, cylindric. Leaves 2, oblong-elliptic, sessile. Flowers white flushed with pink, column yellow.

Sikkim, s.l. ; 300 - 1200 m.

E. spicata (D. Don) Hand-Mazz., Symb. Sin. 7 : 352. 1936.
Octomeris spicata D. Don, Prodr. Fl. Nep. 31. 1825. *E. convallarioides* Lindl., Gen. & Sp. Orchid. 70. 1830 ; Hook. f., F.B.I. 5 : 791. 1890.

Epiphytes. Pseudobulbs flattened, tufted. Leaves 4 - 6, elliptic - lanceolate. Flowers white.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : March - April.

E. vittata Lindl. in J. Linn. Soc. 3 : 51. 1859 ; Hook. f., F.B.I. 5 : 794. 1890.

Epiphytes. Pseudobulbs cylindric. Leaves 2, elliptic - lanceolate. Flowers pale green, fragrant.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : March - April.

ESMERALDA Reichb. f.

Esmeralda clarkei Reichb. f. in Gard. Chron. 2 : 26. 552. 1886.
Arachnanthe clarkei (Reichb. f.) Rolfe in Gard. Chron. 2 : 567. 1888 ;
 Hook. f., F.B.I. 6 : 28. 1890.

Epiphytic pendulous herbs. Flowers white outside, yellow - brown inside.

Sikkim s.l. ; 1800 - 2400 m.

Fl. : Nov.

EULOPHIA R. Br.

Eulophia bicallosa (D. Don) P.F. Hunt. & Summerh. in Kew Bull. 20 : 10. 1966. **Bletia bicallosa** D. Don, Prodr. Fl. Nep. 30. 1825. **Cyrtopera bicarinata** Lindl., Gen. & Sp Orchid. 190. 1833. **Eulophia bicarinata** (Lindl.) Hook. f., F.B.I. 6 : 6. 1890.

var. bicallosa

Terrestrial herbs. Leaves linear, appearing after flowering. Flowers yellowish green, often streaked with pink, midlobe of lip orbicular, crisped.

Sikkim, s.l. ; 600 - 900 m.

Fl. : March - April.

var. major (King & Pantl.) Sarkar in J. Econ. Tax. Bot. 5 : 1007. 1984. **E. bicarinata** var. **major** King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 181. t. 244. 1898.

Differs from var. *bicallosa* in having flowers ca 3 cm across, pale yellowish green ; middle part of lip with 10 - 13 closely set tuberculate ridges of nearly equal length.

Foothills.

Fl. : March - April.

E. candida (Lindl.) Hook. *f.*, F.B.I. 6 : 6. 1890. *Cyrtopera candida* Lindl. in J. Linn. Soc. 3 : 31. 1858.

Terrestrial herbs. Leaves elliptic-lanceolate. Flowers white or greenish.

Sikkim, *s.l.* ; 900 - 1200 m (*cf.* Hooker 1890).

E. densiflora Lindl. in J. Linn. Soc. 3 : 25. 1858 ; Hook. *f.*, F.B.I. 6 : 5. 1890.

Terrestrial herbs. Scape robust. Racemes many flowered.

Foot hills (*cf.* Hooker *f.* 1890).

E. geniculata King & Pantl. in J. Asiatic. Soc. Beng. 64 : 337. 1895.

Terrestrial herbs. Leaves narrowly elliptic. Sepals brown ; petals and lip yellow.

Teesta valley ; *ca* 300 m.

Fl. : Aug.

E. graminea Lindl., Gen. & Sp. Orchid. 182. 1833 ; Hook. *f.*, F.B.I. 6 : 2. 1890.

Terrestrial herbs. Pseudobulbs obpyriform. Leaves linear. Flowers green with reddish nerves ; lip white, red on ridges.

Foot hills.

Fl. : March.

E. mannii (Reichb. *f.*) Hook. *f.*, F.B.I. 6 : 4. 1890. *Cyrtopera mannii* Reichb. *f.* in Flora 1872 : 274. 1872.

Terrestrial herbs, up to 2.2 m tall. Leaves linear. Flowers yellow, petals streaked with dull brown.

In tropical valleys.

Fl. : July - Aug.

E. macrobulbon Par. & Reichb. *f.* in Trans. Linn. Soc. 30 : 144. 1878 ; Hook. *f.*, F.B.I. 6 : 7. 1890.

Terrestrial herbs. Leaves sessile. Flowers yellow, petals streaked with dull brown.

In tropical valleys.

Fl. : July - Aug.

E. nuda Lindl., Gen. & Sp. Orchid. 180. 1833 ; Hook. f., F.B.I. 6 : 5. 1890.

Terrestrial herbs. Leaves elliptic - lanceolate. Flowers green, sepals with brown markings outside ; lip pale green at base, white in middle, pink flushed at apex.

In tropical valleys ; up to 450 m.

Fl. : April - July.

E. sanguinea (Lindl.) Hook. f., F.B.I. 6 : 8. 1890. *Cyrtopera sanguinea* Lindl. in J. Linn. Soc. 3 : 32. 1858. *E. macrorhizon* Hook. f., F.B.I. 6 : 7. 1890 ; King & Pantl., l.c. 179. 1898.

Terrestrial herbs. Leaves absent. Flowers dull brown ; lip white, pink or green.

Sikkim, s.l. ; ca 600 m.

Fl. : May - June.

FLICKINGERIA A.D. Hawkes

' **Flickingeria macraei** (Lindl.) Seid. in Dansk. Bot. Ark. 34 : 39. 1980. *Dendrobium macraei* Lindl., Gen. & Sp. Orchid. 75. 1833 ; Hook. f., l.c. 5. 714. p.p. ; Brühl, l.c. 66.

Epiphytes. Leaf terminal, sessile linear-oblong. Flowers 1 - 2 near leaf base, white.

In tropical valleys.

Fl. : May.

GALEOLA Lour.

Galeola cathcartii Hook. f., F.B.I. 6 : 89. 1890 ; King & Pantl., l.c. 263.

Scandant leafless herbs. Flowers yellow ; lip with red lines at base.

Rangeet valley ; ca 450 m.

Fl. : June.

Rare.

G. falconeri Hook. f., F.B.I. 6 : 88. 1890.

Erect, leafless herbs. Flowers bright yellow.

Sikkim, s.l. ; 1500 - 2100 m.

Fl. : July - Aug.

G. lindleyana (Reichb. f.) Reichb. f., Xen. Orch. 2 : 78. 1865 ; Hook. f., F.B.I. 6 : 88. 1890. *Erythrorchis lindleyana* Reichb. f. in Bonplandia 5 : 37. 1857.

Erect herbs. Stems brownish purple below, pale yellowish upwards. Flowers yellow with a reddish patch on the lip.

Sikkim s.l. ; 1500 - 2100 m.

G. nudiflora Lour., Fl. Cochinch. 2 : 520. 1790, *Galeola hydra* Reichb. f., Xen. Orchid. 2 : 77. 1865 ; Hook. f., F.B.I. 6 : 89. 1890.

Extensive leafless climber. Flowers yellow with red markings at the base of the lip.

Rishap valley.

Fl. : July.

GASTROCHILUS D. Don

Gastrochilus acutifolius (Lindl.) O. Ktze., Rev. Gen. Pl. 2 : 661. 1891. *Saccolabium acutifolium* Lindl., Gen. & Sp. Orchid. 223. 1833 ; Hook. f., F.B.I. 6 : 61. 1890 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 228. t. 304. 1898.

Epiphytes without pseudobulbs. Leaves fleshy, oblong-lanceolate. Flowers pale green, flushed and mottled with dull brown or uniformly yellow ; lip white.

Sikkim, s.l., ca 900 m.

Fl. : Nov. - Dec.

G. affinis (King & Pantl.) Schltr. in Fedde Repert. 12 : 314. 1913. *Saccolabium affine* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 228. t. 304. 1898.

Epiphytes. Flowers green, flushed with brown ; lip yellowish.

Above Pemiongtsi ; ca 2400 m.

Fl. : June.

G. calceolaris (J.E. Smith) D. Don, Prodr. Fl. Nep. 32. 1825.
Saccolabium calceolare (J.E. Smith) Lindl., Gen. & Sp. Orchid. 223. 1833 ; King & Pantl., l.c. 277. 1898.

Pendulous epiphytes. Leaves narrowly oblong, with unequally bifid apices. Flowers pale green with large brown markings ; lip-sac yellow with brown red markings.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : March - April.

G. dasypogon (J.E. Smith) O. Ktze., Rev. Gen. Pl. 2 : 66. 1891.
Aerides dasypogon J.E. Smith in Rees. Cyclop. 39 : 10. 1819. *Saccolabium dasypogon* (J.E. Smith) Lindl., Gen. & Sp. Orchid. 222. 1833 ; Hook. f., F.B.I. 6 : 66. 1890 ; King & Pantl., l.c. 224.

Pendulous epiphytes. Leaves narrowly oblong. Flowers bright yellow.

Sikkim, s.l. ; ca 300 m.

Fl. : Nov. - Dec.

G. distichus (Lindl.) O. Ktze., Rev. Gen. Pl. 2 : 661. 1891.
Saccolabium distichum Lindl. in J. Proc. Linn. Soc. 3 : 36. 1859 ; Hook. f., F.B.I. 6 : 64. 1890.

Epiphytes. Leaves fleshy, lanceolate. Flowers green, with irregular brown markings ; apical lobe of lip yellow.

Sikkim, s.l. ; 1800 - 2400 m.

Fl. : March - May.

G. intermedius (Griff. ex Lindl.) O. Ktze., Rev. Gen. Pl. 2 : 661. 1891. *Saccolabium intermedium* Griff. ex Lindl. in J. Linn. Soc. 3 : 33. 1858 ; Hook. f., F.B.I. 6 : 61. 1890.

Epiphytic herbs. Leaves linear. Flowers yellow, spotted with brownish purple.

Sikkim, s.l. ; 900 m.

Fl. : Aug.

G. pseudodistichus (King & Pantl.) Seid., Orch. Fl. Camb. Laos & Vietn. 1 : 65. 1975. *Saccolabium pseudodistichum* King & Pantl. in J. Asiat. Soc. Beng. 64 : 341. 1895.

Terrestrial herbs. Leaves lanceolate. Flowers greenish with brownish purple spots.

Sikkim, s.l. ; 1800 - 2400 m.

Fl. : Aug. - Oct.

GASTRODIA R. Br.

Gastrodia dyeriana King & Pantl. in J. Asiat. Soc. Beng. 64 : 342. 1895.

Terrestrial leafless herbs. All parts brown except the yellowish white lip.

Sikkim, s.l. ; ca 2100 m.

Fl. : Aug.

GEODORUM G. Jacks

Geodorum densiflorum (Lam.) Schltr. in Fedde Report. 4 : 259. 1929. *Limodorum densiflorum* Lam., Encycl. 3 : 516. 1791 - 92. *Geodorum dilatatum sensu* Hook. f., F.B.I. 6 : 17. 1890, non R. Br. 1813. *G. purpureum* R. Br. in Hort. Kew. ed. 295 : 207. 1813 ; Hook. f., l.c. 16 ; Brühl, l.c. 112.

Terrestrial herbs. Tubers globular, apiculate. Leaves elliptic. Flowers white or pale purple ; callii on the lip yellow.

Found at lower elevations.

Fl. : April - May.

GOODYERA R. Br.

Goodyera foliosa (Lindl.) Benth. ex Clarke in J. Linn. Soc. (Bot.) 25 : 73. 1889 ; Hook. f., F.B.I. 6 : 113. 1890. *Georchis foliosa* Lindl., Gen. & Sp. Orchid. 496. 1840.

Terrestrial herbs, up to 30 cm tall. Leaves ovate-lanceolate. Flowers pink or orange ; lip yellow.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : Oct. - Nov.

G. fumata Thw., Enum. Pl. Zeyl. 314. 1864 ; Hook. f., F.B.I. 6 : 111. 1890.

Terrestrial herbs, up to 1.2 m tall. Leaves obliquely oblong-elliptic. Flowers pale green ; lip and column pale brown.

Sikkim (?) ; ca 1800 m.

Fl. : March.

G. fusca (Lindl.) Hook. f., F.B.I. 6 : 112. 1890. *Aetheria fusca* Lindl., Gen. & Sp. Orchid. 491. 1840.

Terrestrial herbs, up to 30 cm tall. Leaves ovate. Flowers white ; apex of lip green, sac flushed with pinkish yellow.

Lachung valley ; ca 3600 m.

Fl. : Aug.

G. grandis King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 284. 1898.

Terrestrial herbs, up to 50 cm tall. Leaves ovate. Flowers dull red.

Rumtek ; ca 1200 m.

Fl. : Aug.

G. hemsleyana King & Pantl. in J. Asiat. Soc. Beng. 64 : 342. 1895.

Terrestrial herbs, up to 25 cm tall. Leaves 3 - 5, unequal, broadly ovate. Flowers white ; sepals tipped with pink.

Sikkim, s.l. ; 2100 m.

Fl. : July.

G. hispida Lindl. in J. Linn. Soc. 1 : 183. 1857 ; Hook. f., F.B.I. 6 : 114. 1890.

Terrestrial herbs, 10 - 15 cm tall. Leaves elliptic lanceolate. Flowers greenish white.

Teesta valley ; ca 300 m.

Fl. : Sept.

G. procera (Wall. ex Ker-Gawl.) Hook., Exot. Fl. 1 : t. 39. 1823 ; Hook. f., F.B.I. 6 : 111. 1890. *Neottia procera* Wall. ex Kar. - Gawl. in Edw. Bot. Reg. 8 : t. 639. 1822.

Terrestrial herbs, 25 - 75 cm tall. Leaves narrowly elliptic or ovate-elliptic. Flowers white, faintly tinged with green.

Sikkim, s.l. ; up to 1200 m.

Fl. : May.

G. repens (L.) R. Br. in Ait., Hort. Kew., ed. 2. 5 : 198. 1813 ; Hook. f., F.B.I. 6 : 111. 1890. *Satyrium repens* L., Sp. Pl. 945. 1753.

Terrestrial herbs, up to 20 cm tall. Leaves ovate-elliptic, green, reticulated with white. Flowers white flushed with brownish pink.

Sikkim, s.l. ; 2400 - 3300 m.

Fl. : July - Aug.

G. schlechtendaliana Reichb. f. in Linnaea 22 : 861. 1849. *Goodyera secundiflora* Lindl. in J. Linn. Soc. 1 : 82. 1853 ; Hook. f., F.B.I. 6 : 113. 1890.

Terrestrial herbs, 15 - 30 cm tall. Leaves ovate-lanceolate. Flowers yellowish pink ; lip white, tipped with green.

Sikkim, s.l. ; ca 1800 m.

Fl. : Sept. - Jan.

G. vittata Benth. ex Hook. f., F.B.I. 6 : 113. 1890 ; Brühl. l.c. 165.

Terrestrial herbs, up to 20 cm tall. Leaves ovate-lanceolate. Flowers pink.

Lachung valley, 1800 - 3000 m.

Fl. : July - Sept.

GYMNADENIA R. Br.

Gymnadenia orchidis Lindl., Gen. & Sp. Orchid. 278. 1835.

Habenaria orchidis (Lindl.) Hook. f., F.B.I. 6 : 142. 1890. *Orchis habenarioides* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 302. t. 401. 1898.

Terrestrial herbs, up to 50 cm tall. Leaves 4 - 5, elliptic-oblong to linear-oblong, sessile. Flowers pinkish or violet, rarely white.

Lachung valley ; ca 3600 m.

Fl. : Aug.

HABENARIA Willd.

Habenaria aitchisoni Reichb. f. apud Aitch. in Trans. Linn. Soc.

2. Bot. 3. 113. 1888 ; Hook. f., F.B.I. 6 : 152. 1890.

Terrestrial herbs, up to 12.5 cm tall. Leaves 2, sessile, orbicular. Flowers greenish.

Lachen valley ; 3000 - 3300 m.

Fl. : July.

H. albomarginata King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 322. t. 425. 1898.

Terrestrial herbs, less than 20 cm tall. Leaves 2, unequal, orbicular-elliptic to oblong. Flowers green with white margins.

Near Dzongri ; *ca* 3900 m.

Fl. : June.

H. arietina Hook. f., F.B.I. 6 : 138. 1890.

Terrestrial herbs. Leaves narrowly elliptic. Flowers green.

Tumlong, Pemiongchi, Re - chi - la ; 1800 - 2700 m.

Fl. : July - Aug.

H. cumminsiana King & Pantl. in J. Asiat. Soc. Beng. 64 : 343. t. 2. 1895.

Terrestrial herbs, 15 -25 cm tall. Leaves 2 - 4, unequal, broadly elliptic to lanceolate. Flowers green.

Gnathang, Lachung valley ; *ca* 3300 m.

Fl. : July.

H. dentata (Swartz) Schltr. in Fedde Repert. 4 : 125. 1919.
Orchis dentata Swartz in Vet. Acad. Handl. Stockh. 21 : 207. 1800.
Habenaria geniculata D.Don, Prodr. Fl. Nep. 25. 1825 ; Hook. f., l.c. 6 : 138. 1890.

f. dentata

Terrestrial herbs, 30 - 75 cm tall. Leaves oblong-elliptic. Flowers white.

Sikkim, *s.l.* ; 750 - 1500 m.

f. ecalcarata (King & Pantl.) Tuyama in Hara, Fl. E. Himal. 438. 1966. *H. geniculata* var. *ecalcarata* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 320. t. 405, f. 4 - 5. 1898.

Differs from *f. dentata* in having united stigmas, lanceolate-oblong lip and spur being absent.

Sikkim, s.l. ; 750 - 1500 m.

H. malleifera Hook. f., F.B.I. 6 : 143. 1890. *H. furfuracea* Hook. f., l.c. 144.

Terrestrial herbs, 37 - 60 cm tall. Leaves narrowly elliptic. Flowers green.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : Aug. - Sept.

H. nematocaulon Hook. f., F.B.I. 6 : 154. 1890.

Terrestrial herbs, 6.5 - 22.5 cm tall. Leaves elliptic. Flowers white.

Rechi - la ; ca 2700 m.

Fl. : Aug.

H. pectinata (J.E. Smith) D. Don, Prodr. Fl. Nepal. 24 : 1825 ; Hook. f., F.B.I. 6 : 137. 1890. *Orchis pectinata* J.E. Smith, Exot. Bot. t. 99. 1804.

Terrestrial herbs, 30 - 75 cm tall. Leaves linear - lanceolate, distributed in upper part of the stem.

North Sikkim ; ca 1500 m.

Fl. : Aug.

H. stenopetala Lindl., Gen. & Sp. Orchid. 319. 1835 ; Hook. f., F.B.I. 6 : 134. 1890.

Terrestrial herbs, 30 - 105 cm tall ; stem sheathed at lower portion. Leaves distributed above, oblong-lanceolate or ovate-lanceolate. Flowers pale green ; lip brown.

Sikkim, s.l. ; 600 - 1800 m.

Fl. : Aug. - Oct.

H. seshagiriana A.N. Rao in J. Econ. Tax. Bot. 6 : 223. 1895. *H. stenopetala* var. *polytricha* Hook. f. in Ann. Roy. Bot. Gard. Calc. 5 : 64. t. 96. 1895. *H. polytricha* (Hook. f.) Pradhan in Ind. Orchid. 1 : 68. 1976, *non* Rolfe 1896.

Terrestrial herbs. Sepals and petals produced into long capillary tails.

Sikkim, s.l. ; ca 1500 m.

Fl. : Aug. - Sept.

HERMINIUM L.

Herminium angustibre King & Pantl. in J. Asiat. Soc. Beng. 65 : 131. 1896 ; Brühl, l.c. 191. 1926.

Terrestrial herbs, up to 10 cm tall. Leaf solitary, sessile, elliptic. Flowers green.

Lingtu, Singa - le - la ; 3300 - 3600 m.

Fl. : July.

H. jeffreyanum King & Pantl. in J. Asiat. Soc. Beng. 65 : 130. 1896.

Terrestrial herbs, 10 - 20 cm tall. Leaves 2, narrowly oblong. Flowers green.

Sikkim, s.l. ; 2400 - 3000 m.

Fl. : Aug.

H. josephii Reichb. f. in Flora 55 : 276. 1872 ; Hook. f., F.B.I. 6 : 129. 1890.

Terrestrial herbs, 10 - 25 cm tall. Leaves 2, sub-opposite, narrowly elliptic-oblong. Flowers green.

Lachen valley ; 3300 - 3600 m.

Fl. : June - July.

H. lanceum (Thunb. ex Swartz) J. Vuijk. in Blumea 11 : 228. 1961. *Ophrys lancea* Thunb. ex Swartz in Kongl., Vet. Akad. Handl. Stockh. 21 : 223. 1800. *Aceras angustifolia* Lindl. in Edw. Bot. Reg. 10 : t. 1525. 1832. *Herminium angustifolium* (Lindl.) Benth. ex Hook. f., F.B.I. 6 : 129. 1890.

Terrestrial herbs, 25 - 75 cm tall. Leaves 2 - 3, linear, acuminate. Flowers green.

Sikkim, s.l. ; 1350 - 2700 m.

Fl. : July - Aug.

H. macrophyllum (D. Don) Dandy in J. Bot. 70 : 328. 1930. *Neottia macrophylla* D. Don, Prodr. Fl. Nepal. 27. 1825. *Herminium congestum* Lindl. in Edw. Bot. Reg. 18 : Sub. t. 1499. 1832 ; Hook. f., F.B.I. 6 : 130. 1890 ; Brühl, l.c. 193.

Terrestrial herbs, 7.5 - 17.5 cm tall. Leaves 2 - 3, narrowly oblong-elliptic. Flowers green.

Sikkim, s.l. ; 3000 - 3750 m.

Fl. : July - Aug.

H. orbiculare Hook. f., F.B.I. 6 : 130. 1890 ; Brühl, l.c. 191.

Terrestrial herbs, 7 - 15 cm tall. Leaves variable, elliptic-oblong, or linear-lanceolate. Flowers green.

Singale - la ; 3600 m.

Fl. : July.

H. pugioniformae Lindl. ex Hook.f., F.B.I. 6 : 130. 1890 ; Brühl, l.c. 191.

Terrestrial herbs, 5 - 18 cm tall. Leaf solitary, oblong-oblanceolate. Flowers minute, green.

Singale - la ; 3600 m.

Fl. : July.

H. quinquelobium King & Pantl. in J. Asiat. Soc. Beng. 65 : 130. 1896 ; Brühl, l.c. 193.

Terrestrial herbs, 10 - 30 cm tall. Leaves 2, linear-oblong. Flowers green ; lip 5-lobed.

Tankrala, Tendong ; 2100 - 2850 m.

Fl. : Aug.

HERPYSMA Lindl.

Herpysma longicaulis Lindl., Gen. & Sp. Orchid. 506. 1840 ; Hook. f., F.B.I. 6 : 98. 1890.

Terrestrial herbs. Leaves ovate - lanceolate or narrowly elliptic. Flowers white.

Sikkim, s.l. ; 1500 m.

Fl. : Aug.

HETAERIA Bl.

Hetaeria rubens (Lindl.) Benth. in Benth. & Hook. f., Gen. Pl. 3 : 164. 1880 ; Hook. f., F.B.I. 6 : 115. 1890 ; Brühl, l.c. 174.

Terrestrial herbs, 25 - 45 cm tall. Leaves ovate-elliptic, sub-oblique. Flowers green, tipped with pink ; lip white tipped.

Sikkim, s.l. ; 450 - 900 m.

Fl. : Jan - Feb.

KINGIDIUM P.F. Hunt.

Kingidium deliosum (Reichb. f.) Sweet in Amer. Orchid. Soc. Bull. 39 : 1095. 1970. *Phalaenopsis deliosa* Reichb. f. in Bonplandia 2 : 93. 1854. *P. wightii* Reichb. f. in Bot. Zeit. 214. 1854. *Aerides decumbens* Griff., Notul. 3 : 365. 1851 & Ic. Pl. Asiat. t. 320. 1851. *Doritis wightii* (Reichb. f.) Benth. in Benth. & Hook. f., Gen. Pl. 3 : 574. 1883 ; Hook. f., F.B.I. 6 : 32. 1890 ; King & Pantl. in Ann. Roy Bot. Gard. Calc. 8 : 198. 1898.

Epiphytic herbs. Leaves ca 3, pendulous, broadly oblanceolate, blunt, 15 - 20 x 3 - 5 cm. Flowers ca 15 mm across, pale ochraceous yellow with purple markings on lip and column, base of perianth slightly mottled with pink.

Teesta valley, at lower elevations.

Fl. : July.

K. taenialis (Lindl.) P.F. Hunt. in Kew Bull. 24 : 98. 1970. *Aerides taeniale* Lindl., Gen. & Sp. Orchid. 239. 1833. *Doritis taenialis* (Lindl.) Hook. f., F.B.I. 6 : 31. 1890 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 199. 1898.

Epiphytic herbs. Leaves 1 or more, pendulous narrowly elliptic-oblong, 70 - 130 x 15 - 20 mm, acute. Flowers pale purple, ca 20 mm across.

Gangtok (Baluakhani) ; 900 - 1800 m.

Fl. : May - July.

LIPARIS L.C. Rich.

Liparis bhootanensis Griff., Itin. Notes 98. 1848 ; Hook. f., F.B.I. 5 : 700. 1890.

Epiphytes. Pseudobulbs narrowly oblong. Leaf solitary, linear-oblong to lanceolate. Flowers ca 16 mm long, pale brown.

Sikkim, s.l. ; ca 1340 m.

Fl. : Aug.

L. bistriata Parish & Reichb. f. in Trans. Linn. Soc. 30 : 155. 1874 ; Hook. f., F.B.I. 5 : 702. 1890.

Epiphytes. Pseudobulbs elongated, 7.5 - 15 cm long, tufted. Leaves 2, elliptic-lanceolate or elliptic-oblong. Flowers *ca* 12 mm long, pale green.

Rishap ; *ca* 2100 m.

Fl. : July - Aug.

L. caespitosa (Thou.) Lindl., Bot. Ref. 11. subt. 882. 1825. *Malaxis caespitosa* Thou., Orch. Ill. Afr. t. 89. 1882. *L. pusilla* Ridley in J. Linn. Soc. 22 : 294. 1886 ; Hook. f., F.B.I. 5 : 701. 1890. *L. obscura* Hook. f., F.B.I. 5 : 701. 1890.

Epiphytes. Pseudobulbs tufted, ellipsoid or ovoid, *ca* 12 mm long. Leaf solitary, linear-lanceolate or linear-oblanceolate. Flowers pale green, *ca* 3 mm long.

Teesta valley ; 1200 - 2100 m.

Fl. : Aug. - Oct.

L. cathcartii Hook. f. in Hook., Ic. Pl. 19 : f. 1808. 1889 & F.B.I. 5 : 694. 1890.

Terrestrial herbs. Pseudobulbs narrowly ovoid. Leaves 2, sub-opposite, broadly ovate or elliptic ovate. Flowers *ca* 2 cm long, purplish green.

Lachen valley ; 2100 - 3000 m.

L. cordifolia Hook. f. in Hook., Ic. Pl. 19 : t. 1811. 1889 & F.B.I. 5 : 692. 1890.

Terrestrial herbs, dark green. Pseudobulbs crowded, compressed, ovoid. Leaf large, solitary, broadly ovate. Flowers *ca* 18 mm ; lip almost deltoid, broad at apex.

Sikkim, *s.l.* ; 1500 - 1800 m.

Fl. : Oct. - Nov.

L. deflexa Hook. f., F.B.I. 5 : 697. 1890.

Terrestrial herbs. Leaves 2, subopposite, elliptic. Flowers *ca* 7 mm long, yellow.

Sikkim, *s.l.* ; *ca* 600 m.

Fl. : Aug.

L. deliculata Hook. f., Ic. Pl. t. 1889 & F.B.I. 5 : 705. 1890.

Epiphytes. Pseudobulbs ovoid or oblong. Leaves 2, narrowly elliptic. Flowers *ca* 6 mm long., ochraceous.

Sikkim, *s.l.* ; 1800 m.

Fl. : Aug. - Sept.

L. gamblei Hook. *f.*, Ic. Pl. t. 1812. 1889 & F.B.I. 5 : 693. 1890.

Epiphytes. Pseudobulbs in clusters, suborbicular, flattened, green. Leaf solitary, elliptic or elliptic-oblong. Flowers dark green ; lip with brown-purple margins and nerves.

Sikkim, *s.l.* ; 1800 - 1950 m.

Fl. : June.

L. glossula Reichb. *f.* in Linnaea 41 : 44. 1876 ; Hook. *f.*, F.B.I. 5 : 693. 1890.

Terrestrial herbs. Pseudobulbs ovoid, tufted. Leaf solitary, oblong. Flowers *ca* 18 mm long, green ; lip purple.

Sikkim, *s.l.* ; 1800 - 4300 m.

L. nervosa (Thunb.) Lindl., Gen. & Sp. Orchid. 26. 1830. *Ophrys nervosa* Thunb., Fl. Japon. 27 : 1784. *L. bituberculata* Lindl. in Edw. Bot. Reg. 11. sub f. t. 882. 1825 ; Hook. *f.*, F.B.I. 5 : 696. 1890.

Terrestrial herbs. Stems tufted, up to 25 cm tall. Leaves 4 - 5, ovate-elliptic or lanceolate, opposite, oblique, plicate. Flowers *ca* 18 mm long, brownish purple.

Chungthang, Lachen valley ; *ca* 1800 m.

Fl. : June - July.

L. paradoxa (Lindl.) Reichb. *f.*, Walp. Ann. 6 : 218. 1861 ; Hook. *f.*, F.B.I. 5 : 697. 1890. *Empusa paradoxa* Lindl., Bot. Reg. 10. Subt. 825. 1824.

Terrestrial herbs. Pseudobulbs ovoid. Leaves 2 - 3, alternate, elliptic-lanceolate. Flowers *ca* 12 mm long, green ; lip purplish.

Tropical valleys ; up to 1800 m.

Fl. : July - Sept.

L. perpusilla Hook. *f.* in Hook. *f.* Ic. Pl. 19. t. 1856 B. 1889 & F.B.I. 5 : 706. 1890. *L. togashii* Tuyama ex Hara, Fl. E. Himal. 441. 1966.

Epiphytes. Pseudobulbs tufted, ovoid-oblong. Leaves 3 - 4, linear or linear-oblanceolate. Flowers yellow, *ca* 3.5 mm long.

Chungthang ; 1800 - 2900 m.

L. petiolata (D. Don) P.F. Hunt. & Summerh. in Kew Bull. 20 : 52. 1966. *Acianthus petiolatus* D. Don, Prodr. Fl. Nep. 29 : 1825. *Liparis nepalensis* Lindl., Bot. Reg. Sub. f. 882. 1825 ; Hook. f., I.c. 693.

Terrestrial herbs. Pseudobulbs ovate-acuminate, compressed, striate. Leaves 2, broadly ovate to suborbicular. Flowers purplish green ; lip purple.

Sikkim, *s.l.* ; 1050 - 1500 m.

Fl. : May.

L. plantaginea Lindl., Gen. & Sp. Orchid. 29. 1830 ; Hook. f., F.B.I. 5 : 702. 1890.

Epiphytes. Pseudobulbs large, narrowly ovoid-oblong, compressed, smooth, pale green. Leaves subopposite, linear-oblong or oblanceolate. Flowers green, up to 2.5 cm long.

Sikkim, *s.l.* ; 300 - 600 m.

Fl. : June - Aug.

L. platyrhachis Hook. f. in Hook., Ic. Pl. 19. t. 1890. 1899 & F.B.I. 5 : 706. 1890.

Epiphytes. Pseudobulbs tufted, oblong, compressed, *ca* 12 cm long, each bearing 3 - 5 linear-lanceolate leaves. Flowers pale ochraceous.

Chungthang ; 1200 - 1800 m.

Fl. : Sept.

L. pygmaea King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 34. t. 44. 1898.

Terrestrial herbs, up to 3.5 cm tall. Pseudobulbs up to 1.2 cm long, narrowly ovoid, loosely sheathed. Leaves 2, subopposite, ovate. Flowers yellowish, *ca* 6 mm long.

Below Dzongri ; 3900 m.

Fl. : June.

L. resupinata Ridley in J. Linn. Soc. Bot. 22 : 290. 1886. Hook. f., F.B.I. 5 : 705. 1890.

var. *resupinata*

Epiphytes. Pseudobulbs tufted, ovoid, cylindric, sheathed. Leaves 2 - 4, linear-lanceolate, sessile. Inflorescence 15 - 30 cm long, many flowered ; flowers golden brownish ; lip broadly oblong or ovate, sagittate at base.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : Oct. - Dec.

var. *ridleyi* Hook. *f.* in Hook., Ic. Pl. t. 1887. 1889 & F.B.I. 5 : 705. 1890.

Pseudobulbs, leaves and flowers larger than in var. *resupinata*. Racemes flowerless in upper part.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : Oct. - Dec.

L. stricklandiana Reichb. *f.* in Gard. Chron. 1 : 232. 1880 ; Hook. *f.*, l.c. 707. *L. griffithii* Ridley in J. Linn. Soc. 22 : 285. 1886 ; Hook. *f.*, l.c. 700.

Epiphytes. Pseudobulbs ovoid or oblong, compressed. Leaves 2, narrowly oblanceolate. Flowers yellowish green.

Pemiongtchi ; 1500 m.

Fl. : Nov.

L. viridiflora (Bl.) Lindl., Gen. & Sp. Orchid. 31. 1830 ; Hook. *f.*, F.B.I. 5 : 704. 1890. *Malaxis viridiflora* Bl., Bijdr. 392. t. 54. 1825. *L. longipes* Lindl. in Wall., Pl. As. Rar. 1 : 31. t. 35. 1829.

Epiphytes. Pseudobulbs closely arranged in a row, cylindric, stem like. Leaves 2, oblanceolate-oblong. Flowers ca 7 mm, pale yellowish green.

In tropical valleys ; 300 - 1200 m.

Fl. : Oct. - Dec.

LISTERA R. Br. (*nom. cons.*)

Listera alternifolia King & Pantl. in J. Asiatic. Soc. Beng. 65 : 126. 1896.

Terrestrial herbs, 16 - 25 cm tall. Leaves 2, alternate (to sub-opposite), obovate or elliptic. Flowers green.

Lachen valley ; 3000 m.

Fl. : July.

L. brevicaulis King & Pantl. in J. Asiat. Soc. Beng. 65 : 126. 1896.

Terrestrial herbs, 15 - 23 cm tall. Leaves 2, ovate-rotund. Flowers dull olivaceous green ; lip with brown central edge.

Lachen valley ; *ca* 2700 m.

Fl. : July.

L. dentata King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 257. 1898.

Terrestrial herbs, 20 - 30 cm tall. Leaves 2, opposite. Flowers green.

Dzongri ; *ca* 3900 m.

Fl. : July.

L. longicaulis King & Pantl. in J. Asiat. Soc. Beng. 65 : 126. 1896.

Terrestrial herbs, 15 - 35 cm tall. Leaves 2, orbicular-ovate to reniform. Flowers green, with darker veins.

Lachen valley ; *ca* 2100 m.

Fl. : July.

L. micrantha Lindl. in J. Linn. Soc. 1 : 176. 1857 ; Hook. f., F.B.I. 6 : 104. 1890.

Terrestrial herbs, 7.5 - 12.5 cm tall. Leaves 2, opposite, orbicular or cuneate-ovate. Flowers pale green ; lip 3 fid.

Lachen, Lachung valley ; 3000 - 3300 m.

Fl. : July.

L. pinetorum Lindl. in J. Linn. Soc. 1 : 175. 1857 ; Hook. f., F.B.I. 6 : 104. 1890. *L. mucronata* Panigr. & J.J. Wood in Kew Bull. 29 : 731. 1974.

Terrestrial herbs. Leaves broadly cordate or reniform. Flowers pale olivaceous green, flushed with brown ; lip pale brown with a dark band along central ridge.

Lachen valley ; 3000 - 3300 m.

Fl. : July.

L. tenuis Lindl. in J. Linn. Soc. 1 : 176. 1857 ; Hook. f., F.B.I. 6 : 104. 1890.

Terrestrial herbs, 10 - 15 cm tall. Leaves 2, elliptic, ovate. Flowers olivaceous green ; lip pale brown with dark mid rib and faint oblique lines.

Lachen valley ; 3300 - 3600 m.

Fl. : July.

LUISIA Gaud.

Luisia brachystachys Bl., Rumph. 4 : 50. 1849 ; Hook. f., F.B.I. 6 : 23. 1890. *L. indivisa* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 210. 1898.

Epiphytes without pseudobulbs, 12.5 - 17 cm tall. Flowers pale green ; lip entire.

Sikkim, s.l.

Fl. : March.

L. trichorhiza (Hook.) Bl., Mus. Bot. 1 : 63. 1849 ; Hook. f., F.B.I. 6 : 23 : 1890. *Vanda trichorhiza* Hook., Exot. Fl. 1 : t. 72. 1823.

Epiphytes, 10 - 25 cm tall. Leaves fleshy. Flowers produced on short thick tubercles ; sepals pale green with faint purple lines ; lip dark purple.

Sikkim, s.l. ; lower elevations.

Fl. : March - April.

L. volucris Lindl., Fol. Orchid. 1. 1852 ; Hook. f., F.B.I. 6 : 25. 1890.

Pendulous epiphytes, 60 - 150 cm tall. Leaves inserted on one side of stem. Flowers pale green ; lip pale green to purplish brown.

At lower elevations.

Fl. : March - April.

L. zeylanica Lindl., Fol. Orchid. Luisia 3. 1853. *L. teretifolia sensu* Hook. f., F.B.I. 6 : 22. 1890, *non* Gaud. 1829.

Epiphytes, 15 - 30 cm tall. Leaves terete, spreading. Flowers green ; lip and columns purple ; anthers pale yellow.

Sikkim, s.l. ; up to 900 m.

Fl. : May.

MALAXIS Soland ex Swartz

Malaxis acuminata D. Don, Prodr. Fl. Nepal. 29. 1825. *Microstylis wallichii* Lindl., Gen. & Sp. Orchid. 20. 1830 ; Hook. f., F.B.I. 5 : 686. 1888 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 16. 1898.

f. **acuminata**

Terrestrial herbs, ca 7.5 cm tall. Leaves 3 - 4, ovate - lanceolate. Flowers yellowish.

Teesta valley, Rangeet valley, Singhik, Rangpo ; 900 - 2100 m.

Fl. : June - Aug. - Sept.

f. biloba (Lindl.) Tuyama in Hara, Fl. E. Himalaya 443. 1966. *Microstylis biloba* Lindl., Gen. & Sp. Orchid. 20. 1830. *Microstylis wallichii* var. *biloba* (Lindl.) Hook. f., F.B.I. 5 : 606. 1888 ; King & Pantl., l.c. 8 : 16. t. 19. 1898.

Differs from *f. acuminata* in end-lobe of lip being divided up to middle into 2 oblong rounded lobules.

Sikkim, s.l. ; ca 1800 m.

Fl. : July - Sept.

M. aphylla (King & Pantl.) Tang & Wang in Act. Phytotax. Sin. 1 : 1. 1951 *Microstylis aphylla* King & Pantl., l.c. 8 : 18. 1898.

Terrestrial leafless root-parastic herbs, 17.5 - 22.5 cm tall. Flowers yellowish, flushed with dark purple.

Teesta valley ; up to ca 300 m.

Fl. : July.

M. calophylla (Reichb.f.) O. Ktze., Rev. Gen. Pl. 2 : 673. 1891. *Microstylis calophylla* Reichb.f. in Gard. Chron. 2 : 718. 1879. *M. scottii* Hook. f., F.B.I. 5 : 687. 1890. *M. wallichii* var. *brachycheila* Hook. f., F.B.I. 5 : 686. 1888.

Herbs. Leaves 2 - 5, sessile, obliquely ovate, acuminate, crenate, blackish brown with a broad pale dotted border. Scape red, lip green.

Teesta valley ; 300 - 750 m.

Fl. : July.

M. cylindrostachya (Lindl.) O. Ktze., Rev. Gen. Pl. 2 : 673. 1891. *Dienia cylindrostachya* Lindl., Gen. & Sp. Orchid. 22. 1830. *Microstylis cylindrostachya* (Lindl.) Reichb. f. in Walp. Ann. 6 : 207. 1861 ; Hook. f., F.B.I. 5 : 689. 1890.

Terrestrial herbs. Leaf solitary, oblong-elliptic. Flowers yellowish ; green.

Lachen valley ; 3000 - 3600 m.

Fl. : July.

M. josephiana (Reichb. f.) O. Ktze., Rev. Gen. Pl. 2 : 673. 1891. *Microstylis josephiana* Reichb. f. in Bot. Mag. 6325. 1877 ; Hook. f., F.B.I. 5 : 687. 1890.

Terrestrial herbs, *ca* 5 cm tall. Leaves 3 - 5, oblong or elliptic-lanceolate. Flowers ochre-yellow, shaded with brown or yellow with red blotches near the column ; 2 - 3 opening at a time.

Sikkim, *s.l.* ; 450 - 900 m.

Fl. : May.

M. khasiana (Hook. f.) O. Ktze., Rev. Gen. Pl. 673. 1891. *Microstylis khasiana* Hook. f., F.B.I. 5 : 686. 1888.

Terrestrial herbs, *ca* 7.5 cm tall. Leaves 3 - 4, ovate-elliptic, sub-oblique, shining greyish green flushed with purple. Flowers purple.

Sikkim, *s.l.* ; 900 - 1800 m.

Fl. : May - July.

M. latifolia J.E. Sm. in Rees, Cyclop. 22. 40. 3. 1819. *Dienia congesta* (Lindl.) Reichb. f. in Walp. Ann. 6 : 206. 1861 ; Hook. f., F.B.I. 6 : 689. 1890. *Microstylis latifolia* (J.E. Sm.) J.J. Sm., Fl. Buit. 6 : 248. t. 185. 1905.

Terrestrial herbs, 10 - 25 cm tall. Leaves 3 - 6, elliptic-lanceolate, oblique. Flowers red or yellow.

Sikkim, *s.l.* ; up to 750 m.

Fl. : July.

M. maximowicziana (King & Pantl.) Tang & Wang in Act. Phytotax. Sin. 1 : 1. 1951. *Microstylis maximowicziana* King & Pantl. in J. Asiat. Soc. Beng. 64 : 329. 1895.

Terrestrial herbs. Leaves 4 - 5, elliptic to elliptic-lanceolate. Flowers green.

Sikkim, s.l. ; 600 - 1200 m.

Fl. : July.

M. muscifera (Lindl.) O. Ktze., Rev. Gen. Pl. 2 : 673. 1891.
Dienia muscifera Lindl., Gen. & Sp. Orchid. 23. 1830. *Microstylis muscifera* (Lindl.) Ridl. in J. Linn. Soc. 24 : 333. 1888 ; Hook. f., F.B.I. 5 : 689. 1890.

Terrestrial herbs. Leaves 2, one broadly elliptic, the other ovate-lanceolate. Flowers *ca* 4 mm long, yellowish green.

Lachung, Thangu, Thegu, Zemu valley ; 2250 - 4050 m.

Fl. : July - Sept.

M. saprophyta (King & Pantl.) Tang & Wang in Act. Phytotax. Sin. 1 : 1. 1951. *Microstylis saprophyta* King & Pantl. in J. Asiat. Soc. Beng. 65 : 118. 1896.

Terrestrial, leafless, saprophytic herbs, 7.5 - 15 cm tall. Flowers greenish.

Chungthang, Lachen ; *ca* 1800 m.

Fl. : July.

MICROPERA Lindl.

Micropera obtusa (Lindl.) Tang & Wang in Act. Phytotax. Sin. 1 (1) : 94. 1951. *Camarotis obtusa* Lindl. in Bot. Reg. Misc. 73. 1884 ; Chowdhury & Baruah in Ind. Forester 102 : 614. f. 1. 1976. *Sarcochilus obtusus* (Lindl.) Benth. ex Hook. f., F.B.I. 6 : 36. 1890.

Epiphytic herbs. Leaves linear-oblong, coriaceous. Flowers white ; lip with a large yellow spot.

Sikkim, s.l. ; Tropical valleys.

Fl. : Aug.

MONOMERIA Lindl.

Monomeria barbata Lindl., Gen. & Sp. Orclind. 61. 1830 ; Hook f., F.B.I. 5 : 781. 1890.

Epiphytic herbs with woody *ca* 6 mm thick rhizomes. Pseudobulbs ovoid or ellipsoid, 4 - 5 cm long, 5 - 8 cm apart from each other,

shriveled during flowering. Leaf solitary, oblong. Flowers yellow with brown markings.

Sikkim s. l. ; tropical valleys.

Fl. : Feb.

MYRMECHIS (Lindl.) Bl.

Myrmechis pumila (Hook. f.) Tang & Wang in Acta Phytotax. Sin. 1 : 69. 1951. *Odontochilus pumilis* Hook. f., F.B.I. 6 : 99. 1890. *Zeuxine pumila* (Hook. f.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 291. t. 389. 1898.

Terrestrial herbs, 7.5 - 12.5 cm tall. Leaves ovate, acute. Flowers white with green tinge on the claw of lip.

Tendong ; ca 1800 m.

Fl. : July.

NEPHELAPHYLLUM Bl.

Nephelaphyllum nudum Hook. f., F.B.I. 6 : 191. 1890.

Terrestrial creeping herbs. Leaves distant, ovate, acuminate, coriaceous. Scape ca 15 cm long.

Sikkim Himalaya (*fide* Hook. f. 1890.)

NEOTTANTHE (Reichb.) Schltr.

Neottianthe secundiflora (Hook. f.) Schultr. in Fedde Repert. 16 : 291. 1919. *Habenaria secundiflora* Hook. f., F.B.I. 6 : 165. 1890, p.p.

Terrestrial herbs, 10 - 25 cm tall. Leaves 2 - 3, linear-oblong to linear. Flowers pink.

Lachung valley ; 2700 - 3300 m.

Fl. : Aug.-Sept.

NEOTTIA Guettard (*nom. cons.*)

Neottia listeroides Lindl. in Royle, Illus. Bot. Himal. 368. 1839 ; Hook. f., F.B.I. 6 : 103. 1890. *Litsera lindleyana* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 258. t. 343. 1898.

Terrestrial herbs, 20 - 35 cm tall. Flowers reddish brown.

Lachen valley ; ca 3000 m.

Fl. : July.

OBERONIA Lindl.

Oberonia acaulis Griff., Itin. Nat. 76. 1848. *O. myriantha* Lindl., Fol. Orch. Oberon 4. 1859 ; Hook. f., F.B.I. 5 : 679. 1888 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 10. 1898.

Pendulous epiphytes, acaulescent. Leaves ensiform. Inflorescence much longer than leaves. Flowers *ca* 2.5 mm long.

Penlong ; 1300 - 2300 m.

Fl. & Fr. : Oct. - Dec.

O. brachystachys Lindl., Sert. Orch. Sub. t. 8. 1838 *et* Fol. Orch. Oberon. 4. 1859 ; Hook. f., F.B.I. 5 : 677. 1888. *O. demissa* Lindl., Fol. Orch. Oberon. 4. 1859 ; Hook. f., F.B.I. 5 : 680. 1888 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 9. 1898.

Erect, acaulescent epiphytes. Leaves oblong. Inflorescence up to 5 cm long. Flowers *ca* 1.3 mm long.

In tropical valleys.

Fl. : July - Oct.

O. caulescens Wall. ex Lindl., Gen. & Sp. Orchid. 15. 1830 ; Hook. f., F.B.I. 5 : 682. 1888 ; King & Pantl., l.c. 8 : 133 B. 1898 (*quoad nom.*). *O. longilabris* King & Pantl. in J. Asiat. Soc. Beng. 64 : 330. 1895 *et* in Ann. Roy. Bot. Gard. Calc. 8 : 13, pl. 16 A. 1898.

Erect epiphytes, up to 3.5 cm long. Leaves linear, ensiform. Inflorescence up to 9 cm long. Flowers *ca* 3 mm long.

Song, Chongloo, 1200 - 2500 m.

Fl. : July - Aug.

O. denticulata Wight, Icon. 5 (1) : 3, t. 1625. 1851. *O. iridifolia* Lindl., Gen. & Sp. Orchid. 15. 1830, *p.p.* (*nom. superfl. et illeg.*) ; *et*. Fol. Orch. Oberon. 1. 1859 ; Hook. f., F.B.I. 5 : 675. 1888 (incl. var. *denticulata* Hook. f., *angustifolia* Hook. f. and var. *brevifolia* Hook. f.) ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : t. 8. 1898. *O. gammiei* King & Pantl. in J. Asiat. Soc. Beng. II. 66 : 579. 1897.

Acaulescent epiphytes. Leaves oblong, ensiform. Spikes up to 30 cm long. Flowers pale brownish to yellow, in verticiles.

Sikkim s.l. ; *ca* 1400 m.

Fl. : Sept. - Nov.

O. emarginata King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 6. pl. 2A. 1898. *O. micrantha* King & Pantl., l.c. 8 : 6, pl. 5. 1898, *non* A. Rich. 1834.

Epiphytes, acaulescent. Leaves linear. Flowers *ca* 0.5 - 1 m long, in whorls.

Namgah, Tendong ; *ca* 2000 m.

Fl. : July - Sept.

O. ensiformis (Smith) Lindl., Fol. Orch. Oberon. 4. 1859 ; Hook. f., F.B.I. 5 : 679. 1888. *Malaxis ensiformis* Smith, Rees. Cyclop. 22, no. 14. 1812.

Pendulous, acaulescent, epiphytic herbs. Leaves linear, ensiform. Flowers *ca* 3.5 mm long, brownish yellow.

Teesta valley ; *ca* 450 m.

Fl. : Sept. - Jan.

O. falcata King & Pantl. in J. As. Soc. Beng. II, 64 : 329. 1895 & in Ann. Roy. Bot. Gard. Calc. 8 : 12. pl. 14. 1898.

Pendulous epiphytes, caulescent, up to 15 cm long. Leaves linear. Flowers pedicelled, *ca* 2.2 mm long.

Sikkim, *s.l.* ; 3000 m.

Fl. : June - Aug.

O. falconeri Hook. f., Icon. t. 1780. 1888 ; Hook. f., F.B.I. 5 : 678. 1888.

Erect epiphytes, acaulescent. Leaves oblong-lanceolate. Flowers pedicelled, *ca* 2.5 mm long.

Sikkim, *s.l.* ; 500 - 2000 m.

Fl. : Aug. - Sept.

O. jenkinsiana Lindl., Fol. Orch. 20. 1859 ; Hook. f., F.B.I. 5 : 676. 1888 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 7. pl. 7. 1898.

Erect or pendulous epiphytes, acaulescent. Leaves linear. Flowers subsessile, *ca* 1.7 mm long.

Teesta valley ; up to 1200 m.

Fl. ; Sept. - Nov.

C. lobulata King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 4. 1898.

Erect or pendulous epiphytic, acaulescent herbs. Leaves oblong-elliptic. Flowers pedicelled, green.

Teesta valley ; *ca* 300 m.

Fl. : Oct.

O. obcordata Lindl., Fol. Orch. Oberon. 7. 1859 ; Hook. f., F.B.I. 5 : 684. 1888 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 9, pl. 11. 1898. *Oberonia treutleri* Hook. f., Icon. t. 1786. 1888 *et* F.B.I. 5 : 683. 1888.

Erect or pendulous epiphytes, acaulescent. Leaves linear or lanceolate. Flowers pedicelled, *ca* 1.5 mm long.

Sikkim, *s.l.* ; *ca* 2000 m.

O. pachyrachis Reichb. f. *ex* Hook. f., F.B.I. 5 : 681. 1888 ; King & Pantl., Ann. Roy. Bot. Gard. Calc. 8 : 4, pl. 3. 1898.

Pendulous epiphytes, acaulescent. Leaves oblong-ensiform. Flowers sessile, *ca* 1.2 mm long, greenish yellow to orange-brown.

Teesta valley ; 1000 - 1800 m.

Fl. : Sept. - March.

O. prainiana King & Pantl. in J. As. Soc. Beng. II. 64 : 331. 1895 & Ann. Roy. Bot. Gard. Calc. 8 : 3, pl. 1. 1898.

Erect or pendulous epiphytes, acaulescent. Leaves oblong-ovate. Flowers pedicelled, *ca* 1.2 mm long.

Dumsong ; up to 1500 m.

Fl. : April - June.

O. pyrulifera Lindl., Fol. Orch. Oberon. 3. 1859 ; Hook. f., F.B.I. 5 : 678. 1888 ; King & Pantl., Ann. Roy. Bot. Gard. Calc. 8 : 14. pl. 16B. 1898.

Epiphytes, erect, acaulescent. Leaves linear. Flowers pedicelled, 2 mm long, pale or orange yellow.

Tropical valleys ; up to 2000 m.

Fl. : Sept. - Oct.

O. recurva Lindl. in Bot. Reg. 25, Misc. no. 8. 1839 & Fol. Orch. Oberon. 5. 1859 ; Hook. f., F.B.I. 5 : 680. 1888. *O. parvula* King & Pantl., J. As. Soc. Beng. 64 : 330. 1895. *O. croftiana* King & Pantl., Ann. Roy. Bot. Gard. Calc. 8 : 7. 1898.

Erect or pendulous epiphytes, acaulescent. Leaves linear-ensiform. Flowers pedicelled, 1 - 2 mm long, brown or orange, in verticiles.

Tropical valley ; 900 - 2000 m.

Fl. : Oct. - Mar.

OREORCHIS Lindl.

Oreorchis foliosa (Lindl.) Lindl. in J. Proc. Linn. Soc. Bot. 3 : 27. 1858 ; Hook. f., F.B.I. 5 : 709. 1890. *Corallorrhiza foliosa* Lindl., Gen. & Sp. Orchid. 535. 1840.

Terrestrial herbs with corm like pseudobulbs. Leaf solitary, terminal, oblong-lanceolate, acuminate. Flowers reddish.

Lachen valley ; ca 3000 m.

Fl. : July.

O. micrantha Lindl. in J. Linn. Soc. 3 : 27. 1858 ; Hook. f., F.B.I. 5 : 709. 1890.

Terrestrial herbs with corm like pseudobulbs. Leaves usually 2, linear. Flowers pale yellow ; lip red spotted.

Lachen valley ; ca 3000 m.

Fl. : July.

ORNITHOCHILUS (Lindl.) Wall.

Ornithochilus difformis (Wall. ex Lindl.) Schltr. in Fedde Repert. 4 : 277. 1919. *Aerides difformis* Wall. ex Lindl., Gen. & Sp. Orchid. 242. 1833. *O. fuscus* Wall. ex Hook. f., F.B.I. 6 : 76. 1890.

Epiphytes without pseudobulbs. Leaves broadly oblong, flat. Flowers greenish yellow, with brown bands on sepals and petals.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : July - Aug.

OTOCHLUS L.

Otochilus albus Lindl., Gen. & Sp. Orchid. 35. 1830 ; Hook. f., F.B.I. 5 : 843. 1890.

Epiphytes. Stems jointed, branched. Leaves in pairs arising from apex of the terminal node, elliptic-oblong. Flowers white.

Sikkim, s.l. ; 900 - 1500 m.

Fl. : June.

O. fuscus Lindl., Gen. & Sp. Orchid. 35. 1830 ; Hook. f., F.B.I. 5 : 844. 1890.

Pendulous epiphytes with cylindric or fusiform internodes. Leaves linear-oblong. Flowers white with brown saccate base of the lip.

Sikkim, s.l. ; 900 - 1800 m.

Fl. : Dec.

O. porrectus Lindl., Gen. & Sp. Orchid. 36. 1830 ; Hook. f., F.B.I. 5 : 844. 1890, *p.p.*

Epiphytic herbs with jointed stem. Leaves narrowly elliptic. Flowers white with a brown patch on sac of the lip and column.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. : Oct. - Jan.

PACHYSTOMA Bl.

Pachystoma senile (Lindl.) Reichb. f. in Bonplandia 3 : 250. 1858 ; Hook. f., F.B.I. 5 : 812. 1890. *Apatura senilis* Lindl., Gen. & Sp. Orchid. 130. 1831. *A. lindleyana* Wt., Ic. t. 1662. 1851.

Terrestrial, rhizomatous herbs. Leaf appearing after flowering, solitary, linear, *ca* 30 cm long. Flowers greenish white, flushed with pink.

Sikkim, s.l.

Fl. : March.

PANISEA (Lindl.) Lindl.

Panisea demissa (D. Don) Pfitz. in Engl. Pflanzenr. 32 : 141. t. 49. 1907. *Dendrobium demissum* D. Don, Prodr. Fl. Nepal. 34. 1825. *Coelogyne parviflora* Lindl., Gen. & Sp. Orchid. 44. 1830. *Panisea parviflora* (Lindl.) Lindl., Fol. Orchid. 1. 1824 ; Hook. f., F.B.I. 5 : 783. 1890.

Epiphytes with narrow ovoid pseudobulbs. Leaves narrowly elliptic-lanceolate, acute. Flowers white with brown streaks.

Sikkim, s.l. ; 1800 m.

Fl. : Oct.

PANTLINGIA Prain

Pantlingia paradoxa Prain in J. Asiat. Soc. Beng. 65 : 107. 1896.

Terrestrial saprophytic, aphyllous, rhizomatous herbs, up to 10 cm tall. Stem with a ovate bract in the middle. Flowers pale yellow with a purple blotch on the lip.

Chungthang ; ca 1800 m.

Fl. : Sept.

PAPHIOPEDILUM Pfitz.

Paphiopedilum venustum (Sims) Pritz. in Engl., Bot. Jahrb. 19 : 41. 1849. *Cypripedium venustum* Sims in Curtis Bot. Mag. 47. t. 2129. 1820 ; Hook. f., F.B.I. 6 : 173. 1890.

Terrestrial herbs. Leaves 5 - 7, elliptic-oblong, somewhat loriform, acute and minutely bifid at apex, dark green, marbled with pale green above and dull purple beneath. Flowers 5 - 6.25 cm in diam., dorsal sepal white with dark green vertical stripes ; petals green and purplish with a few black spots ; lips yellowish green flushed with pink ; lateral edges yellow.

Tropical valleys ; up to 1200 m.

Fl. : Sept. - Oct.

PAPILIONANTHE Schltr.

Papilionanthe teres (Roxb.) Schltr. in Orchis 9 : 78. 1915. *Dendrobium teres* Roxb., Fl. Ind. 3 : 485. 1832. *Vanda teres* (Roxb.) Lindl., Gen. & Sp. Orchid. 217. 1833 ; Hook. f., F.B.I. 6 : 49. 1890.

Epiphytic herbs. Leaves fleshy, terete. Flowers pink (rarely white) ; spur yellow or ferruginous, marked with purplish brown lines.

Gangtok, Melli, Phodong, Rangpo, Singtam. up to 1700 m.

Fl. : July - Aug. ; *Fr.* : Sept. - Oct.

P. uniflora (Lindl.) Garay in Bot. Mus. Leafl. Harvard Univ. 23 : 370. 1974. *Aerides uniflora* (Lindl.) Summerh. in Kew Bull. 1955 ; 558. 1956. *Mesoclastis uniflora* Lindl., Gen. & Sp. Orchid. 45. 1830. *Aerides longicornu* Hook. f., F.B.I. 6 : 44. 1890.

Epiphytes with leafy stem. Leaves terete. Flowers white with a flush of pink on the spur.

Sikkim, s.l. (*cf. Hara et al. 1978*) ; *ca* 1650 m.

Fl. : Sept.

P. vandarum (Reichb. f.) Garay in Bot. Mus. Leafl. Harvard Univ. 23 : 370. 1974. *Aerides vandarum* Reichb. f. in Gard. Chron. 1867 ; 997. 1867 ; Hook. f., F.B.I. 6 : 44. 1890.

Epiphytes. Leaves 15 - 25 cm long. Flowers white.

Sikkim (*cf. Hara et al. 1978*).

PERISTYLVUS Bl.

Peristylus affinis (D. Don) Seid. in Dansk. Bot. Ark. 31 (3) : 48. 1977. *Habenaria affinis* D. Don, Prodr. Fl. Nepal. 25. 1825 ; Hook. f., l.c. 6 : 166. 1890. *Habenaria goodyeroides* var. *affinis* (D. Don) Lindl., Gen. & Sp. Orchid. 299. 1835.

Terrestrial herbs. Leaves ovate-elliptic. Flowers greenish white or cream coloured.

West Sikkim ; *ca* 600 m.

Fl. : June.

P. aristatus Lindl., Gen. & Sp. Orchid. 300. 1835. *Habenaria aristata* (Lindl.) Trin. in J. Ceylon. Branch. Roy. Asiat. Soc. 9 : 91. 1885 ; Hook f., F.B.I. 6 : 158. 1890.

Terrestrial herbs, 30 - 50 cm tall. Leaves 3 - 5, unequal, elliptic-lanceolate. Flowers green.

Chungthang, Tumlong, Pashating ; 1800 - 2100 m.

Fl. : Aug.

P. constrictus (Lindl.) Lindl., Gen. & Sp. Orchid. 300. 1835. *Herminium constrictum* Lindl. in Edw. Bot. Reg. 18, Sub. t. 1499. 1832. *Habenaria constricta* (Lindl.) Hook. f., F.B.I. 6 : 161. 1890.

Terrestrial herbs, 50 - 75 cm tall. Leaves 5 - 6, in a whorl near stem tip, ovate-elliptic. Flowers brown ; lip white.

Teesta valley ; *ca* 450 m.

Fl. : June - July.

P. fallax Lindl., Gen. & Sp. Orchid. 298. 1835. *Herminium fallax* (Lindl.) Hook. f., F.B.I. 6 : 129. 1890. *Habenaria fallax* (Lindl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 325. t. 428. 1898.

Terrestrial herbs, 10 - 30 cm tall. Leaf solitary, narrowly elliptic, acute. Flowers green.

Lachen & Lachung valleys, Gnathang ; 3000 - 3600 m.

Fl. : July.

P. goodyeroides (D. Don) Lindl., Gen. & Sp. Orchid. 299. 1835. *Habenaria goodyeroides* D. Don, Prodr. Fl. Nep. 25. 1825 ; Hook. f., F.B.I. 6 : 161. 1890.

Terrestrial herbs, 30 - 60 cm tall. Leaves close below the apex but not whorled, ovate-elliptic, acute. Flowers white or cream coloured.

Sikkim Himalaya ; ca 900 m.

Fl. : May - July.

P. parishii Reichb.f. in Trans. Linn. Soc. 30 : 139. 1874. *Habenaria parishii* (Reichb. f.) Hook. f., F.B.I. 6 : 161. 1890.

Terrestrial herbs, 30 - 45 cm tall. Leaves 3 - 5, ovate or oblong. Flowers brownish green.

Sikkim, s.l. ; ca 1050 m.

Fl. : Aug.

P. prainii (Hook. f.) Kranz., Orchid. Gen. Sp. 514. 1898. *Habenaria prainii* Hook. f., F.B.I. 6 : 159. 1890.

Terrestrial herbs, 30 - 45 cm tall. Leaves 3 - 5, oblong to ovate-oblong. Flowers brownish green.

Sikkim, s.l. ; 1500 m.

P. pseudophrys (King & Pantl.) Mitra, Fl. Pl. E. India 1 : 274. 1958. *Habenaria pseudophrys* King & Pantl. in J. Asiat. Soc. Beng. 65 : 133. 1896.

Terrestrial herbs, 45 - 70 cm tall. Leaves 4, elliptic-oblong.

Chungthang ; 1650 m.

Fl. : July.

P. tentaculatus (Lindl.) J.J. Smith, Orch. Java. 35. 1905. *Glossula tentaculata* Lindl., Bot. Reg. t. 862. 1824. *Habenaria tentaculata* (Lindl.) Reichb. f., Otia. Bot. Hamb. 34. 1878.

Terrestrial herbs, 15 - 45 cm. Leaves 2 - 3, unequal, oblong-elliptic or linear-oblong. Flowers green.

Gorubathan (South Sikkim Dist.) ; 450 m.

Fl. : Aug.

PHAIUS Lour.

Phaius flavus (Bl.) Lindl., Gen. & Sp. Orchid. 128. 1831. *Limodorum flavum* Bl., Bijdr. 8 : 375. 1825. *Phaius maculatus* Lindl., Gen. & Sp. Orchid. 127. 1831 ; Hook. f., F.B.I. 5 : 817. 1890.

Herbs, up to 60 cm tall. Pseudobulbs ovoid cylindric. Leaves 4 - 6, elliptic to lanceolate, plicate. Flowers yellow with darker stripes ; terminal lobe of lip brown.

Tropical valleys.

Fl. : April.

P. mishmensis Reichb. f. in Bonplandia 5 : 43. 1857 ; Hook. f., F.B.I. 5 : 817. 1890.

Herbs, up to 1.2 m tall. Pseudobulbs naked below. Leaves elliptic-lanceolate, acute. Flowers rose coloured ; spur yellow.

Sikkim, s.l. ; 1700 - 1800 m.

Fl. : Sept. - Oct.

P. tancarvilleae (Banks ex L. Herit.) Bl., Mus. Bot. Lugd. Bat. 2 : 177. 1856 ("*tankervillii*"). *Limodorum tancarvilleae* Bank ex L'. Herit., Sert. Angl. 28. 1789. *Phaius wallachii* Lindl. in Wall. Pl. Asiat. Rar. 2 : 46, t. 158. 1831 ; Hook. f., F. B.I. 5 : 816. 1890.

var. **tancarvilleae**

Herbs 60 - 105 cm tall. Pseudobulbs conical, sheathed. Leaves elliptic-lanceolate, acute. Sepals and petals brown inside, white outside ; lip pale purple passing into white on edges and yellow on tube.

Tropical valleys.

Fl. : March - April.

var. **pulchra** (King & Pantl.) Karthik. in Karthik. et al., Fl. Ind. Enum. Monocot. 170. 1989. *Phaius blumei* var. *pulchra* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 108. 1898.

Differs from var. *tancarvilleae* in spur being less than 6 mm and the mouth of lip contracted and trilobulate.

Tropical valleys.

Fl. : March - April.

P. woodfordii (Hook.) Merr. in J. Arn. Arbor. 24 : 211. 1948.
Bletia woodfordii Hook., Bot. Mag. t. 2719. *Phalus maculatus* sensu
 Hook. f., F.B.I. 5 : 817. 1890, *p.p.non* Lindl. 1831.

Herbs with ovoid-cylindric pseudobulbs. Leaves 4 - 6, elliptic-lanceolate. Flowers yellow.

Sikkim, s.l.

Fl. : April.

PHALAEENOPSIS Bl.

Phalaenopsis mannii Reichb. f. in Gard. Chron. 902. 1871 ; Hook. f., F.B.I. 6 : 30. 1890.

Epiphytes without pseudobulbs. Leaves oblong lanceolate, obtuse. Flowers yellow.

Teesta valley ; up to 1200 m.

Fl. : May.

P. lobbii (Reichb. f.) Sweet, Genus Phalaenopsis (The Orchid Digest) 53. 1980. *P. parishii* var. *lobbii* Reichb. f. in Saunders Refug. Bot. 2. sub. t. 85. 1870. *P. listeri* Berkeley in Gard. Chron. ser. 3. 1 : 280. 1887.

Epiphytic herbs without pseudobulbs. Leaves broadly elliptic, obliquely notched at apex. Flowers white with yellow and brown spots on column.

Teesta valley ; 360 - 1200 m.

Fl. : March - April.

PHOLIDOTA Lindl. ex Hook.f.

Pholidota articulata Lindl., Gen. & Sp. Orchid. 38. 1830 ; Hook. f., F.B.I. 5 : 844. 1890. var. *griffithii* King & Pantl., I.c. 8 : 147. 1898. *Pholidota griffithii* Hook. f., Ic. Pl. t. 1881. 1889 & F.B.I. 5. 845. 890.

Epiphytes. Stems jointed. Leaves narrowly oblong. Flowers white, tinged with brownish pink ; floral bracts pale brown.

Sikkim, s.l. ; 1200 m.

Fl. : June - Aug.

P. imbricata (Roxb.) Lindl. in Hook. f., Fl. Exot. 2. t. 138. 1825 ; Hook. f., F.B.I. 5 : 845. 1890. *Cymbidium imbricatum* Roxb., Fl. Ind. 3 : 460. 1832.

Epiphytes with jointed stem. Pseudobulbs caespitose, cylindric. Flowers pale brown.

Gangtok ; 1000 - 1500 m.

Fl. : May - Aug.

P. protracta Hook. f. in Hook., Ic. Pl. 19 : t. 1877. 1889 & F.B.I. 5 : 845. 1890.

Epiphytes. Stem elongate, clothed with long scarious sheaths. Flowers green ; lip with 4 small bright patches of dull yellow on its anterior lobe.

Dentam - Pamianchi ; 1500 - 2250 m.

Fl. : Oct. - Dec.

P. recurva Lindl., Gen. & Sp. Orchid. 37. 1830 ; Hook. f., F.B.I. 5 : 844. 1890.

Epiphytes, stem jointed ; internodes yellowish, angled. Leaves elliptic-oblong, acute. Flowers pale brown.

Sikkim, s.l. ; 600 - 1200 m.

Fl. : Aug. - Sept.

P. undulata Lindl. in Edw. Bot. Reg. 14 : t. 1213. 1829. *P. rubra* Lindl., Gen. & Sp. Orchid. 37. 1830. Hook. f., F.B.I. 5 : 847. 1890.

Epiphytes. Leaves elliptic-lanceolate. Racemes decurved, lax.

Sikkim, s.l. ; 1200 - 1800 m.

PHREATIA Lindl.

Phreatia elegans Lindl. , Gen. & Sp. Orchid. 63. 1830 ; Hook. f., F.B.I. 5 : 810. 1890.

Plants caespitose. Pseudobulbs tufted, ovoid, compressed. Leaves narrowly elliptic, subacute, or acute. Flowers minute, white.

Chungthang, ca 1800 m.

Fl. : Aug.

PLATANTHERA L.C. Rich.

Platanthera backertana (King & Pantl.) Kranzl., Orchid. Gen. Sp. 1 : 632. 1898. *Habenaria backeriana* King & Pantl. in J. Asiat. Soc. Beng. 65 : 132. 1896 ; Bruhl, l.c. 182.

Terrestrial herbs, 22 - 45 cm tall. Leaves several, scattered, sessile ; lower one oblong, upper 3 - 4 linear-lanceolate. Flowers green.

Lachen valley, Singalela range ; 2700 - 3600 m.

Fl. : July.

P. biermanniana (King & Pantl.) Kranzl., Orchid. Gen. Sp. 1 : 636. 1898. *Habenaria biermanniana* King & Pantl., l.c. 64 : 343. 1895.

Terrestrial herbs. Leaves 4 - 5, caudine, amplexicaule, lanceolate.

Singa - le - la ; *ca* 3000 m.

Fl. : July.

P. clavigera Lindl., Gen. & Sp. Orchid. 289. 1835. *Habenaria densa* Wall. ex Lindl., l.c. 326. 1835; Hook. f., F.B.I. 6 : 153. 1890.

Terrestrial herbs. Leaves 3 - 5, narrowly elliptic. Flowers green.

Lachung valley, Singa - le - la range ; 2400 - 3600 m.

Fl. : July - Aug.

P. dyeriana King & Pantl. in J. Asiat. Soc. Beng. 65 : 133. 1896. *Habenaria dyeriana* King & Pantl. in J. Asiat. Soc. Beng. 65 : 133. 1896.

Terrestrial herbs, less than 30 cm tall. Leaves 3 - 4, sessile, scattered along the stem. Flowers green.

Lachen valley ; *ca* 3300 m.

Fl. : Aug.

P. excelliana Soo in Ann. Hist. Nat. Mus. Natl. Hung. 26 : 259. 1929. *Habenaria oligantha* Hook. f., F.B.I. 6 : 154. 1890.

Terrestrial herbs, 7.5 - 30 cm tall. Leaves usually solitary, oblong-ob lanceolate. Flowers green.

Lachen valley, Gnathang, Dzongri ; 3000 - 3600 m.

Fl. : July - Sept.

P. juncea (King & Pantl.) Kranzl., Orchid. Gen. Sp. 1 : 942. 1901. *Habenaria juncea* King & Pantl. in J. Asiat. Soc. Beng. 65 : 132. 1896.

Terrestrial herbs, *ca* 30 cm tall. Leaf solitary, narrowly oblong.

Lachen, Lachung, Gnathang, Singa - le - la range ; 3300 - 3600 m.

Fl. : Aug.

P. latilabris Lindl., Gen. & Sp. Orchid. 289. 1835. *Habenaria latilabris* (Lindl.) Hook. f., F.B.I. 6 : 153. 1890.

Terrestrial herbs, 20 - 45 cm tall. Leaves 5, elliptic-oblong. Flowers green ; lip greenish yellow.

Lachung, Lachen valley ; 2400 - 3000 m.

Fl. : July - Sept.

P. leptocaulon (Hook. f.) Soo in Ann. Nat. Hist. Mus. Hung. 26 : 360. 1929. *Habenaria leptocaulon* Hook. f., F.B.I. 6 : 154. 1890.

Terrestrial herbs, 20 - 40 cm tall. Leaves 2, unequal, narrowly elliptic. Flowers green.

Lachung valley ; 2700 - 3300 m.

Fl. : July - Sept.

P. sikkimensis (Hook. f.) Kranzl., Orchid. Gen. Sp. 1 : 621. 1899. *Habenaria sikkimensis* Hook. f., F.B.I. 6 : 155. 1890.

Terrestrial herbs, 25 - 30 cm tall. Leaves narrowly elliptic - oblong. Flowers green.

Lachung valley, Lingtoo - Zelap la, Singa - le - la ; *ca* 3600 m.

Fl. : Aug. - Sept.

P. stenantha (Hook. f.) Soo in Ann. Hist. Nat. Mus. Natl. Hung. 26 : 363. 1929. *H. stenantha* Hook. f., F.B.I. 6 : 153. 1890.

Terrestrial herbs, 30 - 60 cm tall. Leaves 3 - 4, oblong-lanceolate. Flowers green ; lip yellowish.

Sikkim, *s.l.* ; 2400 - 3600 m.

Fl. : July - Sept.

PLEIONE D. Don

Pleione hookeriana (Lindl.) Williams, Orchid Grow. Man. ed. 6 : 548. 1885. *Coelogyné hookeriana* Lindl., Fol. Orchid. Coelogyné 14 : 1854 ; Hook. f., F.B.I. 5 : 842. 1890.

Pseudobulbs ovoid or ellipsoidal. Flowers white flushed with violet ; lip with 5 - 7 thin parallel longitudinal ridges and reddish mottlings.

Dombong valley ; up to 3600 m.

Fl. : June.

P. humilis (J.E. Smith) D. Don, Prodr. Fl. Nep. 37. 1825. *Epidendrum humile* J.E. Smith, Exot. Bot. 2 : 75. t. 98. 1806. *Coelogyne humilis* (J.E. Smith) Lindl., Coll. Bot. Sub. t. 37. 1825 ; Hook. f., F.B.I. 5 : 840 . 1890.

Pseudobulbs ovoid-oblong. Flowers white ; central part of lip yellow with brown spots, margins purple speckled.

Tendong Peak ; 2100 - 2550 m.

Fl. : May - June.

P. maculata (Lindl.) Lindl. in Paxt., Fl. Gard. 2 : 5. t. 39. f. 1. 1851. *Coelogyne maculata* Lindl., Gen. & Sp. Orchid. 43. 1830 ; Hook. f., F.B.I. 5 : 841. 1890.

Pseudobulbs enclosed in a dense network of fibres. Leaves broadly oblanceolate, appearing after flowering. Flowers white, some times with pink lines ; lip with purple yellow markings.

Sikkim, s.l. ; 480 - 900 m.

Fl. : May.

P. praecox (J.E. Smith) D. Don, Prodr. Fl. Nepal. 37. 1825. *Epidendrum praecox* J.E. Smith, Exot. Bot. 2 : 73. 1806. *Coelogyne praecox* (J.E. Smith) Lindl., Coll. Bot. Sub. t. 37. 1825 ; Hook. f., F.B.I. 5 : 840. 1890.

Pseudobulbs, 2.5 - 4 cm in diam., sparsely sheathed by loose fibres. Flowers rose coloured, rarely white.

Dentam - Pamianchi ; 1800 - 2400 m.

Fl. : Oct. - Nov. (-May).

PODOCHILUS Bl.

Podochilus cultratus Lindl., Gen. & Sp. Orchid. 234. 1833 ; Hook. f., F.B.I. 6 : 80. 1890.

Plants up to 15 cm long, clothed with equitant flattend, falcately lanceolate, acute leaves. Racemes slender, terminal, 2 - 4 - flowered. Flowers white, slightly flushed with pink.

Singtam - Rangpo ; 600 - 900 m.

Fl. : Sept. - Oct.

P. khasianus Hook. f., F.B.I. 6 : 81. 1890.

Plants up to 15 cm tall. Leaves linear-lanceolate, acute, coriaceous. Racemes terminal, erect. Flowers 3 - 4, distichous, white, flushed with yellow.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. : Sept.

POMATOCALPA Breda

Pomatocalpa armigerum (King & Pantl.) Tang & Wang in Acta Phytotax. Sin. 1 : 98. 1951. *Cleisostoma armigera* King & Pantl. in J. Asiat. Soc. Beng. 65 : 123. 1896.

Epiphytes without pseudobulbs. Leaves fleshy, semiterete, channeled in front. Flowers pale straw coloured, flushed with pink.

Sikkim, s.l.

Fl. : Sept.

P. wendlandorum (Reichb.f.) J. J. Sm. in Nat. Tijdschr. Med. Ind. 72 : 108. 1912. *Cleisostoma wendlandorum* Reichb. f. in Otto. & Dietr., Allg. Gart. 219. 1956 ; Hook. f., F.B.I. 6 : 74. 1890.

Epiphytes with very short stem. Leaves crowded, pendulous, obtuse and obliquely 2-lobed at apex. Flowers yellow, flushed with pink.

Sikkim, s.l. ; 450 m.

Fl. : May - June.

PONERORCHIS Reichb.f.

Ponerorchis chusua (D. Don) Soo in Acta Bot. Acad. Sci. Hung. 12 : 352. 1966. *Orchis chusua* D. Don, Prodr. Fl. Nepal. 23. 1825 ; Hook. f., F.B.I. 6 : 127. 1890.

var. **chusua**

Terrestrial herbs, 10 - 25 cm tall. Leaves 1 - 2 ; narrowly oblong. Flowers dark purple.

Singa-le-la range, Tallum Samdong in Lachen Valley ; 3000 - 3900 m.

Fl. : July - Aug.

var. **nana** (King & Pantl.) R.C. Srivastava *Ponerorchis nana* (King & Pantl.) Soo in *Acta Sci. Hung.* 12 : 353. 1966. *Orchis chusua* var. *nana* King & Pantl. in *Ann. Roy. Bot. Gard. Calc.* 8 : 303. t. 402 A. 1898. *Chusua roborowskyi* (Maxim.) P.F. Hunt. var. *nana* (King & Pantl.) P.F. Hunt. in *Kew Bull.* 26 : 176. 1971.

Terrestrial herbs, 5 - 7.5 cm tall. Leaf solitary. Flowers dark purple ; middle lobe of lip shallowly lobed ; spur subcompressed.

Singa-le-la range, Lachen valley ; 3000 - 3900 m.

Fl. & Fr. : July - Aug.

PORPAX Lindl.

Porpax fibuliformis (King & Pantl.) King & Pantl. in *Ann. Roy. Bot. Gard. Calc.* 8 : 114. t. 159. 1898. *Eria fibuliformis* King & Pantl. in *J. Asiat. Soc. Beng.* 64 : 336. 1895.

Small epiphytic herbs with crowded or overlapping pseudobulbs. Leaves in pairs, oblanceolate, sessile. Flowers dull red.

In tropical valleys.

Fl. : Oct.

P. meirax (Par. & Reichb.f.) King & Pantl. in *Ann. Roy. Bot. Gard. Calc.* 8 : 114. t. 158. 1898. *Cryptochilus meirax* Par. & Reichb.f. in *Trans. Linn. Soc.* 30 : 148. 1874. *Eria meirax* Hook.f., F.B.I. 5 : 786. 1890. *p.p.*, non N.E. Br. 1880.

Small epiphytes with clustered, depressed-turbinate, subdiscoid pseudobulbs. Leaves 2, elliptic-oblong, shortly petiolate. Flowers dull brown.

Sikkim, *s.l.* ; 450 - 750 m.

Fl. : Oct.

PTEROCERAS Hasselt ex Hassk.

Pteroceras suaveolens (Roxb.) Holttum in *Kew Bull.* 14 : 271. 1960. *Aerides suaveolens* Roxb., *Fl. Ind.* 3 : 473. 1832. *Sarcochilus suaveolens* (Roxb.) Hook.f., F.B.I. 6 : 33. 1890 ; Smith & Cave in *Rec. BSI* 5 : 243. 1911. *Ornitharium striatum* Lindl. in *Pat.*, *Fl. Gard.* 1: 188 f. 117. 1853. *Thrixspermum teres* Reichb.f., *Xen. Orch.* II. 121. 123 (excl. syns).

Epiphytes without pseudobulbs. Leaves distichous or absent during flowering time. Flowers yellow, speckled with red, fragrant.

Teesta valley ; ca 1200 m.

RHYNCHOSTYLIS Bl.

Rhynchostylis retusa (L.) Bl., Bijdr. 286. t. 49. 1825 ; Hook. f., F.B.I. 6 : 32. 1890. *Epidendrum retusum* L., Sp. Pl. 953. 1753. *Saccolabium rheedi* Wt., Ic. t. 1745-46. 1851. DROUPADI MALA (Hindi).

Epiphytes. Leaves curved, keeled. Flowers pink, variously marked with purple or violet.

Gangtok-Saramsa, Rumtek ; ascending to 1200 m.

Fl. : June - July.

RISLEYA King & Pantl.

Risleya atropurpurea King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 246. 1898.

Terrestrial, aphyllous herbs. Flowers minute, deep blackish purple.

Dzongri ; ca 3900 m.

Fl. : June - July.

RITALA King & Pantl.

Ritala himalaica (Hook. f.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 157. t. 214. 1898. *Ceratostylis himalaica* Hook. f., F.B.I. 5 : 826. 1890.

Pendulous epiphytes. Leaf solitary, linear-oblong, obliquely notched and acute at apices. Flowers pale yellow.

Tendong ; ca 1500 m.

Fl. : May.

ROBIQUETIA Gaud.

Robiquetia spathulata (Bl.) J.J. Smith in Nat. Tijdschr. Ned. Ind. 72 : 115. 1912. *Cleisostoma spathulatum* Bl., Bijdr. 364 : 1825. *C. spicatum* Lindl. in Bot. Reg. t. 32. 1847.

Pendulous epiphytes. Leaves oblong, coriaceous. Flowers yellow, mottled and banded with brownish purple ; column white.

Teesta valley ; 300 m.

Fl. : June.

R. succisa (Lindl.) Seid. & Garay in Bot. Tidsskr. 67 : 119. 1972.
Sarcanthus succisus Lindl. in Bot. Reg. t. 1014. *Saccolabium buccosum* Reichb. f. in Gard. Chron. 938. 1871 ; Hook. f., F.B.I. 6 : 58. 1890.

Pendulous, epiphytic herbs. Leaves narrowly oblong, obliquely lobed at apex. Flowers lax.

Sikkim, s.l. ; 600 m.

Fl. : May - June.

SACCOLABIOPSIS J.J. Sm.

Saccolabiopsis pusilla (Lindl.) Seid & Garay in Bot. Tikdskr. 67 : 118. t. 33. 1972 ; Hook. f., F.B.I. 6 : 56. 1890. *Saccolabium pusillum* Lindl. in J. Linn. Soc. 3 : 36. 1858. *Saccolabium pumilio* Reichb. f. in Walp. Ann. 6 : 886. 1861 ; Hook. f., F.B.I. 6 : 56. 1890.

Epiphytes. Leaves linear-oblanceolate. Flowers cream coloured.

Sikkim, s.l. ; ca 450 m.

SATYRIUM Swartz (*nom. cons.*)

Satyrium ciliatum Lindl., Sp. Orchid. 341. 1838. *S. nepalense* var. *ciliata* (Lindl.) Hook. f., F.B.I. 6 : 168. 1890 ; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 338 - 339. 1898. KHIRU (Tibetan).

Terrestrial herbs, up to 28 cm tall. Leaves oblong-lanceolate. Flowers rose coloured.

Lachung, 2400 - 3000 m.

Fl. : Sept. - Oct.

Uses : Tubers are eaten by Lepcha & Bhutia people after roasting (cf. Raju & Srivastava 1990).

S. nepalense D. Don, Prodr. Fl. Nep. 26. 1825 ; Hook. f., F.B.I. 6 : 168. 1890 ; King & Pantl., l.c. 8 : 338 - 390. 1898. *S. nepalense* f. *albiflorum* Tuyama in Hara, Fl. E. Himal. 450. 1966.

Terrestrial herbs, 50 - 75 cm tall. Tubers ellipsoidal, singly borne at the end of stolons. Leaves almost fleshy, narrowly elliptic, subacute. Spikes 5 - 15 cm long. Flowers rose-pink, rarely white.

Karponang - Changu ; 2400 - 3000 m.

Fl. : Sept. - Oct.

Notes : The flowers posses faint musk odour.

SCHOENORCHIS Bl.

Schoenorchis gemmata, (Lindl.) J.J. Sm. in Nat. Tijdss. Ned. Ind. 72 : 100. 1912. *Saccolabium gemmatum* Lindl. in Edw. Bot. Reg. 24, Misc. 50. 1838 ; Hook. f., F.B.I. 6 : 55. 1890. *Cleisostoma gemmata* (Lindl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 234. t. 313. 1898.

Pendulous, epiphytes without pseudobulbs. Leaves linear, curved, margins infolded, minutely 3 - toothed at apex. Flowers purple.

Gangtok, Ratong - chu - Tingling bridge ; 600 - 1800 m.

Fl. : May - July (-Aug.).

SMITINANDIA Holttum

Smitinandia micrantha (Lindl.) Holttum in Gard. Bull. Singapore 25 : 106. 1969. *Saccolabium micranthum* Lindl., Gen. & Sp. Orchid. 220. 1833 ; Hook. f., F.B.I. 6 : 59. 1890. *Cleisostoma micrantha* (Lindl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 234. t. 312. 1898. *Ascocentrum micranthum* (Lindl.) Holttum in Gard. Bull. Singapore 11 : 275. 1947.

Epiphytes with stout compressed stem, up to 20 cm long, covered by leaf sheaths. Leaves narrowly oblong, keeled, obliquely truncate and bifid at apex. Flowers white with large pink spots ; lip pink or purple.

Teesta valley, up to 900 m.

Fl. : July - Aug.

SPATHOGLOTTIS Bl.

Sapthoglossis ixoides (D. Don) Lindl., Gen. & Sp. Orchid. 120. 1831 ; Hook. f., F.B.I. 5: 814. 1890. *Cymbidium ixoides* D. Don, Prodr. Fl. Nep. 36. 1825.

Terrestrial herbs. Pseudobulbs globular, ovoid, *ca* 1.5 cm long. Leaves 2 - 3, linear, acuminate. Flowers yellow.

Lachung valley ; *ca* 3000 m.

Fl. : July - Aug.

SPIRANTHES L.C. Rich.

Spiranthes sinensis (Pers.) Ames, Orch. 2 : 53. 1908. *Neottia sinensis* Pers., Syn. Pl. 2 : 511. 1807. *N. australis* R. Br., Prodr. 319. 1810. *S. australis* (R. Br.) Lindl. in Bot. Rep. 10 : 5. 823. 1824.

Terrestrial herbs, 15 - 45 cm tall. Roots tuberous. Leaves 4 - 5, alternate, clustered near the base of stem, linear. Flowers pink.

Kyongsla, Phodong, Lachen ; 600 - 2700 m.

Fl. : April - May & Aug. - Sept.

STEREOCHILUS Lindl.

Stereochilus hirtus Lindl. in J. Proc. Linn. Soc. 3 : 58. 1858.
Sarcochilus hirtus (Lindl.) Benth. in Benth. & Hook. f., Gen. Pl. 3 : 576. 1883 ; Hook. f., F.B.I. 6 : 35. 1890.

Epiphytes. Leaves narrowly oblong, apically bifid. Flowers pink ; lip white and pink.

Tendong ; 1800 m.

Fl. : June.

SUNIPIA Lindl.

Sunipia andersonii (King & Pantl.) P.F. Hunt. in Kew Bull. 26 : 183. 1971. *Ione andersonii* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 159. t. 217. 1898.

Epiphytes with ovoid pseudobulbs. Leaves sessile, narrowly elliptic - oblong, apices obtuse, shallowly notched. Flowers yellowish.

Sikkim, s.l. ; ca 1800 m.

S. bicolor Lindl., Gen. & Sp. Orchid. 179. 1833. *Ione bicolor* (Lindl.) Lindl., Fol. Orchid. 3 : 1853 ; Brühl, l.c. 207. *Bulbophyllum bicolor* (Lindl.) Hook. f., F.B.I. 5 : 770. 1890.

Epiphytes. Pseudobulbs obpyriform. Leaf linear, obliquely bifid at apex. Flowers ca 6 mm long.

Sikkim, s.l. ; 1800 - 2100 m.

Fl. : Oct. - Dec.

S. intermedia (King & Pantl.) P.F. Hunt. in Kew Bull. 26 : 184. 1971. *Ione intermedia* King & Pantl. in J. Asiat. Soc. Beng. 65 : 120. 1896.

Terresrial herbs. Pseudobulbs ovoid. Leaf linear, apically notched. Flowers pale green.

Tendong ; 1800 m.

S. paleacea (Lindl.) P.F. Hunt. in Kew Bull. 26 : 184. 1971. *Ione paleacea* Lindl., Fol. Orchid. 2. 1853. *Bulbophyllum paleaceum* (Lindl.) Hook. f., F.B.I. 5 : 769. 1890 ; Brühl, l.c. 208.

Terrestrial herbs. Pseudobulbs ovoid or obpyriform. Leaf narrowly oblong. Flowers translucent, greenish white, pink nerved ; lip purple.

Sikkim, s.l. ; 1800 m.

Fl. : Oct. - Nov.

S. racemosa (J.E. Smith) Tang & Wang in Acta Phytotax. Sin. 1 : 90. 1951. *Stelis racemosa* J. E. Smith in Rees, Cyclop. 34. Stelis No. 10. 1816. *Sunipia scarisa* Lindl., Gen. & Sp. Orchid. 179. 1833 ; Hook. f., F.B.I. 5 : 722. 1890. *Ione scariosa* (Lindl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 161. t. 219. 1898.

Pseudobulbs ovoid, spongy. Leaves broadly lanceolate. Flowers in two rows, greenish, flushed with purple.

Sikkim, s.l. ; 1080 - 1200 m.

Fl. : April - May.

THRIXSPERMUM Lour.

Thrixspermum pygmaeum (King & Pantl.) Holt. in Kew Bull. 14 : 275. 1960. *Sarcochilus pygmaeus* King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 207. t. 277. 1898 ; Brühl, Guide Orch. Sikkim 125. 1926.

Pendulous epiphytes, ca 2.5 cm long. Leaves 2 - 3, fleshy, linear-oblong, acuminate.

Near Dentam, Pemiongtsi (Pemianchi) ; ca 900 m.

Fl. : April - May . Rare.

TROPIDIA Lindl.

Tropidia angulosa (Lindl.) Bl., Coll. Orchid. 122. 1858 ; Hook. f., F.B.I. 6 : 92. 1890. *Cnemidia angulosa* Lindl., Gen. & Sp. Orchid. 463. 1840. *Govindovia nervosa* Wt., Ic. t. 2090. 1853. *Tropidia bellii* Blatt. & McC. in J. Bombay Nat. Hist. Soc. 35 : 730. 1932.

Terrestrial herbs, 20 - 30 cm tall. Leaves 2, broadly elliptic, plicate. Flowers white, flushed with pale ochre-colour.

Teesta valley ; 300 - 600 m.

T. curculigoides Lindl., Gen. & Sp. Orchid. 497. 1840 ; Hook. f., F.B.I. 6 : 93. 1890.

Terrestrial herbs. Stem 30 - 60 cm tall. Leaves narrowly elliptic, acuminate. Flowers white, tinged with yellow.

Sikkim, s.l. : 300 m.

Fl. : Nov.

TAENIOPHYLLUM Bl.

Taeniophyllum crepidiforme (King & Pantl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 245. 1898. *Sarcochilus crepidiformis* King & Pantl. in J. Asiat. Soc. Beng. 64 : 340. 1895.

Minute leafless epiphytic herbs. Racemes erect, *ca* 7 mm long. Flowers greenish ; lip white, stigma purple margined.

In tropical valleys.

Fl. : Sept.

T. retrospiculatum (King & Pantl.) King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 244. 1898. *Sarcochilus retrospiculatus* King & Pantl. in J. Asiat. Soc. Beng. 64 : 340. 1895.

Small, leafless, epiphytic herbs. Flowers green.

Sikkim, *s.l.* ; 1500 m.

Fl. : June.

TAINIA Bl.

Tainia cordata Hook. f., F.B.I. 6 : 193. 1890 ; Karthik. *et al.*, Fl. Ind. Enum. Monocot. 172. 1989.

Terrestrial herbs. Pseudobulbs dark purple. Leaf 30 x 11.5 cm, 7-nerved, oblique at base ; petiole dark purple. Flowers dark brown and red ; lip yellow.

Sikkim Himalaya (Ic. in Hort. Calcutta).

Notes : Karthikeyan *et al.* (1989) mentioned its distribution 'E. Himalaya' probably on the basis of Hooker (1890). There is no further record of its re-collection from any other locality.

T. hookeriana King & Pantl. in J. Asiat. Soc. Beng. 64 : 336. 1895.

Terrestrial herbs. Pseudobulbs ovoid. Leaf plicate, oblong-lanceolate. Flowers greenish with brown lines ; lip white, oblong, lamellate ; lamellae yellow, spotted with pink.

Fl. : April.

T. latifolia (Lindl.) Benth. ex Hook. f., F.B.I. 5 : 820. 1890 ; King & Pantl., I.c. 8 : 103. 1898. *Amia latifolia* Lindl., Gen. & Sp. Orchid. 130. 1831.

Terrestrial herbs. Pseudobulbs olive green, narrowly fusiform. Leaf plicate, elliptic, acute ; petioles 8.7 - 12.5 cm, channelled. Flowers dark brown ; lip yellow.

In tropical valleys ascending to 750 m.

Fl. : March - April.

T. minor Hook. f., F.B.I. 5 : 821. 1890.

Terrestrial herbs. Pseudobulbs narrowly fusiform. Leaves plicate, elliptic. Flowers purplish brown with darker streaks, lip white speckled brown purple at lower half, yellow towards tip, column green.

Chungthang ; 1200 - 1800 m.

Fl. July.

T. wrayana (Hook. f.) J.J. Sm. in Bull. Jard. Bot. Buitzen. 2, 8 : 6. 1912. *Ipsea wrayana* Hook. f., F.B.I. 5 : 812. 1890. *Nephelaphyllum grandiflorum* Hook. f., F.B.I. 6 : 192. 1890.

Terrestrial herbs. Pseudobulbs narrowly fusiform, ca 7.5 cm long. Leaves ovate-elliptic ; petioles not distinguishable from the pseudobulbs. Flowers ca 2.5 cm long ; floral bracts linear-lanceolate, nerves dark brown ; lip subpanudriform, white, pink dotted with an yellow blotch at base and tip, mentum and column yellow.

Teesta valley ; 300 - 450 m.

Fl. : March.

THELASIS Bl.

Thelasis longifolia Hook. f., F.B.I. 6 : 87. 1890.

Epiphytes. Pseudobulbs conical. Leaves oblanceolate, blunt and shallowly bifid at apex. Flowers green.

In tropical valleys.

Fl. : July - Aug.

T. pygmaea (Griff.) Lindl. in J. Proc. Linn. Soc. 3 : 63. 1859 ; Hook. f., F.B.I. 6 : 86. 1890. *Euprobosus pygmaea* Griff. in Calcutta J. Nat. Hist. 5 : 372. t. 26. 1844 ; Wt., Ic. t. 1732. 1851.

Epiphytes. Pseudobulbs ovoid-globose. Leaves linear-oblong, fleshy. Flowers pale green.

Sikkim, s.l.

Fl. : Aug. - Sept.

THUNIA Reichb.f.

Thunia alba (Lindl.) Reichb.f. in Bot. Zeit. (Berlin) 10 : 64. 1852.
Phaius albus Lindl., Gen. & Sp. Orchid. 128. 1830 ; Hook. f., F.B.I. 5 : 818. 1890, *p.p.*

Terrestrial herbs, without pseudobulbs, 30 - 90 cm. Leaves many, sessile, elliptic or oblong, linear, lanceolate. Racemes terminal, drooping, 4 - 6-flowered. Flowers with white or yellow lip having purple or orange veins and 7 - 9 orange ridges on upper surface.

Rangeet valley ; 600 - 1200 m.

Fl. : May.

TIPULARIA Hutt.

Tipularia josephii Reichb. f. ex Lindl. in J. Linn. Soc. 1 : 174. 1857 ; Hook. f., F.B.I. 5 : 708. 1890 ; King & Pantl., l.c. 151 - 152.

Pseudobulbs subglobular, pointed, ca 4.5 mm long. Leaf solitary, broadly ovate, acute, plaited, 7 - nerved. Racemes erect, arising from the base of pseudobulbs, 20 - 30 cm long. Flowers ca 8 mm long, lateral lobes of lip with thick oblong calli.

Sikkim, s.l. ; 2200 - 2900 m.

Fl. : Aug. - Oct.

TRICHOTOSIA Bl.

Trichotosia dasypylla (Par. & Reichb. f.) Kranzl. in Pflanzenr. 50 : 138. 1911. *Eria dasypylla* Par. & Reichb. f. in Trans. Linn. 30 : 147. 1874 ; Hook. f., l.c. 808.

Epiphytic herbs. Flowers yellowish green.

Deorali (Cult.) ; 300 - 600 m (-1400 m).

Fl. : May - June.

T. pulvinata (Lindl.) Kranzl. in Pflanzenr. 50 : 138. 1911. *Eria pulvinata* Lindl. in J. Linn. Soc. 3 : 56. 1858 ; Hook. f., l.c. 807. *E. rufinula* Reichb. f. in Bamb. Gartenzeit. 19 : 13. 1863 ; Hook. f., l.c. 807.

Epiphytic herbs. Flowers whitish pink.

Deorali (Cult.) ; 800 - 1400 m.

Fl. : May.

TYLOSTYLLIS Bl.

Tylostylis discolor (Lindl.) Hook. f. in Ann. Roy. Bot. Gard. Calc. 5 : 22. 1895. *Eria discolor* Lindl. in J. Linn. Soc. 3 : 51. 1858. *E. pulchella sensu* Hook. f., F.B.I. 5 : 802. 1890, *p.p.* non Lindl. 1841.

Herbs. Stem consisting of 4 - 6 pseudobulbs in series. Leaves arising at the joints, coriaceous, sessile. Racemes few flowered. Flowers ochre-yellow, dull white outside ; lip with a purplish brown basal patch.

Sikkim, *s.l.* ; up to 1050 m.

Fl. : Feb.

UNCIFERA Lindl.

Uncifera acuminata Lindl. in J. Linn. Soc. (Bot.) . 3 : 40. 1858. *Saccolabium acuminatum* (Lindl.) Hook. f., F.B.I. 6 : 65. 1890.

Epiphytic herbs. Leaves ovate or elliptic-lanceolate. Flowers pale green.

Sikkim, *s.l.* ; 1800 - 2100 m.

U. lancifolia (King & Pantl.) Schltr. in Orch. 583. 1914. *Saccolabium lancifolium* King & Pantl. in J. Asiat. Soc. Beng. 65 : 122. 1896.

Epiphytic herbs. Leaves linear. Flowers pale yellow.

Rissisum ; *ca* 1800 m.

U. obtusifolia Lindl. in J. Linn. Soc. 3 : 40. 1858. *Saccolabium obtusifolium* (Lindl.) Hook. f., F.B.I. 6 : 65. 1890. *Gastrochilus obtusifolius* (Lindl.) A.S. Rao & Mukherjee in Rec. B.S.I. 20 (2) : 213. 1973.

Epiphytic herbs. Leaves linear-oblong. Flowers pale yellow.

Sikkim, *s.l.* ; 1050 - 1500 m.

VANDA W. Jones ex R. Brown

Vanda alpina Lindl., Fol. Orchid. Vanda 10. 1853 ; Hook. f., F.B.I. 6 : 53. 1890.

Epiphytic herbs. Leaves flat, folded inwards. Flowers greenish or yellowish ; lip without spur.

Gangtok, Namste, Tendong ; 1000 - 1800 m.

Fl. : July.

V. cristata Lindl., Gen. & Sp. Orch. 216. 1833 ; Hook. f., l.c. 53 ; King & Pantl., l.c. 8 : 216. t. 287. 1898.

Epiphytes. Leaves narrowly oblong, conduplicate, truncate and bilobed at apex. Flowers yellowish or olive-green with purplish brown blotches.

Gangtok, Namthang, Tendong ; ca 1500 m.

Fl. : May - June

V. testacea (Lindl.) Reichb.f. in Gard. Chron. 2 : 166. 1877. *Aerides testaceum* Lindl., Gen. & Sp. Orchid. 238. 1833. *V. parviflora* Lindl. in Edw. Bot. Reg. 30. Misc. 45. 1844 ; Hook. f., l.c. 6 : 50. 1890 ; King & Pantl., l.c. 215. 1898.

Epiphytic herbs. Leaves distichous, linear. Lip spurred, bluish purple.

Rangpo ; 350 - 700 m.

V. pumila Hook. f., F.B.I. 6 : 53. 1890.

Epiphytic herbs. Leaves narrowly oblong, unequally bilobed at apex, dilated towards base. Flowers pale yellowish or white ; midlobe of lip pale streaked with purple.

Sikkim, s.l. ; ca 600 m.

VANDOPSIS Pfitz.

Vandopsis undulata (Lindl.) J.J. Smith in Nat. Tijdschr. Ned. Ind. 72 : 77. 1912. *Vanda undulata* Lindl. in J. Linn. Soc. 3 : 42. 1859. *Stauropsis undulatus* (Lindl.) Benth. ex Hook. f., F.B.I. 6 : 27. 1890.

Epiphytes. Leaves narrowly oblong, blunt, retuse. Flowers white flushed with pink, yellowish green striped with pink.

Dentam ; 1500 - 2100 m.

Fl. : April - May.

Notes : Young shoots, rachis and peduncle are green and spotted with purplish brown.

YOANIA Maxim.

Yoania prainii King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 175. 1898.

Saprophytic, aphyllous, succulent, 5 - 10 cm tall, erect, herbs. Rhizomes densely clothed with small scales. Flowers white.

Chungthang ; ca 1800 m.

ZEUXINE Lindl. *nom. cons.*

Zeuxine affinis (Lindl.) Benth. *ex* Hook. f., F. B. I. 6 : 108. 1890.
Monochilus affine Lindl., Gen. & Sp. Orchid. 487. 1840.

Terrestrial herbs, 15 - 30 cm tall. Leaves ovate-lanceolate. Flowers whitish green ; lip yellow ; anthers pink.

Rangeet valley ; ca 450 m.

Fl. : April.

Z. flava (Lindl.) Trimen in J. Ceylon Br. Roy. Asiat. Soc. 9 : 90. 1885 ; Hook. f., F.B.I. 6 : 108. 1890. *Monochilus flavum* Wall. *ex* Lindl., Gen. & Sp. Orchid . 487. 1840. *Hetaeria flava* Lindl. *ex* Wall., Numer. List. 247. 1840.

Terrestrial herbs, 15 - 25 cm tall. Leaves ovate or ovate-lanceolate, clustered at top of the stem.. Petals white ; lip pink at base, yellow at apex.

Dentam - Pamianchi ; 1500 - 2000 m.

Fl. : May - June.

Z. franchetiana King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 292. 1898.

Terrestrial herbs, 15 - 22 cm tall. Leaves few, scattered, linear-lanceolate. Flowers white with a green dash near the base of lip.

Sikkim (?) ; ca 1950 m.

Fl. : Aug.

Z. goodyeroides Lindl., Gen. & Sp. Orchid. 486. 1840 ; Hook. f., F.B.I. 6 : 107. 1890. *Monochilus goodyeroides* (Lindl.) Lindl. in J. Proc. Linn. Soc. Bot. 1 : 187. 1857.

Terrestrial herbs, 10 - 22 cm tall. Leaves 2 - 4, oblong-ovate, acute, green, midrib white. Flowers pink, turning to white.

Sikkim, s.l. ; 1800 m.

Fl. : Sept. - Oct.

Z. nervosa (Wall. *ex* Lindl.) Trimen in J. Ceylon Br. Roy. Asiat. Soc. 9 : 90. 1885 ; Hook. f., F.B.I. 6 : 108. 1890. King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8. 289. t. 385. 1898. *Monochilus nervosum* Wall. *ex* Lindl., Gen. & Sp.Orchid. 487. 1840.

Terrestrial herbs, 15 - 22 cm tall. Leaves 2 - 3, obliquely ovate-lanceolate, acute. Sepals green, petals white ; lobes of lip yellow.

Sikkim, s.l. ; ca 600 m.

Fl. : April.

Z. pulchra King & Pantl. in J. As. Soc. Beng. 65 : 127. 1896.

Terrestrial herbs, 15 - 20 cm tall. Leaves 3 - 4, crowded, ovate, subacute, coriaceous, blackish purple above, midrib white. Flowers pale brown ; lip white above, yellow below.

Lachung valley ; ca 2250 m.

Fl. : Aug.

Z. reflexa King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 : 291. 1898.

Terrestrial herbs, 17 - 22 cm tall. Leaves narrowly ovate, acute. Sepals green, petals white ; lip yellow, sac white.

Sikkim (?) ; 1050 m.

Fl. : April.

Z. strateumatica (L.) Schlechter in Fedde Repert. Beih. 1 : 77. 1911. *Orchis strateumatica* L., Sp. Pl. 943. 1753. *Pterygodium sulcatum* Roxb., Fl. Ind. 3 : 452. 1832. *Zeuxine sulcata* (Roxb.) Lindl. in Royle, Ill. Bot. Himal. Mount. 1 : 368. 1839 ; Hook. f., F.B.I. 6 : 106. 1890 ; King & Pantl. in l.c. 8 : 286. t. 381. 1898.

Terrestrial herbs, 5 - 25 cm tall. Leaves linear, acuminate. Flowers white or pink ; lip purple at base, yellow at apex.

Sikkim, s.l. ; up to 1200 m.

Fl. : Jan.

ZINGIBERACEAE

(S. Kumar)

Alpinia Roxb. *nom. cons.*

Alpinia bracteata Roxb., Fl. Ind. 1 : 63. 1820 ; Baker in Hook. f., F.B.I. 6 : 255. 1892.

Herbs. Leaves oblong-lanceolate, villous beneath. Brancheoles pale green. Flowers with white lip with variegated red and yellow centre. Capsules globose.

Sikkim, s.l. ; ca 300 m.

Fl. & Fr. : May - Oct.

A. nigra (Gaertn.) Burtt in Notes Roy. Bot. Gard. Edinb. 35 : 213. 1977. *Zingiber nigrum* Gaertn., Fruct. 1 : 35. 5. 12. 1788. *Alpinia allughas* (Retz.) Rosc. in Trans. Linn. Soc. London 8 : 346. 1807 ; Baker in Hook. f., F.B.I. 6 : 253. 1892.

Biennial or perennial herbs with tuberous rhizomes. Leaves glabrous, acuminate. Brancheoles small, sessile oblong-lanceolate, glabrous. Flowers with white bilobed lip having a pair of pink lines at centre. Capsules globose, up to 5 cm in diam.

Teesta valley.

Fl. & Fr. : May - Oct.

Notes : Young roots are eaten as vegetable.

A. malaccensis (Burm. f.) Rosc. in Trans. Linn. Soc. 8 : 345. 1807 ; Baker in Hook. f., F.B.I. 6 : 255. 1892. *Maranta malaccensis* Burm. f., Fl. Ind. 2. 1768.

Herbs with several stout stems. Leaves long petioled, oblong-lanceolate, villous beneath. Brancheoles white or pale brownish. Flowers with yellow lip having deep red and yellow centre. Capsules globose, yellowish to orange-red.

Dikchu, Chungthang ; up to 1350 m.

Fl. & Fr. : March - May.

Notes : An essential oil viz. 'Amadi' is obtained from rhizomes.

AMOMUM Roxb. *nom. cons.*

Amomum aromaticum Roxb., Fl. Ind. 1 : 44. 1820 ; Baker in Hook. f., F.B.I. 6 : 241. 1892. MORANG HATCHI (Hindi, Beng.).

Herbs. Rhizomes faintly aromatic, white. Leaves linear-lanceolate, caudate-acuminate, glabrous, aromatic. Flowers white, yellow and white with a medium yellow stripe and few pinkish veins radiating towards base. Capsules trigonous, smooth, fleshly.

Rishap, Melli.

Fl. & Fr. : Throughout the year.

Notes : Seeds are used as a condiment and for flavouring food.

A. corynostaichyum Wall., Pl. Asiat. Rar. 1 : 48. t. 58. 1830 ; Baker in Hook. f., F.B.I. 6 : 238. 1892.

Herbs. Leaves petioled, lanceolate, glabrous. Spikes globose. Flowers with white subrhomboid lip having yellow median band and red streaks at the base. Capsules ellipsoid, faintly many ridged, hairy.

Sikkim, s.l. ; ca 750 m.

Fl. & Fr. : June - July.

A. dealbatum Roxb., Fl. Ind. 1 : 43. 1820 ; Baker in Hook. f., F.B.I. 6 : 239. 1892.

var. dealbatum

Perennial herbs with aromatic rhizomes and leafy stems. Leaves subsessile, oblong - lanceolate. Bracts red. Flowers with a white lip having yellow blotch at the base. Capsules oblong-subglobose, with 9-crenulate winged ribs, pinkish red at maturity.

Sithong, Singhik, Dikchu, Jinglam ; 500 - 1500 m.

Fl. & Fr. : April - May.

Notes : It is used as a substitute for large cardamom.

var. sericeum (Roxb.) Baker in Hook. f., F.B. I. 6 : 239. 1892.
A. sericeum Roxb., Fl. Ind. 1 : 46. 1820.

Herbs. Leaves white, sericeous beneath. Flowers white, lip variegated with pink veins on a yellow ground. Capsules ovate-oblong, not winged.

Rangpo valley, up to 1000 m.

Fl. & Fr. : April - May.

A. kingii Baker in Hook. f., F.B.I. 6 : 241. 1892.

Herbs with stout leafy stems. Leaves oblong-lanceolate. Flowers white. Lip obscurely 3-lobed, tinged with yellow, narrowed suddenly to a claw.

Sikkim, s.l.

Fl. & Fr. : June - July. Endemic (?)

Notes : It is represented by a single collection available at CAL.

A. subulatum Roxb., Pl. Corom. 3 : 75. t. 277. 1819 ; Baker in Hook. f., F.B.I. 6 : 240. 1892.

Herbs with leafy stem. Leaves subsessile, oblong-lanceolate, glabrous. Flowers white. Lip yellow with deep yellow veins. Capsules ellipsoid, flattened, longitudinally striate with crenulate wings, red brown.

Cultivated throughout tropical and subtropical regions ; up to 1800 m.

Fl. & Fr. : April - May.

Notes : It is one of the major cash crops of Sikkim. Seeds are used for flavouring food.

CAULOKAEMPFERIA K. Larsen

Caulokaempferia secunda (Wall.) K. Larsen in Bot. Tidssk. 60 : 170. 1964 ; Rao & Verma in Bull. Bot. Surv. India 14 : 125. 1972. *Kaempferia secunda* Wall. in Roxb., Fl. Ind. 1 : 19. 1820 ; Baker in Hook. f., F.B.I. 6 : 223. 1892.

Small herbs with leafy stem. Leaves sessile-subsessile, oblong-ovate, glabrous. Flowers pink or red. Lips notched. Capsules 1 cm long ; seeds whitish.

Sikkim, s.l. ; up to 1200 m.

C. sikkimensis (King ex Baker) K. Larsen in Saert. Bot. Tidssk. 60 : 169. 1964. *Kaempferia sikkimensis* King ex Baker in Hook. f., F.B.I. 6 : 223. 1892.

Small herbs with slender leafy stem. Leaves sessile, lanceolate. Spikes 1-flowered. Flowers white ; corolla lobes dark lilac ; lip suborbicular, white.

Ranipool, Rangpo, Gangtok.

Fl. & Fr. : July - Aug.

CAUTLEYA (Benth.) Hook. f.

Cautleya cathcartii Baker in Hook. f., F.B.I. 6 : 208. 1890.

Perennial robust herbs. Leaves sessile, linear-lanceolate - oblong, acuminate, glabrous. Spikes 15 - 20 - flowered. Bracts green. Flowers bright yellow. Capsule red. Seeds white.

Sikkim, s.l.

Fl. & Fr. : May - July.

C. gracilis (J.E. Smith) Dandy in Journ. Bot. 70 : 328. 1954. *Roscoea gracilis* J.E. Smith in Trans. Linn. Soc. 13 : 460. 1812. *C. lutea*

(Royle) Hook. f. in Bot. Mag. t. 699. 1888 ; Baker in Hook. f., F.B.I. 6 : 208. 1890. *Roscoea lutea* Royle, Illustr. Bot. Himal. 361, t. 89, f.2. 1839.

Herbs with thick rhizomes. Leaves sessile, lanceolate. Spikes terminal, lax. Bracts purplish green. Flowers yellow. Capsules red. Seeds conical, ridged, dark blue or black.

Lachung, Chungthang, Dikchu.

Fl. & Fr. : June - Oct.

C. robusta Baker in Hook. f., F.B.I. 6 : 208. 1890.

Perennial robust herbs. Leaves sessile, oblong, acute. Spikes dense, many-flowered. Bracts reddish. Flowers deep orange-yellow. Capsule depressed globose. Seeds ovoid or compressed, pink, enclosed in red or yellow aril.

Sikkim, s.l.

Fl. & Fr. : June - Oct.

C. spicata (J.E. Smith) Baker in Hook. f., F.B.I. 6 : 209. 1890. *Roscoea spicata* J.E. Smith in Trans. Linn. Soc. 13 : 461. 1812.

Perennial herbs. Leaves up to 10 cm broad. Spikes shortly peduncled. Bracts red, tubular. Flowers bright yellow. Capsule globose, red. Seeds black.

Fl. & Fr. : June - Nov.

COSTUS L.

Costus lacerus Gagnep. in Bull. Soc. Bot. France 3, 50 : 261. 1903 ; Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 59. 1978.

Robust herbs. Leaves sessile, ovate-lanceolate, subacuminate, Spikes sessile, oviform. Bracts coarsely hairy, ciliate, margins torn. Flowers creamy white. Seeds 4 - 5 angled, sky-blue.

Teesta valley.

Fl. & Fr. : July - Oct.

C. speciosus (Koen.) J.E. Smith in Trans. Linn. Soc. London 1 : 249. 1800 ; Baker in Hook. f., F.B.I. 6 : 249. 1892. *Banksia speciosa* Koen. *apud* Retz., Obs. Bot. 3 : 75. 1783.

Robust herbs. Leaves petioled, oblanceolate-obovate. Spikes compact. Bracts ovate, cuspidate. Flowers white ; lip white with a

median yellow band and crimped margins. Capsule globose-oblong, red. Seeds 4 - 5 angled, black.

Rangeet, Rangpo, Riang, Gangtok ; up to 1700 m.

Fl. & Fr. : June - Oct.

Notes : Roots edible, bitter, and used in medicine. Plants are of ornamental value as well.

CURCUMA Roxb. *nom. cons.*

Curcuma angustifolia Roxb. in *Asiat. Res.* 11 : 338. t. 3. 1810 ; Baker in *Hook. f.*, *F.B.I.* 6 : 210. 1890.

Perennial herbs. Rhizomes small, globose, pale yellow inside. Leaves petioled, up to 8 cm broad. Coma-bracts purple, forming a cupule. Flowers bright yellow.

Sikkim, *s.l.* ; up to 1700 m.

Notes : It is cultivated for production of East Indian Arrowroot starch.

C. aromatica Salisb., *Parad. London* 20 : t. 96. 1808 ; Baker in *Hook. f.*, *F.B.I.* 6 : 210. 1890.

Perennial herbs. Rhizomes tuberous, strongly aromatic, pale yellow inside. Leaves petioled, oblong. Coma-bracts pink. Flowers green.

Teesta valley, Rongsong.

Fl. & Fr. : May - June.

Notes : It is used in medicine and perfume and is cultivated.

C. caesia Roxb. in *Asiat. Res.* 11 : 334. 1810 ; Baker in *Hook. f.*, *F.B.I.* 6 : 212. 1890.

Herbs. Rootstocks large, ovoid ; tubers sessile, pale grey inside. Leaves petioled, green with a broad purple-brown band in middle. Spikes densely flowered. Coma-bracts red. Flowers pale yellow. Capsules trigonous ; seeds arillate, white.

Sikkim, *s.l.* ; ca 600 m.

Fl. & Fr. : May - June.

Notes : It is cultivated for perfume industry.

C. longa L., *Sp. Pl.* 1 : 2. 1753 ; Baker in *Hook f.*, *F.B.I.* 6 : 214. 1890.

Herbs. Rhizomes tuberous, strongly aromatic, pale yellow to deep orange, yellow inside. Coma-bracts pinkish white. Flowers white with creamy white top having deep yellow band.

Cultivated up to 1700 m.

Fl. & Fr. : June - Nov.

C. zedoaria (Berg.) Rosc. Monandv. Pl. t. 109. 1828 ; Baker in Hook. f., F.B.I. 6 : 210. 1890. *Amomum zedoaria* Berg. Mat. Med. 41. 1788.

Herbs. Rhizomes pale yellow, faintly aromatic inside. Leaves petioled, oblong with purple tinge on both surfaces. Coma-bracts bright red. Flowers pale yellow.

Ranipol, Teesta valley ; up to 2000 m.

Fl. & Fr. : April - May.

Notes : Cultivated throughout India for Starch production.

CURCUMORPHA Rao & Verma

Curcumorpha longiflora (Wall.) Rao & Verma in Bull. Bot. Surv. India 13 : 339. 1971. *Gastrochilus longiflora* Wall., Pl. Asiat. Rar. 1 : 22. t. 25. 1829 ; Baker in Hook. f., F.B.I. 6 : 251. 1892.

Herbs. Rhizome short, faintly aromatic. Leaves oblong-lanceolate, petioled. Spikes purplish green. Flowers white.

Teesta valley.

Fl. & Fr. : July - Sept.

GLOBBA L.

Globba andersonii Clarke ex Baker in Hook. f., F.B.I. 6 : 202. 1890.

Perennial herbs with leafy stem. Leaves oblong-lanceolate, caudate, pubescent beneath. Panicle lax, rhachis hairy. Flowers bright yellow. Capsule warty.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

G. clarkei Baker in Hook. f., F.B.I. 6 : 201. 1890 ; Rao & Verma, l.c. 11 : 246. 1969.

Perennial herbs. Leaves lanceolate, caudate, hairy at base beneath. Rhachis glabrous. Flowers orange yellow. Capsules wrinkled, faintly 3-lobed, oblong. Bulbils tiny, white or brownish replacing flowers at apex.

Sikkim, s.l. ; up to 2400 m.

Fl. & Fr. : June - Oct.

G. multiflora Wall. ex Baker in Hook. f., F.B.I. 6 : 202. 1890 ; Rao & Verma, l.c. 14 : 118. 1972.

Perennial herbs. Leaves oblong-lanceolate, caudate hairy beneath. Panicles with hairy rachis. Flowers yellow. Capsules subglobose.

Sikkim, s.l. ; ca 1200 m.

Fl & Fr. : May - Sept.

G. racemosa J. E. Smith, Exot. Bot. 2 : 115. t. 117. 1804 ; Rao & Verma, l.c. 14 : 118. 1972. *G. orixensis* Roxb. in Asiat. Res. 11: 358. 1810.

var. **racemosa**

Perennial herbs. Leaves oblong-lanceolate, subcaudate. Rachis glabrous. Flowers yellow or orange with dark yellow or purple tinged tips. Capsules globose.

Rhenock, Riang, Gangtok.

Fl. & Fr. : May - Oct.

var. **hookeri** (Clarke ex Baker) S. Kumar, stat. et comb. nov. *G. hookeri* Clarke ex Baker in Hook. f., F.B.I. 6 : 202. 1890.

Differs from var. *racemosa* in having broader leaves, very hairy on the midrib beneath ; bright yellow flowers with unspotted lip.

Kolbong, Riang, Rangsong, Dikchu.

Fl. & Fr. : May - Sept.

G. siamensium (Koen.) Rao & Verma in Bull. Bot. Surv. India 14 : 118. 1972. *Hura siamensis* Koen. in Retz., Obs. Bot. 3 : 49. 1783. *G. versicolor* J.E. Smith, Exot. Bot. f. 117 ABC. 1804 ; Baker in Hook. f., F.B.I. 6 : 205. 1890. *G. expansa* Wall. ex Baker, l.c.

Perennial herbs. Leaves oblong, caudate. Branches and rhachis slender. Flowers yellow tinged with lilac smaller corolla tube. Capsules cylindric smooth.

Sikkim, s.l.

Fl. & Fr. : June - Oct.

HEDYCHIUM Koen.

Hedychium aurantiacum Wall. *ex* Rosc., Monanar. Pl. t. 39. 1828 ; Baker in Hook. f., F.B.I. 6 : 232. 1892.

Perennial rhizomatous aromatic herbs. Leaves lanceolate, acuminate. Spikes lax. Flowers bright orange-yellow, with a deep orange blotch at mouth.

Rhenock; *ca* 600 m.

Fl. & Fr. : July - Feb.

H. coccineum Buch. - Ham. *ex* J.E. Smith in Rees., Cyclop. 17 : 5. 1811 ; Baker in Hook. f., F.B.I. 6 : 231. 1892. *H. coccineum* var. *longifolium* Baker, *l.c.*

var. coccineum

Perennial herbs. Leaves linear-lanceolate, purplish villous beneath. Flowers small, bright crimson.

Riang, Chungthang, Great Rangeet.

Fl. & Fr. : June - Dec.

var. squarrosum (Buch.-Ham. *ex* Wall.) Baker in Hook. f., F.B.I. 6 : 231. 1892. *H. squarrosum* Buch. - Ham. *ex* Wall. in Hook., Kew J. Bot. 5 : 372. 1853.

Differs from var. *coccineum* in having glabrous narrow leaves and small flowers.

Sikkim, s.l. ; 2400 - 2700 m.

Fl. & Fr. : June - Dec.

H. coronarium Koen. in Retz., Obs. Bot. 3 : 73. 1783 ; Baker in Hook. f., F.B.I. 6 : 225. 1892 ; Rao & Verma, *l.c.* 11 : 121. 1971.

Perennial sweet scented herbs. Leaves oblong-lanceolate, acuminate, appressed hairy beneath. Spikes densely flowered. Flowers white with a yellow blotch at mouth.

Sikkim, s.l. ; up to 130 m.

Fl. & Fr. : June - Oct.

Notes : Very variable in shape of staminodes and flowers.
Ornamental.

H. densiflorum Wall. in Hook., Kew Journ. Bot. 5 : 368. 1853 ;
Baker in Hook. f., F.B.I. 6 : 227. 1892 ; Rao & Verma, l.c. 14 : 130.
1972.

Perennial herbs. Leaves linear-lanceolate, glabrous beneath.
Spikes moderately dense flowered, sessile. Flowers small, orange-yellow, turning reddish at maturity.

Lachen ; up to 2400 m.

Fl. & Fr. : June - Oct.

H. ellipticum Buch.-Ham. ex J. E. Smith in Rees Cyclop. 17. n.
2. 1811 ; Baker in Hook. f., F.B.I. 6 : 228. 1892 ; Rao & Verma in Bull.
Bot. Surv. India 14 : 131. 1972.

Perennial, terrestrial or epiphytic herbs. Leaves oblong, glabrous.
Spikes dense flowered. Flowers fragrant, white, turning yellow with age.

Chungthang, Teesta valley, Rhenock, Dikchu ; 600 - 1500 m.

Fl. & Fr. : June - Sept.

Notes : Ornamental.

H. gardnerianum Sheppard ex Ker.-Gawl. in Edw. Bot. Reg. 8.
t. 774. 1824 ; Baker in Hook. f., F.B.I. 6 : 230. 1892 ; Rao & Verma,
l.c. 14 : 133. 1972.

Perennial herbs with leafy stems. Leaves subsessile, oblong,
caudate, white puberulent beneath. Spikes lax-flowered. Flowers bright
yellow. Capsules persistent. orange-red within.

Sikkim, s.l. ; 900 - 2400 m.

Fl. & Fr. : July - Dec.

H. gracile Roxb., Fl. Ind. 1 : 14. 1820 ; Baker in Hook. f., F.B.I.
6 : 229. 1892 ; Rao & Verma, l.c.; *H. gracile* var *glacum* Baker, l.c.

Perennial herbs. Leaves subsessile, oblong-lanceolate. Spikes
moderately dense-flowered. Flowers white or pale yellow tinged.
Capsules globose, glabrous.

Lachung, Gangtok, Great Rangeet.

Fl. & Fr. : May - Dec.

H. greenii W.W. Smith in Rec. B.S.I. 4 : 272. 1913 ; Rao & Verma, l.c. 14 : 130. 1972..

Perennial herbs. Leaves oblong, sparsely hairy. Spikes purplish red. Flowers red. Bulbils present.

Sikkim, s.l.

Fl. & Fr. : July - Sept.

H. spicatum Buch.-Ham. ex J.E. Smith in Rees Cyclop. 17 : 3. 1811 ; Baker in Hook. f., F.B.I. 6 : 227. 1892.

var. spicatum

Perennial herbs. Leaves oblong-lanceolate, glabrous. Spikes densely flowered. Flowers white, with deeply bifid lip. Capsules subglobose, glabrous.

Sikkim, s.l. ; up to 3000 m.

Fl. & Fr. : July - Sept.

var. acuminatum (Rosc.) Wall. in Hook., Kew Journ. Bot. 5 : 328. 1853 ; Baker in Hook. f., F.B.I. 6 : 227. 1892 ; Rao & Verma in Bull. Bot. Surv. India 14 : 130. 1972. *H. acuminatum* Rosc., Monandr. Pl. t. 47. 1828.

Differs from var *spicatum* in having abruptly acuminate leaves, lax few-flowered spikes and convolute bracts.

Phodonchen, Lepcha Zagat (Dzongu).

H. thyrsiforme Buch.-Ham. ex J.E. Smith in Rees Cyclop. 17. 1811 ; Baker in Hook. f., F.B.I. 6 : 230. 1892 ; Rao & Verma, l.c. 14 : 132. 1972.

Perennial aromatic herbs. Leaves oblong-lanceolate, villous beneath. Spikes compact, densely flowered. Flowers white, fragrant ; lip deeply bifid with a yellow blotch at base.

Rangeet valley ; up to 1200 m.

Fl. & Fr. : Aug. - Sept.

HEMIORCHIS Kurz

Hemiorchis pantlingii King in Ann. Roy. Bot. Gard. Calc. 5 : 163. t. 198 - 199. 1896 ; Rao & Verma, l.c. 14 : 119. 1972.

Perennial, rhizomatous herbs. Leaves oblong-lanceolate, produced after flowering. Peduncles sheathed. Hispidulous corolla and

yellow lip with purplish median line and orange tip. Capsules puberulous, 3 - valved.

Sikkim, s.l. ; up to 1200 m.

Fl. & Fr. : March - April.

HORNSTEDTIA Retz.

Hornstedtia linguiformia K. Schum. in Engl. Pflanzenr. 20 : 196. 1904. *Amomum linguiforme* (Shult.) Baker in Hook. f., F.B.I. 6 : 235. 1892. *Elettaria linguiformis* Schult., Mant. Pl. 1 : 24. 1822.

Perennial herbs with creeping rhizomes. Leaves subsessile, lanceolate, acuminate, glabrous. Spikes ellipsoid. Flowers red ; lip yellow having a red median band.

Sikkim, s.l. ; up to 1500 m.

Fl. & Fr. : March - Aug.

KAEMPFERIA L.

Kaempferia angustifolia Rosc. in Trans. Linn. Soc. 8 : 351. 1807 ; Baker in Hook. f., F.B.I. 6 : 219. 1890.

Herbs with tuberous rootstock. Leaves sessile, lanceolate. Spikes sessile. Flowers white with small bracts and lilac lip.

Sikkim, s.l.

Fl. : May - June.

K. involucrata King ex Baker in Hook. f., F.B.I. 6 : 221. 1890 ; Rao & Verma, l.c. 14 : 124. 1972.

Herbs with subglobose tubers. Leaves petioled, oblanceolate. Flowers white, enclosed in cup-like structure formed of two bracts on a flat receptacle ; lip white with a yellow blotch in the centre.

Sikkim, s.l. ; up to 1000 m.

Fl. : May - June.

K. rotunda L., Sp. Pl. 3. 1753 ; Baker in Hook. f., F.B.I. 6 : 222. 1890 ; Rao & Verma, l.c. 14 : 124. 1972. BHUICHAMPA (Nep.).

Herbs with aromatic rhizomes. Leaves lanceolate, appressed hairy beneath. Spikes subsessile. Bracts purplish white. Calyx purplish-dotted. Corolla white ; lip lilac or red, bilobed.

Rishap.

Fl. : March - May.

ROSCOEAE J.E. Smith

Roscoea alpina Royle, Ill. Bot. Himal. 361. t. 89. 1839 ; Baker in Hook. f., F.B.I. 6 : 207. 1890. *R. alpina* var. *minor* Royle, l.c. t. 89. *R. longifolia* Baker, l.c. 208.

Herbs with small, slender stem. Leaves 2 - 3, sessile, sublinear-lanceolate-oblong, acuminate. Flowers purple, in 2 - 5 - flowered sessile spikes.

Sikkim, s.l. ; 2400 - 3200 m.

Fl. & Fr. : July - Oct.

R. auriculata K. Schum. in Engl., Pflanz. 4 : 118. 1904 ; Cowley in Kew Bull. 36 : 757. 1982.

Herbs. Leaves linear to broadly elliptic, auriculate at base. Flowers deep purple with some parts white, deflexed lip and staminodes with V - shaped purple marking at base. Seeds brownish, slightly contracted near middle.

Lachung, Chungthang, Panthgang, Bakhim, Lamteng.

Fl. & Fr. : May - Sept.

R. purpurea J.E. Smith, Exot. Bot. 11 : 97. t. 108. 1804 ; Baker in Hook. f., F.B.I. 6 : 207. 1890 ; Rao & Verma, l.c. 14 : 119. 1972 ; Cowley in Kew Bull. 36 : 752. 1982. *R. purpurea* var. *exilis* Baker, l.c. 206. *R. purpurea* var. *procera* (Wall.) Baker, l.c. 208.

Herbs with fusiform root. Stem leafy. Leaves oblong-lanceolate. Spikes sessile, terminal, 2 - 4 - flowered. Flowers purple.

Lachung, Bakhim, Lamteng ;

Fl. & Fr. : May - Sept.

ZINGIBER Boehm.

Zingiber capitatum Roxb. in Asiat. Res. 11 : 348. 1810 ; Baker in Hook. f., F.B.I. 6 : 248. 1892 ; Rao & Verma, l.c. 14 : 135. 1972.

var. **capitatum**

Perennial rhizomatous herbs with leafy stem. Leaves sessile, linear-lanceolate, pubescent beneath. Spike terminal dense-flowered.

Flowers yellow with pale yellow unspotted lip. Capsule oblong, bright red. Seeds brownish black.

Sikkim terai.

Fl. & Fr. : July - Sept.

var. **elatum** (Roxb.) Baker in Hook. *f.*, F.B.I. 6 : 249. 1892.
Zingiber elatum Roxb., Fl. Ind. 1 : 57. 1820.

Differs from var. *capitatum* in having thin linear grasslike leaves ; stouter spikes ; larger flowers and yellow lip with brownish tinge ; lateral staminodes very short, free.

Sikkim, *s.l.*

Z. clarkei King *ex* Benth. in Benth. & Hook. *f.*, Gen. Pl. 3 : 646. 1882 ; Hook. *f.*, F.B.I. 6 : 248. 1892.

Perennial herbs with leafy stem. Leaves oblong-lanceolate, finely pubescent beneath. Spikes lateral, cernuous. Flowers yellow with purple brown tinge.

Sikkim, *s.l.* ; up to 1600 m.

Fl. & Fr. : Aug. - Dec.

Z. chrysanthum Rosc., Monandr. Fl. t. 86. 1827 ; Baker in Hook. *f.*, F.B.I. 6 : 243. 1892.

Very large herbs. Leaves oblong-lanceolate, pubescent beneath. Spikes globose-ovoid with very short peduncles. Corolla tube bright red. Lip yellow unspotted. Capsules ellipsoid-oblong. Seeds brown, subglobose.

Sikkim, *s.l.* ; ca 1800 m.

Fl. & Fr. : July - Sept.

Z. officinale Rosc. in Trans. Linn. Soc. 8 : 348. 1807 ; Baker in Hook. *f.*, F.B.I. 6 : 248. 1892.

Herbs. Rhizomes aromatic, pale yellow inside, bearing sessile tubers. Stems leafy. Leaves linear-lanceolate. Corolla tube greenish. Lip purplish black.

Sikkim, *s.l.* ; up to 1200 m.

Fl. : Sept. - Oct.

Notes : Cultivated. It is an important cash crop of Sikkim.

Z. purpureum Rosc. in Trans. Linn. Soc. 8 : 348. 1807. Z. cassumunar Roxb. in Asiat. Res. 11 : 347. t. 5. 1810 ; Baker in Hook. f., F.B.I. 6 : 248. 1892.

Herbs. Rhizomes aromatic, deep yellow inside. Stems leafy. Leaves sessile, lanceolate-linear, acuminate, pubescent along midrib beneath. Spikes ellipsoid. Flowers yellow. Capsules globose.

Sikkim, s.l. ; up to 1500 m.

Fl. & Fr. : July - Aug.

Z. rubens Roxb. in Asiat. Res. 11 : 348. 1810 ; Baker in Hook. f., F.B.I 6 : 243. 1892.

Herbs with stout leafy stem. Leaves sessile, oblong-lanceolate, acuminate, puberulous beneath along midrib. Spikes subglobose. Corolla lobes red ; lip yellowish white with red spots and streaks. Capsules ellipsoid, reddish.

Sikkim, s.l. ; up to 1500 m.

Fl. & Fr. : Aug. - Sept.

Z. zerumbet (L.) Rosc. ex J.E. Smith, Exot. Bot. 2 : 105. t. 112. 1804. *Amomum zerumbet* L., Sp. Pl. 1. 1753 ; Hook. f., F.B.I. 6 : 247. 1892.

Herbs with tuberous rhizomes and stout leafy stem. Leaves oblong-lanceolate, glabrous beneath. Spikes very dense, oblong. Flowers with bright yellow lip. Capsules white, oblong.

Sikkim, s.l.

Fl. & Fr. : Aug. - Oct.

CANNACEAE

(R.C. Srivastava)

CANNA L.

Canna edulis Ker-Gawler in Edw. Bot. Regist. 9 : t. 775. 1824.
PHUL-TARUL (Nep.).

Stout herbs, up to 60 cm tall. Leaves tinge purple. Flowers red or yellow.

Jorethang, near Legship, Yoksum, Phamtam, Namthang ; 1200 - 1800 m.

Fl. & Fr. : April - June.

Cultivated for its edible rhizomes.

C. indica L., Sp. Pl. 1 : 1753. *C. orientalis* Roscoe, Monandr. Pl. t. 12. 1828. *C. indica* var. *orientalis* (Roscoe) Baker in Hook. f., F.B.I. 6 : 260. 1892.

var. indica

Stout herbs, up to 1.5 m tall. Leaves elliptic-oblong, pale green. Flowers red or yellow, with 3 outer staminodes, *ca* 4 x 0.5 cm.

Singtam, Rangpo, Jorethan, Gangtok, Tadong ; up to 1000 m.

Fl. & Fr. : April - June. Cult.

var. speciosa (Sims) Baker in Hook. f., F.B.I. 6 : 261. 1892. **C. speciosa** Sims in Curtis Bot. Mag. 49 : t. 2317. 1822.

Herbs, up to 1.5 m high. Leaves broadly lanceolate, with abruptly contracted to cuspidate apex. Inflorescence *ca* 30 cm, with short, suberect branches. Flowers dull crimson, orange streaked.

Jorethang, Gangtok ; 500 - 1750 m.

Fl. & Fr. : June - Sept.

Canna x generalis L.H. Bailey is cultivated at Gangtok, Yoksum, Legship etc.

MUSACEAE

(B. Krishna)

MUSA L.

Musa balbisiana Colla, Mem. Gen. Musa. 56. 1820. *M. sapientum sensu* Baker in Hook.f., F.B.I. 6 : 262. 1892, *p.p. non* L. 1753. *M. paradisiaca* ssp. *sapientum* (L.) K. Schum. in Engl. Pflanzenr. 45 : 20. 1900.

Perennial large herbs with branching rhizomes. Pseudostems 2 - 6 m high. Flowers 10 - 20, in a single row, pruinose, yellow. Fruits pale yellow.

Cultivated throughout Sikkim up to 1750 m.

Fl. & Fr. : Jan. - Nov.

M. hookeri King *ex* Cowan & Cowan, Trees Bengal 135. 1929.

Large perennial herbs with branching rhizomes. Pseudostems 3.6 - 4.5 m tall. Fruits compact, few, 4 - 5 cm long.

Sikkim, s.l. ; ca 1667 m.

Fl. & Fr. : March - Sept.

M. sikkimensis Kurz in J. Agri. Hort. Soc. India 14 : 164. 1867 ; Baker in Hook. f., l.c. 262. 1892.

Perennial herbs with stolons. Pseudostems short, 1 - 1.5 m long. Flowers 6 - 8, in two rows, yellow. Fruits 7 - 10 cm long.

Throughout Sikkim on hill slopes.

Fl. & Fr. : Feb. - Oct.

M. velutina Wendl. & Drude in Regel Garten. Flora 24 : 65. t. 823. 1875 ; Baker in Hook. f., l.c. 263. *M. dasycarpa* Kurz in J. Agri. Hort. Soc. India 14 : 301. 1867 ; Baker in Hook.f., l.c. 263.

Perennial large herbs with branched rhizomes. Pseudostems 1.2 - 1.6 m high. Flowers 2 - 3, in a single row, orange-yellow. Fruits subsessile, pinkish red, hairy.

In Tessta valley, Singtam.

Fl. & Fr. : Jan. - Nov. Ornamental.

IRIDACEAE

(R.C. Srivastava)

BELAMCANDA A. Adans.

Belamcanda chinensis (L.) DC. in Redoute, Liliac. 3 : t. 121. 1857 ; Hook.f., F.B.I. 6 : 277. 1892. *Ixia chinensis* L., Sp. Pl. 36. 1753.

Herbs, up to 1.5 m tall, with brown spotted orange flowers.

Jorethang, Yoksum, N. of Tashiding, Tingbong ; up to 2200 m.

Fl. & Fr. : June - Aug. Cult.

CROCOSMIA Planch.

Crocosmia x crocosmiflora (Lemoine) N.E. Br. is grown in gardens at Gangtok, Yoksum etc. localities.

GLADIOLUS L.

Gladiolus callianthus Marais, **G. natalensis** (Ecklon) Hooker, **G. undulatus** L. and many hybrids of **G. insignis** Paxt. & **G. hortulamus** L.H. Bailey are grown in gardens.

IRIS L.

Iris clarkei Baker ex Hook. f., F.B.I. 6 : 275. 1892. *I. himalaica* Dykes in Gard. Chron. 45 : 3, 36. 1909.

Herbs, up to 60 cm tall. Leaves linear or ensiform. Flowers lilac, blotched with violet ; throat yellow.

Changu, Dikchu, Gnathong, Kyangosla, Kupup, Karponang, Laghep, Padamchan, Yakla ; 3000 - 3500 m.

Fl. & Fr. : April - May.

I. decora Wall., Pl. Asiat. Rar. 1 : 77. t. 86. 1830. *I. nepalensis* D. Don, Prodr. Fl. Nep. 54. 1825 ; Hook. f., F.B.I. 6 : 273. 1892.

Herbs up to 30 cm tall. Leaves linear. Flowers pale-liliac.

Chungthang ; 1800 - 4000 m.

Fl. & Fr. : June- Aug.

I. goniocarpa Baker in Gard. Chron. 2 : 710. 1876 ; Hook. f., F.B.I. 6 : 274. 1892.

Herbs, up to 15 cm tall. Leaves narrow linear. Flower heads solitary.

Lachung, Thangu, Yatung ; 3600 - 4400 m.

Fl. : June - July.

I. tectorum Maxim.

Herbs, vegetative shoots with fans of 3 - 4 curved, ensiform, acute leaves. Flower blue violet.

Yoksum ; 2370 - 2750 m.

Fl. & Fr. : April - May.

AMARYLLIDACEAE

(D.C.S. Raju, R.C. Srivastava & S. Singh)

CRINUM L.

Crinum amoenum Roxb. ex Ker - Gawler in J. Sci. Arts. London 3 : 106. 1817 ; Hook. f., F.B.I. 6 : 282. 1892.

Bulbous herbs. Leaves ensiform. Scape up to 60 cm long. Umbels 6 - 12 - flowered.

Yoksum, Sedonchen to Rongli ; 700 - 2100 m.

Fl. & Fr. : April - June (-Aug.).

C. powelii Baker is grown as an ornamental.

HYMENOCALLIS Salisb.

Hymenocallis littoralis Salisb. in Trans. Hort. Soc. London 1 : 338. 1812.

Herbs, up to 60 cm tall. Flowers white with a corona.

Namli ; ca 1000 m.

Fl. : July - Aug. Cult.

PANCRATIUM L.

Pancratium verecundum Aiton, Hort. Kew. 1 : 412. 1789 ; Hook. f., F.B.I. 6 : 285. 1892.

Bulbous herbs. Leaves thin, lanceolate. Scape longer or almost equalling leaves, 2 - 6 - flowered. Spathe lanceolate. Corolla lobes joined by corona.

Namli ; ca 500 m.

Fl. : Aug.

ZEPHYRANTHES Herb.

Zephyranthes candida Herb. in Bot. Mag. t. 2607. 1826.

Herbs. Leaves linear. Flowers large, solitary, funnel-shaped, white.

Gangtok, ca 1750 m.

Fl. : -Aug.

Z. carinata Herbert in Bot. Mag. 51 : t. 2594. 1825.

Herbs. Leaves several, flat. Flowers large, solitary, funnel-shaped, usually pink, borne on a leafless stalk.

Gangtok, Yoksum, Sambuk, Legship, Tendang, Namthang ; 600 - 2500 m.

Fl. & Fr. : April - July.

HYPOXIDACEAE

(B. Krishna)

CURCULIGO Gaertn.

Curculigo capitulata (Lour.) O. Kuntze, Rev. Gen. Pl. 2 : 703. 1891. *Leucoium capitulatum* Lour., Fl. Cochinch. 199. 1790. *Curculigo recurvata* Dryand in Ait., Hort. Kew. ed. 2 : 253. 1811 ; Hook. f., F.B.I. 6 : 278. 1892.

Stout herbs. Leaves variable, lanceolate. Flowers yellow.

Burtuk basti, Lower Sichey Basti, Saramsa.

Fl. & Fr. : May - July.

C. crassifolia (Baker) Hook. f., F.B.I. 6 : 279. 1892. *Molineria crassifolia* Baker in J. Linn. Soc. Bot. 17 : 121. 1879.

Stout herbs. Leaves elliptic-lanceolate. Flowers creamy yellow.

Silign Tzokupa (North Sikkim).

Fl. & Fr. : May - June.

C. gracilis (Kurz) Wall. ex Hook. f., F.B.I. 6 : 278. 1892. *Molineria gracilis* Kurz in Ann. Mus. Bot. Lugd. Bat. 4 : 177. 1867.

Stout herbs. Leaves lanceolate. Flowers lightyellow.

Samdong.

Fl. & Fr. : May - June.

C. orchiodies Gaert., Fruct. 1 : 63. t. 13. 1788 ; Hook. f., F.B.I. 6 : 279. 1892. *Curculigo brevifolia* Dryand in Ait., Hort. Kew. ed. 2. 2 : 253. 1811.

Small herbs. Leaves linear or linear-lanceolate. Flowers yellow or orange.

Teesta valley.

Fl. & Fr. : Aug. - Oct.

AGAVACEAE

(R.C. Srivastava)

AGAVE L.

A. sislana Persine in U.S. Senate Document 300 : 36. 1838. HATHIBAR (Nep.).

Robust shrubs. Leaves fleshy, linear-lanceolate, dark green, with tiny recurved spines at margins and a blackish brown, *ca* 2 mm long apical spine. Inflorescence 3 - 4 m tall.

Saramsha, on way to Pakyong, Turung, Rangpo, Singtam.

Fl. & Fr. : Feb. - April.

A. vera-cruz Miller., Dict. ed. n. 7. 1768. HATIBAR (Nep.).

Robust shrubs with large leaves. Inflorescence appearing periodically, 3 - 4 m tall. Flowers white.

Tarku, Namthi, Singtam, Yoksum, Jorethang ; up to 1500 m.

Fl. & Fr. : June.

Notes : Planted.

DRACAENA Vandelli *ex* L.

Dracaena angustifolia Roxb., Fl. Ind. 2 : 155. 1832 ; Hook. f., F.B.I. 6 : 327. 1892. MAKAI PATE (Nep.).

Erect, woody shrubs, up to 3 m tall, white. Leaves erect or patent, linear-lanceolate with broad sheathing base. Flowers pale yellow.

Saramsa ; up to 1400 m.

Fl. & Fr. : April - May.

FURCRAEA Vent.

Furcraea foetida (L.) Haw., Syn. Pl. Succ. 73. 1812. *Agave foetida* L., Sp. Pl. 323. 1753.

Large succulent shrubs. Leaves in rosettes of 30 - 40, lanceolate, up to 90 x 7.5 cm, coriaceous, more or less fleshy. Inflorescence up to 4.5 m tall. Flowers greenish, in loose panicles with drooping branches. Bulbils usually borne on flower clusters.

Jorethang (on the foothills of Salgiri) ; *ca* 550 m.

Fl. & Fr. : Sept. - Oct.

Notes : A variegated horticultural variety is also planted around Jorethang town.

YUCCA Dill. *ex* L.

Yucca aloifolia L., Sp. Pl. 319. 1753.

Large shrubs. Leaves hard, flattened, tipped with a needle like thorn. Flowering stems up to 3 m tall. Flowers white, hanging like bells from the erect flowering stem.

Gangtok (near Paljour Stadium) ; ca 1700 m. Cult.

DIOSCOREACEAE

(B. Mitra)

DIOSCOREA L.

Dioscorea alata L., Sp. Pl. 1035. 1753 ; Hook. f., F.B.I. 6 : 296. 1892. BOK (Lep.)

Twiners. Stem twining to the right, glabrous, prickly at the base. Leaves mostly opposite, broadly ovate ; basal nerves 5.

Cultivated throughout Sikkim ; up to 2500 m.

D. esculenta (Lour.) Burkill in Gard. Bull. Singap. 1 : 396. 1917 ; Prain & Burkill in Ann. Roy. Bot. Gard. Calc. 14 : 80. t. 35 - 36. 1936. *Oncus esculentus* Lour., Fl. Cochinch. 194. 1790. *Dioscorea sativa* L., Sp. Pl. 1033. 1753, p.p. ; Hook f., F.B.I. 6 : 295. 1892.

Twiners. Stem twining to the left, glabrous, purple with prickles. Leaves alternate, broadly cordate ; basal nerves 5 - 7. Capsules oblong, membranous.

Sikkim s.l. ; up to 912 m. cultivated.

D. glabra Roxb., Fl. Ind. ed. 2, 3 : 804. 1832.

Twiners. Stems twining to the right, glabrous, abundantly prickly at base. Leaves 7 - 9, broad oblong - lanceolate, rounded or cordate at base, basal nerves usually 7.

Sikkim s.l. ; up to 1216 m.

Fl. & Fr. : Aug. - Dec.

D. hamiltonii Hook. f., F.B.I. 6 : 295. 1892 ; Prain & Burkill in Ann. Roy. Bot. Gard. Calc. 14 : 301. t. 122. 1938. PUUMBOK ; (Lep.)

Twiners. Stems twining to the right, glabrous, not prickly at base. Leaves opposite, cordate or often subsagittate ; basal nerves 7 - 9. Capsules oblong.

Sikkim, s.l. ; ca 1200 m.

Fl. & Fr. : Aug. - Dec.

D. hispida Dennst., Schless. Hort. Malabr. 15. 1818. *Dioscorea deamona* Roxb., Fl. Ind. 3 : 805. 1832 ; Hook f., F.B.I. 6 : 289. 1892. RUKLOO (Lep.).

Twiners. Stems twining to the left, prickly. Leaves trifoliolate ; leaflets elliptic - oblong or obovate. Capsules pale brown, wings semi-elliptic ; bulbils absent.

Pakyong - Rhenock, S. of Peminagsti ; up to 2000 m.

Fl. & Fr. : July - April.

D. kamooneensis Kunth, Enum. Pl. 5 : 395. 1850 ; Hook. f., F.B.I. 6 : 290. 1892.

Twiners. Stems twining to the left, unarmed, dark brownish red, white hairy when young. Leaves alternate, 3 - 5 - foliolate ; leaflets 7 - 12 cm long, basal nerves 2 - 3. Capsules oblong, hairy till fully ripened.

Lachung valley, Chungthang, Rangpo valley ; 1200 - 1800 m.

Fl. & Fr. : Aug.

D. prazeri Prain & Burkhill in J. Asiat. Soc. Beng. 73. 2, suppl. : 2. 1904 & in Ann. Roy. Bot. Gard. Calc. 14 : 29. 202. t. 5. 1936. *Dioscorea sikkimensis* Prain & Burkhill in Rec. B.S.I. 4 : 77. 84. 134. 1910.

Twiners. Stems twining to the left, unarmed. Leaves alternate, ovate-lanceolate, basal nerves 7 - 9. Capsules orbicular.

Sikkim, s.l. ; ca 1700 m.

LILIACEAE

(R.C. Srivastava)

ALETRIS L.

(Sometimes treated under Melanthiaceae)

Aletris gracilis Rendle in J. Bot. 44 : 41. 1906.

Small herbs. Leaves linear, 7 - 10 in rosettes. Flowers pink ; pedicels glabrous.

Yakla, Latong, Maimaichu Lake ; 3000 - 3900 m.

Fl. & Fr. : Aug.

A. pauciflora (Klotzsch.) Hand - Mazz., Symb. Sin. 7 : 1220. 1936. *Stachyopogon pauciflorus* Klotzsch. in Klotzsch. & Garcke, Bot.

Reise Pr. Waldemar 49. t. 94. 1862. *A. nepalensis* Hook. f., F.B.I. 6 : 264. 1898, nom. illegit

Small, scapose herbs. Leaves linear, 5 - 10 in rosettes. Flowers pinkish white ; pedicels hairy.

Chola range, Lachen, Zemu valley, Dzongri, Changu ; 2500 - 4900 m.

Fl. & Fr. : June - Sept.

A. sikkimensis Hook. f., F.B.I. 6 : 265. 1892.

Herbs, much larger than *A. pauchiflora* with short narrowed leaves and leafy scape with 30 - 60 cm long raceme. Flowers greenish - white to orange-green.

Thangu, Yumthang, Karponang, Lachem, Zemu, Lilonakh ; 2400 - 4200 m.

Fl. & Fr. : July - Sept.

ALLIUM L.

A. cepa L., Sp. Pl. 300. 1753 ; Hook. f., F.B.I. 6 : 377. 1892. PYAJ (Nep., Hindi), ONION (Eng.).

Bulbous herbs. Leaves subdistichous, fistular. Flowers white.

Yumthang ; cultivated up to 1750 m.

Fl. & Fr. : April - July.

A. fasciculatum Rendel in J. Bot. 44 : 42. 1906. *A. gageanum* W.W. Smith *apud* W.W. Smith & Cave in Rec. BSI 4 : 248. 1911.

Herbs, up to 30 cm tall. Bulbs narrowly cylindric, 2 - 5 mm wide. Leaves linear, flat. Flowers white or greenish yellow.

Dongbong, Nathula, Kupup, Jelep-la, Thanka-la, upper valley of Llonakh ; 2400 - 4800 m.

Fl. & Fr. : Aug. - Oct.

Notes : Found in loose alluvial soil, by the side of streams.

A. hookeri Thw., Enum. Pl. Zeyl. 339. 1864 ; Hook. f., F.B.I. 6 : 341. 1892.

Herbs, up to 50 cm high. Bulbs narrowly cylindric. Leaves 4-5, linear, flat. Spathe deciduous. Umbels subglobose, many flowered, lax, up to 5 cm in diam. Flowers white.

Bitu (on way to Kabi from Gangtok) ; 1900 - 3050 m.

Fl. & Fr. : July - Aug. Cult.

A. macranthum Baker in J. Bot. 3 : 293. 1847 ; Hook. f., F.B.I. 6 : 345. 1892.

Small erect herbs with narrow bulbs. Leaves linear, gradually acuminate, keeled. Flowers purple.

Thangu, Lachen, Serbathang ; 3600 - 4200 m.

Fl. & Fr. : July - Aug.

A. prattii C.H. Wright *apud* Forbes Hemsley in J. Linn. Soc. Bot. 36 : 124. 1903. *A. victorialis* *auct. non* L. : Hook. f., F.B.I. 6 : 342. 1892, *p.p.* *A. victorialis* var. *angustifolium* Hook. f., *tom. cit.* 343. 1892. *A. prattii* var. *ellipticum*. Wang. & Tang. in Bull. Fam. Mem. Instl. Priol. Ser. 2 : 297. 1937. KOK - PA (Sikkimese).

Bulbous herbs. Leaves flat, elliptic. Flowers pinkish purple.

Changu, Kalep to Thangu, Zemu, Llonakh, Laghep, Bikbari, Giagong, Dzongri, Yumthang ; 2400 - 4500 m.

Fl. & Fr. : May - Oct.

A. prezealaskianum Regel., Allior. Monogr. 164. 1875. *A. jacquemontii* Regel. *Op. Cit.* 162. 1875 ; Hook. f., F.B.I. 6 : 342. 1892.

Herbs with linear leaves. Heads globose or subglobose. Flowers lilac.

Sikkim, *s.l.* ; 3900 - 4200 m.

Fl. & Fr. : May - Sept.

A. sativum L., Sp. Pl. 296. 1753 ; Hook. f., F.B.I. 6 : 37. 1892 ; LAHSUN (Nep.), GARLIC (Eng.).

Herbs with flat leaves. Scales slender. Sapthes long - beaked.

Cultivated up to 1400 m.

Fl. & Fr. : June - Sept.

A. sikkimense Baker in J. Bot. 12 : 292. 1874 ; Hook. f., F.B.I. 6 : 341. 1892.

Herbs with cylindric bulbs. Leaves linear. Flowers blue.

Kareng - Chholamo Kongralama, Thangu, Llonakh ; 3400 - 4800 m.

Fl. & Fr. : July - Sept.

A. wallichii Kunth, Enum. Pl. 4 : 443. 1834 ; Hook. f., F.B.I. 6 : 341. 1892.

Herbs with small clustered bulbs. Leaves 2, oblong-linear. Flowers purple-pink.

Chamnago, Gnathang, Lachung, Karponang, Dzongri, Laghep, Manamitso Lake, Changu, Lachen - Thangu ; 2400 - 4650 m.

Fl. & Fr. : June - Oct.

ALOE L.

Aloe vera Burm. f., Fl. Ind. 83. 1768. GHEWKUMARI (Nep.).

Herbs with fleshy leaves and orange flowers.

Gangtok, Tadong ; 1200 - 1600 m.

Fl. & Fr. : April - May. Planted.

Notes : Medicinal herb.

ASPARAGUS L.

Asparagus filicinus Buch. - Ham. *apud* D. Don, Prodr. Fl. Nepal. 49. 1825 ; Hook. f., F.B.I. 6 : 314. 1892.

Herbs profusely branched. Stems unarmed. Cladodes 6 - 8 mm. Pedicels slender.

Dongbong, Yoksum ; up to 1800 m.

Fl. & Fr. : May - July.

Asparagus racemosus Wills, Sp. Pl. 2 : 152. 1799 ; Hook. f., F.B.I. 6 : 316. 1892. SATAWARI, SATMULA (Nep.).

var. **racemosus**

Profusely branched climbing shrubs with recurved spines. Cladodes terete, curved. Flowers small, white.

Namli, Gangtok, North of Jorethang ; 600 - 1800 m.

Fl. & Fr. : April - Aug.

var. **subacerosa** Baker in J. Linn. Soc. Bot. 14 : 624. 1875 ; Hook. f., F.B.I. 6 : 315. 1892.

Climbing undershrubs. Cladodes stout. Racemes small, few-flowered. Rachis and pedicels axillary.

Sikkim, s.l. ; up to 1200 m.

Fl. & Fr. : April - June.

ASPHODELUS L.

Asphodelus tenuifolius Cav. in Anal. Cienc. Nat. 3 : 46. t. 27. 1801; Hook. f., F.B.I. 6 : 322. 1892.

Herbs. Leaves radical, linear. Flowers greenish white.

Turung, Sada ; up to 1200 m.

Fl. & Fr. : July - Oct.

CAMPYLANDRA Baker

Campylandra aurantiaca Baker in J. Linn. Soc. Bot. 14 : 582. t. 20. 1875. *Tupistra aurantiaca* (Baker) Wall. ex Hook. f., F.B.I. 6 : 325. 1892.

Herbs with long stout roots. Leaves flaccid, coriaceous, sessile. Flowers yellow, in spike.

Yoksum - Bakhim, Damthang to Temi, Rabong - la ; 1200 - 2450 m.

Fl. & Fr. : Oct. - May

CARDIOCRINUM Lindl.

Cardiocrinum giganteum (Wall.) Makino in Bot. Mag. Tokyo 27 : 125. 1913. *Lilium giganteum* Wall., Tent. Fl. Nepal. 21, t. 12 - 13. 1824, excl. syn. ; Hook. f., F.B.I. 6 : 349. 1892.

Herbs, up to 2 m tall. Leaves broadly cordate-ovate. Flowers purple-white, borne in racemes.

Lower Chakung Chu, Zemu, Lachung-Yumthang, Bakhim, Lachen, Thomba, Muguthang ; 1800 - 3300 m.

Fl. & Fr. : May - Aug.

CHLOROPHYTUM Ker-Gawler

Chlorophytum arundinaceum Baker in J. Linn. Soc. Bot. 15 : 323. 1876 ; Hook. f., F.B.I. 6 : 333. 1892. *C. breviscapum* auct. non Dalz : Hook. f., l.c. 333. 1892, p.p.

Herbs. Leaves oblanceolate. Scapes 15 - 50 cm long, stout. Flowers white.

Melli - Teesta ; 500 - 1200 m.

Fl. & Fr. : July - Aug.

C. breviscapum Dalz. in Hook., Kew J. Bot. 2 : 141. 1850 ; Hook. f., F.B.I. 6 : 353. 1892.

Herbs. Leaves linear-lanceolate, crispulate. Scapes stout, 7.5 - 15 cm, rarely branched, naked, shorter than leaves. Flowers in dense racemes.

Melli - Teesta ; ca 900 m.

Fl. : June - Aug.

C. khasianum Hook. f., F.B.I. 6 : 334. 1892. *C. napalense* Baker in J. Linn. Soc. Bot. 15 : 330. 1876, *p.p.*

Herbs. Leaves sessile, linear, flat. Flowers white, in dense-flowered racemes.

Sikkim, s.l. ; 1200 - 3200 m.

Fl. & Fr. : July - Sept.

C. nepalense (Lindl.) Baker in J. Linn. Soc. Bot. 15 : 330. 1876, *p.p.* *Phalanquium nepalense* Lindl. in Trans. Hort. Bot. London. 6 : 276. 1826. *Chlorophytum undulatum* Wall. ex Hook. f., F.B.I. 6 : 335. 1892, *nom. illegit.*

Herbs. Leaves linear. Scapes naked, 30 - 90 cm long, paniculately branched. Flowers white, drooping, solitary or in clusters.

Chungthang, Namchi, Yoksum, Reshi, Lachung, Lachen, Kabi ; 1400 - 2500 m.

Fl. & Fr. : July - Aug.

C. comosum (Thunb.) Jacques **variegatum** (Eng. - Spider plant) is cultivated at Legship, Yoksum, Tashiding localities and is characterised by its variegated leaves and arching scapes bearing plantlets.

CLINTONIA Rafin.

(Sometimes treated under Uvulariaceae)

Clintonia undensis Trautv. & Meyer var. **alpina** (Kunth ex Baker) Hara in J. Jap. Bot. 38 : 72. 1963. *C. alpina* Kunth ex

Baker in J. Linn. Soc. Bot. 14 : 585. 1875 ; Hook. f., F.B.I. 6 : 361. 1892.

Herbs, with creeping rootstocks. Leaves narrow costate. Flowers white.

Tsoka, Bakhim - Dzongri, Chola, Dikchu, Zemu valley, Lachen, Lachung, Kupup ; 2200 - 3600 m.

Fl. & Fr. : May - Aug.

CORDYLINE Comm. *ex A.L. Juss. nom. cons.*

Cordyline terminalis Kunth, Enum. Pl. 5 : 25. 1825 ; Hook. f., F.B.I. 6 : 331. 1892.

Gangtok ; ca 1700 m.

Fl. & Fr. : Aug. - Sept.

DIANELLA Lamk.

(Sometimes included in Phormiaceae)

Dianella ensifolia (L.) DC. in Redoute, Lil. 1 : t. 1. 1802 ; Hook. f., F.B.I. 6 : 337. 1892. *Dracaena ensifolia* L., Mant. Pl. 1 : 63. 1767.

Herbs, up to 1 m high. Leaves gradually tapering to acute at apex, midrib keeled. Scapes 30 - 60 cm long. Flowers white or yellowish.

Sikkim, s.l. ; up to 1800 m.

Fl. : April - July.

D. caerulea Sims.

Herbs up to 1 m high. Ultimate partial inflorescence spikelike, bearing more than 20 flowers on an elongate axis.

Cultivated at Namli ; up to 150 m.

Fl. : April - June.

DISPORUM Salisb. *ex D.Don*

(Sometimes treated under Uvulariaceae)

Disporum calcaratum D.Don in Trans. Linn. Soc. Lond. 18 : 516. 1841 ; Hook. f., F.B.I. 6 : 359. 1892. *Disporum wallichii* D.Don. *tom. cit.* 516. 1841.

Erect herbs, up to 70 cm tall. Leaves ovate-lanceolate. Umbels 4 - 6 - flowered. Flowers pale yellow ; tepal spurs 4 - 5 mm long.

Karponang ; 2100 - 2500 m.

Fl. & Fr. : May - Sept.

D. cantoniense var. **parviflorum** (Wall.) Hara, Fl. E. Himal. 407. 1966. *Uvularia parviflora* Wall. in Asiat. Res. 13 : 378. 1820.

Erect herbs, up to 1 m tall. Leaves ovate-lanceolate. Umbels 4 - 9 - flowered. Flowers pale violet with yellow anthers ; tepal spurs up to 1.5 cm long.

Yumcho La, Zemu, Llonakh, Latong, Singhik, Chungthang, Changu, Gnathang ; 2700 - 4800 m.

Fl. & Fr. : May - June.

D. leucanthum Hara in J. Jap Bot. 47 : 114. 1972.

Herbs, up to 40 cm high. Leaves ovate, acute to cuspidate. Flowers white.

Chung - thang ; 1530 - 2800 m.

Fl. : April - May.

FRITILLARIA L.

Fritillaria cirrhosa D. Don, Prodr. Fl. Nepal. 51. 1825 ; Hook.f., F.B.I. 6 : 353. 1892.

Herbs. Leaves scattered, lanceolate. Flowers greenish yellow, tessellated.

Changu, West of Tankra - la, Thangshing, Lingtoo, Dzongri, Bikbari, Alukthang ; 1500 - 4800 m.

Fl. & Fr. : April - Oct.

F. delavayii Franch. in J. Bot. Paris 12 : 222. 1898.

Herbs. Bulbs globose. Leaves 4 - 5, crowded at 2 - 3 cm above stem - base, sessile. Flowers yellowish green, with or without red spots.

Lachung, Tankra-la ; 3000 - 5000 m.

Fl. & Fr. July - Aug.

Notes : Roasted bulbs are eaten in Sikkim.

GAGEA Salisb.

Gagea lutea (L.) Ker - Gawl. in Curtis Bot. Mag. 30 t. 1200. 1809 ; Hook. f., F.B.I. 6 : 355. 1892. *Ornithogalum luteum* L., Sp. Pl. 306. 1753. *Gagea elegans* Wall. ex D. Don in Royle, Ill. Bot. Himal.

1. t. 75. f. 2. 1839. *G. lutea* auct. non Schultes f. : Hook.f., F.B.I. 6 : 35. 1892.

Small, bulbiferous herbs with lanceolate leaves and yellow flowers.

Zemu - chu, Thangu, Soonderdurya glacier, Lachen, Pentong ; up to 4300 m.

Fl. & Fr. : May - Aug.

GLORIOSA L.

Gloriosa superba L., Sp. Pl. 305. 1753 ; Hook.f., F.B.I. 6 : 358. 1892.

Climbing herbs. Leaves lanceolate, with cirrhose apices. Flowers bright yellow-red, large.

Sarmasa ; 400 - 2200 m.

Fl. & Fr. : Aug. - Sept. Probably grown.

HEMEROCALLIS L.

Hemerocallis fulva L., Sp. Pl. ed. 2, 1 : 462. 1762 ; Hook. f., F.B.I. 6 : 326. 1892.

Herbs with costate, linear leaves and large yellow, orange or red flowers.

Gangtok ; up to 3600 m.

Fl. & Fr. : May - Aug. Cultivated ; also as an escape.

IPHIGENIA Kunth

Iphigenia indica (L.) Kunth, Enum. Pl. 4 : 213 1843 ; Hook. f., F.B.I. 6 : 357. 1892. *Melanthium indicum* L., Mant. Pl. Alt. 226. 1771. *Anguillaria indica* (L.) R. Br., Prodr. Fl. Nov. Holl. 1 : 273. 1810.

Erect, bulbous herbs. Leaves few, scattered, linear. Flowers purple.

Sikkim, s.l. ; 1800 - 2000 m.

Fl. & Fr. : June - Sept.

KNIPHOFIA Moench.

Several species of *Kniphofia* are grown as ornamental of Yoksum, Rabong - la, Singtam.

LILIUM L.

Lilium nepalense D. Don in Mem. Werner. Nat. Hist. Soc. 3 : 412. 1821 ; Hook. f., F.B.I. 6 : 350. 1892. BHANGOO (Nep.).

Slender herbs, up to 1 m tall. Leaves lanceolate. Flowers drooping, sweet scented, orange-yellow, purple dotted.

Sikkim, s.l. ; 2300 - 3400 m.

Fl. & Fr. : June - Aug.

L. wallichianum J.A. & J.H. Schultes, Syst. Veg. 7 (2) : 1689. 1830 ; Hook.f., F.B.I. 6 : 349. 1892.

Bulbous herbs, up to 2 m tall. Leaves linear. Flowers 1 - 2, infundibuliform, white.

Singhik ; 1100 - 3000 m.

Fl. & Fr. : May - Sept.

Notes : Bulbs are eaten as vegetable by the local people.

Lilium lancifolium (*L. tigrinum* Ker - Gawler) is grown as an ornamental at Gangtok, Rumtek, near Tashiding and various other parts of the state.

LLOYDIA Salisb.

Lbydia delavayi Franch. in J. de. Bot. 12 : 193 & 195. 1898.

Herbs, up to 12 cm tall. Bulbs formed of needle like radical-leaf bases. Flowers solitary, terminal, yellow.

North of Dzongri, Momay Samdong Changu, Thangu, Jelep-la, Bikbari ; 4000 - 4500 m.

Fl & Fr. : May - June.

L.deliculata Noltie

Herbs, up to 5 cm high. Bulbs narrowly ovoid. Leaf solitary, basal, filiform, long than stems. Flowers white with purplish midrib and yellowish nectary.

Clangu - Kupup - Nathula, Lachen, Samiti ; 3650 - 4450 m.

Fl : June - July.

L flavenutans Hara in J. Jap. Bot. 49 : 202. 1974. *L. serotina* auct. no: Reich. ex Hook. f., F.B.I. 6 : 354. 1892, p.p.

Herbs, up to 24 cm tall. Bulbs narrowly ovoid with long neck, formed of 2 - 8 existing radical leaves. Flowers 1 or 2, nodding, yellow.

Jelep-la, Migothang, Mome Samdong, Gamothang, Thangshing-Lam Pokhri, Nathula, Changu - Kupup, Thangu, Muguthang, Llonak-la ; 3800 - 4700 m.

Fl. & Fr. : May - Sept.

C. longiscapa Hook. f., Icon. Pl. 9 : t. 834. 1852. *L. serotina* auct. non L. : Hook. f., F.B.I. 6 : 354. 1892, p.p.

Herbs up to 20 cm tall. Bulbs narrowly ovoid. Leaves filiform. Flowers white with violet veins.

Semthang, Yampung, Yumtso - la ; 4000 - 5000 m.

Fl. & Fr. : May - Aug.

L. serotina (L.) Reichb., Fl. German. Excur. 1 : 102. 1830 ; Hook. f., F.B.I. 6 : 354. 1892, p.p. *Bulbocodium serotinum* L., Sp. Pl. 294. 1753. *Lloydia himalensis* Royle, Ill. Bot. Himal. t. 93. f. 2. 1839. 338. 1840.

var. *serotina*

Small, bulbous herbs with 1 - 2 m, erect, campanulæ white flowers with violet veins.

Lachen, Thangu, Zemu, Llonakh valley ; 3500 - 5100m.

Fl. & Fr. : May - Aug.

var. *parva* (Marquand & Shaw) Hara, Fl. E. Himal. 2 : 16. 1971. *L. serotina* f. *parva* Marquand & Shaw in J. Linn. Soc. Bot. 48 : 228. 1919.

Herbs, smaller than var. *serotina*. Scape 2 - 6 cm. Perianth 6.5 - 8 mm ; filaments up to 3.5 mm, anthers more or less circular, ca 1 mm.

Muguthang, Lower Eastern Lungnak ; 4100 - 4700 m.

Fl. & Fr. : May - Aug.

var. *sikkimensis*

Changu, Nathula ; 3900 - 4200 m.

. **L. yunnanensis** Franch. *L. himalensis* sensu Dasgupta & Deb in Ind. J. For. 9 : 108. 1986, p.p. non Royle 1840.

Herbs, up to 10 cm high. Leafy bracts up to 6. Flower solitary, drooping, white, with reddish brown veins.

Changu, Bikbari, Dzongri, Namdee, Onglathang ; 3050 - 4400 m.

Fl. : May - July.

MAIANTHEMUM G.H. Weber ex Wiggers

Maianthemum fuscum (Wall.) La Frankie in Taxon 35 : 588. 1986. **Smilacina fusca** Wall., Pl. Asiat. Rar. 3 : 37. t. 257. 1832 ; Hook.f., F.B.I. 6 : 323. 1892. **Tovaria fusca** Baker in J. Linn. Soc. 14 : 568. 1875.

Herbs, up to 50 cm tall, slender. Leaves ovate-cordate, petioled. Panicles deltoid, 7 - 15 cm long. Perianth brown purple.

Bakhim, Thangu, Zemu ; 2400 - 3300 m.

Fl. & Fr. : May - June.

M. oleraceum (Baker) La Frankie in Taxon 35 : 589. 1986. **Tovaria oleracea** Baker in J. Linn. Soc. Bot. 14 : 569. 1875. **Smilacina oleracea** (Baker) Hook.f., F.B.I. 6 : 323. 1893.

var. **oleraceum**

Pendulous herbs. Flowers pink.

Lachen, Phodonchen, Sedonchen to Kabi, Lingtu, Tsoka to Yoksum, Zemu, Prag - Chu, Karponang ; 2500 - 4000 m.

Fl. & Fr. : May - June.

Notes : It is a favourite pot herb of Lepchas.

var. **acuminatum** (Wang & Tang) Karthik, l.c. 97. **Smilacina acuminata** (Wang & Tang) Hara in Hara *et al.*, Enum. Fl. Nepal. 1 : 78. 1978. **O. oleracea var. acuminata** Wang & Tang in Bull. Fam. Mem. Inst. Biol. 7 : 288. 1937.

Herbs, up to 1 m high. Leaves lanceolate, acuminate. Flowers deep wine-red to blackish. Inflorescence lax, axis zigzag ; pedicels long.

Tsoka to Jamlingthang, Lingtu ; ca 3200 m.

Fl. & Fr. : May - June.

M. oligophylla (Baker) Karthik. in Karthik. *et al.*, Fl. Ind. Enum. Monocot. 97. 1989. **Tovaria oligophylla** Baker in J. Linn. Soc. 14 : 565. 1875. **Smilacina oligophylla** (Baker) Hook. f., F.B.I. 6 : 323. 1892.

Perennial herbs, up to 30 cm tall, with an underground rhizome, oblong leaves and purple flowers.

Lachen, Gnathang to Sedonchen ; 3000 - 4000 m.

Fl. & Fr. : May - June.

M. pallidum (Royle) Karthik., l.c. 97. *Smilacina pallida* Royle, Illus. Bot. Himal. 1 : 380. 1839 ; Hook. f., F.B.I. 6 : 323. 1892. *Tovaria pallida* (Royle) Baker in J. Linn. Soc. 14 : 566. 1875.

Herbs 60 - 90 cm tall. Leaves 8 - 12, oblong, 10 - 15 cm long. Flowers white or purple.

Zemu ; 2400 - 3900 m.

Fl. & Fr. : May - July.

M. purpureum (Wall.) La Frankie in Taxon 35 : 589. 1986. *Smilacina purpurea* Wall., Pl. Asiat. Rar. 2 : 38, t. 144. 1831. *S. pallida* auct. non Royle : Hook.f., F.B.I. 6 : 323. 1892, p.p. *Tovaria purpurea* (Wall.) Baker in J. Linn. Soc. Bot. 14 : 566. 1875.

Stout herbs with creeping rootstocks. Leaves 8 - 12, oblong, up to 15 cm long, densely pubescent beneath. Flowers white.

Zemu, Lingtu, Yumthang, Dzongri Changu ; 2600 - 4200 m.

Fl. & Fr. : April - Aug.

NOMOCHARIS Franch.

Nomocharis flavidum (Rendle) Hara in Ohashi, Fl. E. Himal. 3 : 132. 1975. *Fritillaria flava* Rendle in J. Bot. 44 : 45. 1906.

Bulbous herbs. Flowers creamy-white, streaked with yellow or green.

Thangshing, Jamlingthang, Yatung, Bikbari, Samiti, Thangu ; ca 4500 m.

Fl. & Fr. : June - Sept.

N. nana (Klotz.) Wilson, Lilies E. Asia 13. 1925. *Lilium nanum* Klotz. in Bot. Ergeb. Veldem. Reise 53. 1862 ; Hook. f., F.B.I. 6 : 352. 1892.

Bulbous herbs. Leaves linear, blunt. Flowers small, drooping, campanulate, deep reddish purple.

West of Dzongri, Chamnags, Chemathang, Gnathong, Yumchhola, Zemu, Llonakh ; 3700 - 4800 m.

Fl. & Fr. : June - Sept.

NOTHOLIRION Wall.

Notholirion bulbiferum (Ling.) Stearn in Kew Bull. 1950. 421. 1950. *Paradisea bulbifera* Ling. in Limper. f. in Fedde Repert., Beih. 12 : 316. 1992.

Herbs, up to 1 m high. Leaves linear-lanceolate. Flowers lilac, mauve or pale purple.

Chamnago, Dikchu valley ; 3200 - 4100 m.

N. macrophyllum (D. Don) Boiss., Fl. Orient. 5 : 190. 1882. *Fritillaria macrophylla* D. Don, Prodr. Fl. Nepal. 51. 1825. *Lilium roseum* Wall. ex Hook. f., F.B.I. 6 : 352. 1892, p.p.

Small herbs, 30 - 50 cm tall. Leaflets oblong or linear.

Dikchu, Lachung, Tankra, Chungthang, Lachen ; 2100 - 5400 m.

Fl. & Fr. : May - Nov.

N. thomsonianum (Royle) Stapf. in Kew Bull. 1934 : 95. 1934. *Fritillaria thomsonianum* Royle, Illus. Bot. Himal. t. 92. f. 1. 1839. *Lilium roseum* Wall. ex Hook. f. in Curtis Bot. Mag. 79. t. 4725. 1853 ; Hook. f., F.B.I. 6 : 352. 1892, p.p.

Small, bulbous herbs. Flower pink to pale lilac.

Sikkim, s.l. ; 2700 - 3000 m.

OPHIOPOGON Ker - Gawler

(Sometimes included in Haemodoraceae)

Ophiopogon clarkei Hook.f., F.B.I. 6 : 268. 1892.

Scapigerous herbs. Stems prostrate, stout. Leaves lanceolate. Flowers white or pale lilac, with a terminal tuft of secundly curved linear leaves. Scape short, few-flowered.

Sikkim, s.l. ; 2400 - 2800 m.

O dracaenoides (Baker) Hook.f., F.B.I. 6 : 268. 1892. *Fluggea dracaenoides* Baker in Trimen, J. Bot. 174. 1874.

Scapigerous herbs. Stems prostrate, stout. Leaves lanceolate. Flowers white or pale lilac.

Sikkim, s.l. ; 1200 - 1800 m.

Fl. & Fr. : July - Sept.

O. intermedius D. Don, Prodr. Fl. Nep. 48. 1825 ; Hook. f., F.B.I. 6 : 269. 1892.

Short stemmed, scapigerous herbs. Leaves tufted, linear, toothed margined. Flowers lilac tinged white.

Yoksum Bakhim - Tsoka, Gangtok, Karponang, Thangu, Zemu, Changu, Lachen ; 1200 - 3000 m.

Fl. & Fr. : July - Sept.

O. wallichianus (Kunth) Hook. f., F.B.I. 6 : 269. 1892. *Fluggea wallichiana* Kunth, Enum. Pl. 5 : 303. 1850. *Ophiopogon intermedium* D. Don var. *parviflora* Hook. f., F.B.I. 6 : 269. 1892. *O. parviflorus* (Hook. f.) Hara in J. Jap. Bot. 40 : 21. 1965.

Scapigerous herbs. Leaves very narrow, secundly curved. Scape very slender, decurved. Flowers white or pale purplish.

Lachen, Phedang - Choka, Karponang, Burmiok, Kabi, Yoksum to Bakhim ; 2000 - 2800 m.

Fl. & Fr. : June - Aug.

PARIS L.

Paris polyphylla Smith in Rees. Cyclop. 26 : Paris n. 2. 1813 ; Hook.f., F.B.I. 6 : 362. 1892. SATUWA (Nep.).

ssp. *polyphylla* var. *polyphylla*

Erect, unbranched herbs. Leaves whorled, long pointed. Flowers yellow-green.

Penlong - La, Chunghang - Lachen, Fungong, Bakhim - Dzongri, Changu, Sada ; 1800 - 3300 m.

Fl. & Fr. : April - July.

ssp. *polyphylla* var. *appendiculata* Hara, Fl. E. Himal. 410. 1966.

Differs from var. *polyphylla* in having long apiculi to the anther connectives (3.5 mm or more).

Bikbari, Migothang - Nayathang, above Tsoka, Bakhim - Dzongri ; 2500 - 4000 m.

Fl. & Fr. : May - June.

ssp. *polyphylla* var. *wallichii* Hara in Ohashi, Fl. E. Himal. 3 : 134. f. 14b. 1975.

Differs in its leaves which are ovate-lanceolate, 3.5 - 6 cm wide, with truncate base and 1.2 - 4 cm long petioles.

Penlong La ; 1800 - 3000 m.

Fl. & Fr. : May - June.

var. *stenophylla* Franchet in Nouv. Arch. Mus. Paris 2, 10 : 97.
1888.

Differs in its leaves which are sessile, 10-15 in number and *ca* 12 mm wide.

Draigchu ; *ca* 2500 m.

Fl. & Fr. : July - Aug.

ssp. *marmorata* (Stearn) Hara in J. Fat. Sci. Univ. Tokyo Bot. 10 : 176. t. 2B-C, 6C. 1969. *Paris marmorata* Stearn in Bull. Br. Mus. Nat. Hist. (Bot.) 2 : 79. f. 11, t. 8. 1956. *P. polyphylla* var. *brachysteamon* Franchet in J. de. Bot. 12 : 191. 1898.

Herbs, similar to ssp. *polyphylla* but smaller in all parts.

Bakhim - Dzongri ; 2500 - 3000 m.

Fl. & Fr. : May - June.

PELIOSANTHES Andrews

Peliosanthes griffithii Baker in J. Linn. Soc. Bot. 17 : 506. 1879 ; Hook.f., F.B.I. 6 : 266. 1892.

Scapigerous herbs. Leaves long petioled, strongly 5-nerved. Scape very short. Racemes many-flowered. Flowers yellow.

Sikkim, *s.l.* ; 300 - 700 m.

Fl. & Fr. : Dec. - April.

P. macrophylla Wall. ex Baker in J. Linn. Soc. 17 : 505. 1879 ; Hook.f., F.B.I. 6 : 266. 1892. CHILLEDHOTISARA (Nep.).

Large, stout herbs. Leaves elliptic-lanceolate, 13 - 15 - nerved. Scape long or short. Racemes many - flowered. Flowers greenish or purple.

Sikkim, *s.l.* ; 300 - 700 m.

Fl. & Fr. : Nov. - Dec.

P. teta Andrews in Bot. Rep. 10 : t. 605. 1808.

Herbs differing from the above 2 species in having more than one flower per bract.

Sikkim, s.l.

POLYGONATUM Mill.

Polygonatum brevistylum Baker in J. Linn. Soc. Bot. 14 : 556. 1975 ; Hook. f., F.B.I. 5 : 319. 1892. *P. oppositifolium* var. *decipiens* Baker in F.B.I. 6 : 320. 1892.

Herbs. Stem grooved. Leaves usually all alternate, oblong, subcoriaceous. Peduncles 2 - 4 - flowered. Flowers yellow ; lobes green. Berries (fruits) ellipsoid.

Chola, Bakhim ; 2100 - 3000 m.

Fl. & Fr. : June - Sept.

P. cathcartii Baker in J. Linn. Soc. Bot. 14 : 559. 1875 ; Hook. f., F.B.I. 6 : 320. 1892.

Herbs. Stem tall, glabrous, terete. Leaves all opposite, subpetioled, ovate-oblong or lanceolate. Peduncle 2 - 4 - flowered. Flowers green or yellowish.

Bakhim to Tsoka, Lachen ; 2400 - 3200 m.

Fl. & Fr. : May - Aug.

P. cirrhifolium (Wall.) Royle, Ill. Bot. Himal. 1 : 380. 1839 ; Hook. f., F.B.I. 6 : 322. 1892. *Convallaria cirrhifolia* Wall. in Asiat. Res. 13 : 382. t. 1820.

Weak, flexuous, climbing herbs with whorled sessile leaves and white (tinged with purple) flowers.

Chungthang, Changu, Chola, Dentam, Pamianchi, Dzongri, Gamothang, Zemu, Lachen ; 1700 - 4400 m.

Fl. & Fr. : May - Sept.

P. hookeri Baker in J. Linn. Soc. (Bot.) 14 : 550. 1875 ; Hook. f., F.B.I. 6 : 320. 1892.

Dwarf herbs. Leaves opposite or lower ones alternate, crowded, sessile, linear. Flowers solitary, violet.

Yumthang, Samiti, Legep, Changu, Thangu, Thela, Llohnak, Lachen, Dzongri, above Thangsheng, Chaunrikiang ; 2900 - 5790 m.

Fl. : April - July.

P. kansuense Maximowicz ex Batalin.

P. curvistylum Hua in J. de. Bot. 6: 434. 1892.

Herbs 20 - 40 cm high. Leaves in whorls, 1 - 4 in each whorl, 2.9 - 10 mm wide, hooked when young. Flowers purple.

Dzongri, Sherabthang, Lamtung, Laghep, Bikbari ; 2700 - 4300 m.

Fl. May - July.

P. leptophyllum (D. Don) Royle.

Herbs. Leaves 3.3 - 12 mm wide, gradually tapering to very fine, acute apex. Peduncles 7 - 21 mm long. Pedicels 4 - 10 mm long. Berries up to 7 mm.

Lachen, Yampung, Zemu valley, Yakla, Matong, Tumbok; 2400 - 4250 m.

Fl. : June - July.

P. nervulosum Baker in J. Linn. Soc. 14 : 557. 1875; Hook.f., F.B.I. 6 : 319. 1892.

Herbs. Stems up to 55 cm high, zigzag above, leafless below. Scales oblong, whitish. Leaves alternate, lanceolate to elliptic, abruptly caudate. Flowers yellow.

Lachung ; 2100 - 3000 m.

Fl. : June - July.

P. oppositifolium (Wall.) Royle, Ill. Bot. Himal. 1 : 380. 1839 ; Hook.f., F.B.I. 6 : 320. 1892. *Convallaria oppositifolia* Wall. in Asiat. Res. 13 : 380. t. 1820. RUKKHIROLA (Sikkimese name).

Herbs, up to 1 m tall. Leaves oblong-lanceolate. Flowers many, axillary, light-pink.

Dentam ; *ca* 1500 m.

Fl. & *Fr.* : April - June.

P. punctatum Royle, Ill. Bot. Himal. 1 : 380. 1839 ; Hook.f., F.B.I. 6 : 321. 1892.

Epiphytic herbs with angled and grooved stems. Leaves oblong-lanceolate. Flowers white, spotted with liliac.

Joropokree, Chola, Latong, Lachen, Tendong Peak ; 2000 - 2600 m.

Fl. & Fr. : April - June.

P. Sibiricum F. Delaroche

Herbs. Leaves in whorls of 3 - 9, linear-lanceolate, coiled at tips.
Flowers purple tinged.

Laghep, Lachen, Lachung, Yumthang, Latong, Dzongri, Pemiongsti,
Chungthang, Yakthang, Jamlingthang to Bikbari, Thangshing to Lam-
pokri ; 1800 - 4200 m.

Fl. & Fr. : March - July.

P. singalilense Hara in J. Jap. Bot. 40 : 102. 1965.

Herbs. Stems up to 45 cm high. Leaves opposite or in groups of 3
upwards, lowest solitary, lanceolate, contracted below apices. Flowers
yellow-greenish-white.

Laghep, Thangsing to Lampokri Dzongri, Lingtoo, Jamlinghang to
Bikbari, Muguthang, Nayathang ; 3700 - 4400 m.

Fl. & Fr. : April - June.

P. verticillatum (L.) All., Fl. Pedemont. 1 : 131. 1785 ; Hook. f.,
F.B.I. 6 : 321. 1892. *Gonvallaria verticillata* L., Sp. Pl.. 315. 1753.

Erect herbs with sessile, whorled leaves. Flowers greenish white.

Changu, Thangu, Chola Range, Lachen, Jambinghang Zemu
valley ; 2700 - 4200 m.

Fl. & Fr. : May - Sept.

SANSEVIERIA Thunb.

Sansevieria cylindrica Boj. in Hort. Maurit. 349. 1839.

Perennial herbs. Leaves cylindrical, in a basal rosette, thick
cartilaginous, 90 - 100 cm long, dark green, often banded with paler lines,
acuminate. Scapes with inflorescence, shorter than leaves, racemose, up
to 30 cm long. Flowers ca 18 mm long, creamy white, tinged with
pink.

Jorethang, Rangpo ; up to 1500 m. Cult.

S. trifasciata Hort. ex Prain, Bengal Pl. 2 : 1054. 1903.

Perennial herbs. Leaves in a basal rosette, tufted, linear - lanceolate,
wide ensiform, flat. Flowers ca 12 mm long.

Jorethang, Gangtok, Mamli, Rangpo ; up to 1500 m. Cult.

STREPTOPUS Michaux.

Streptopus parasimplex Hara & Ohashi in J. Jap. Bot. 48 : 101.
f. 2 B. 1973.

Herbs, stoloniferous, up to 20 cm high, unbranched. Flowers white ; filaments equalling anthers.

Near Dzongri, Migothang to Nayathong ; 3100 - 4000 m.

Fl. & Fr. : June - July.

S. simplex D. Don, Prodr. Fl. Nepal. 48. 1825 ; Hook. f., F.B.I. 6 : 322. 1892. GARI - MUCHI (Nep.).

Herbs with creeping rootstock. Leaves amplexicaul, ovate-cordate. Flowers solitary or 2 - nate, axillary, white ; filaments shorter than anthers.

Chola, Changu, Phadomchen, Gnathang, Lachen to Naram village, Tsoka to Dzongri, Lingtu, Zemu ; 2400 - 4200 m.

Notes : Found among moss on rocks in shade. Eaten as vegetable.

THEROPOGON Maxim.

Theropogon pallidus (Kunth) Maxim. in Bull. Acad. Imp. Sci. St. Petersb. 15 : 90. 1870 ; Hook. f., F.B.I. 6 : 324. 1892. *Ophipogon pallidus* Wall. ex Kunth, Enum. Pl. 5 : 300. 1850. JUNGLI-PIAZ (Nep.).

Herbs with radical, recurved, glossily leaves. Flowers in terminal racemes, drooping, pink with white-tinge.

Lachung - Thanka La, Lachen, Zemu, Chungthang ; 1800 - 2700 m.

Fl. & Fr. : June - Sept.

TOFIELDIA Huds.

(Sometimes treated under Melanthiaceae)

Tofieldia himalaica Baker in J. Linn. Soc. Bot. 17 : 489. 1879 ; Hook. f., F.B.I. 6 : 358. 1812.

Herbs, with creeping rootstock. Leaves radical, equitant, narrowly linear. Flowers greenish white.

Dikchu, Leghep, Yumchoo La, Zemu, Lachung, Yakla, Changu ; 3200 - 3900 m. Occassional.

Fl. & Fr. : May - Sept.

TRICYRTIS Wall. *nom. cons.*

(Sometimes treated under Uvulariaceae)

Tricyrtis maculata (D. Don) J.F. Macbr. in Contr. Gray Herb., n.s. 3, 53 : 5. 1918. *Composa maculata* D. Don, Prodr. Fl. Nepal. 51. 1825. *Tricyrtis pilosa* Wall., Tent. Fl. Nep. 62, t. 46. 1826 ; Hook. f., F.B.I. 6 : 359. 1892.

Glandular-pubescent herbs with creeping rootstock. Leaves aplexicaul. Flowers white, spotted with purple.

Lachen, Lachung, Chungthang ; 1300 - 2500 m.

Fl. & Fr. : July - Aug.

TRILLIDIUM Kunth

Trillidium govanianum (D. Don) Kunth, Enum. Pl. 5 : 120. 1850. *Trillium govanianum* D. Don in Royle, Ill. Bot. Himal. 348. t. 93. f. 1. 1839 ; Hook. f., F.B.I. 6 : 361. 1892.

Herbs with radical leaves and drooping, purple flowers.

Gnathang, Lachen, Migothang to Nayathang Zemu valley ; 2700 - 4000 m.

Fl. & Fr. : July - Aug.

TRILLIUM L.

Trillium tschonoskii Maxim. in Bull. Acad. St. Petersb. 29 : 218. 1884 ; Hook. f., F.B.I. 6 : 361. 1892. *T. obovatum* Pursh. *sensu* Rao in Bull. Bot. Surv. India 5 : 165 - 205. 1964.

Perennial herbs with underground rhizome. Leaves sessile, broadly subrhomoboidally ovate or orbicular. Flowers light purple.

Lachen to Naram village, Rookah, Lachung, Lachen - Zemu valley ; 2700 - 4000 m.

Fl. & Fr. : July - Sept. Common at moist places.

TUPISTRÀ Ker - Gawl.

Tupistra clarkei Hook. f., F.B.I. 6 : 325. 1892.

Plants larger than *T. nutans*. Leaves 10 - 12.5 x 5 - 8 cm. Spikes decurved. Flowers pale greenish yellow or purplish ; mouth of perianth lobes not closed by stigma.

Sikkim, s.l. ; 600 - 1500 m.

T. nutans Wall. ex Lindl. in Edw., Bot. Reg. 15 : t. 1223. 1829 ; Hook. f., F.B.I. 6 : 324. 1892. NAKIMA (Nep.).

Herbs. Leaves radical, oblanceolate. Flowers dull brown-purple, borne in 7 - 15 cm long spikes, peduncles decurved ; mouth of perianth lobes closed by stigma.

Pemiangsti - Tingling bridge, on way to Phodong ; up to 1500 m.

Fl. & Fr. : March - April.

Notes : Inflorescence is eaten cooked.

YPSILANDRA Franch.

Ypsilandra yunnanensis W.W. Smith & Jeffrey var. **himalaica** Hara in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 80. 1978.

Herbs. Leaves in basal rosettes, oblanceolate. Flowers pale blue.

Sikkim, s.l. ; 4000 - 4300 m (*cf.* Hara *et al.* 1978)

SMILACACEAE

(R.C. Srivastava)

Smilax L.

Smilax aspera L., Sp. Pl. 1028. 1753 ; Hook. f., F.B.I. 6 : 306. 1892.

Large prickly climbing shrubs with small white fragrant flowers and blue - black berries.

Tingling ; 1200 - 2600 m.

Fl. & Fr. : Oct. - June.

S. aspericaulis Wall. ex DC., Monogr. Phan. 1 : 195. 1878 ; Hook. f., F.B.I. 6 : 306. 1892. *Smilax verruculosa* Merr. in Philipp. J. Sci. 5 : 173. 1910 & Enum. Fl. Pl. 1 : 209. 1922. *S. bracteata* Presl. var. *verruculosa* (Merr.) T. Koyama in Quart. J. Taiwan Mus. 10 : 18. 1957. DATUM (Nep.).

Climbing shrubs. Leaves oblong-lanceolate, cuspidate. Flowers creamy white.

Gangtok, Singhik to Mangan, Yoksum to Mintok - Khola ; 1200 - 2600 m.

Fl. & Fr. : Oct. - May.

S. elegans Wall. ex Kunth, Enum. Pl. 5 : 163. 1850. *S. galucophylla* Klotzsch in Klötzsch & Gareke, Bot. Reise Pr. Waldemar 45, t. 91. 1862. —

S. parviflora Wall. ex Hook.f., F.B.I. 6 : 304. 1892. *Smilax longibracteata* Hook. f. *tom. cit.* 305. 1892.

Slender, terete, smooth, unarmed herbs. Leaves ovate-cordate or lanceolate, 2.5 - 7.5 cm long, acuminate. Flowers minute, purple.

Lachen ; 1800 - 3100 m.

Fl. & Fr. : May - Aug.

S. ferox Wall. ex Kunth, Enum. Pl. 5 : 251. 1850 ; Hook.f., F.B.I. 6 : 307. 1892.

Shrubs up to 2 m tall. Leaves elliptic, ovate-obovate-oblong. Berries green-purple.

Sikkim, s.l.; 1100 - 2700 m.

Fl. & Fr. : April - June.

S. hajrae R.C. Srivastava, *nom. nov.* *S. rigida* Wall. ex Kunth, Enum. Pl. 5 : 164. 1850, *nom. illegit.*, *non* Solander in Russel, Nat. Hist. Aleppo (ed. 2) 27. 1794. Hook.f., F.B.I. 6 : 304. 1892.

Scandent shrubs. Leaves subsessile, orbicular-ovate or ovate-cordate. Flowers small pale-brown, in 1 - 4 - flowered umbels. Berries up to 12 mm in diam, black.

Gnathang to Phodonchen, Lachen, Bakhim - Dzongri, Yoksum. 2100 - 2900 m.

Fl. & Fr. : May - Oct.

S. lanceifolia Roxb., Fl. Ind. 3 : 792. 1832. Hook.f., F.B.I. 6 : 308. 1892.

Climbing shrubs. Leaves orbicular, oblong or lanceolate. Flowers ca 6 mm in diam.

Yoksum, Burmiok, Rishi - Rinchinpong, Rhenok ; 900 - 2100 m.

Fl. & Fr. : Nov. - March.

S. menispermoidea A. DC., Mongr. Phan. 1 : 108. 1878. *S. elegans* Wall. ex Hook.f., F.B.I. 6 : 305. 1892, *non* Wall. ex A.DC. 1878.

Slender, terete, small herbs. Leaves variable, ovate-cordate ; petiole sheaths 1.2 - 2.5 cm long. Umbels few-flowered. Berries ca 8 mm in diam., blue-black.

Lachen, Chungthang, Lachung, Dzongri, Yakla, Karponang ; 1800 - 3400 m.

Fl. & Fr. : May - Aug.

S. ocreata A.DC., Mongr. Phan. 1 : 193. 1878. *S. roxburghiana* Wall. ex Hook.f., F.B.I. 6 : 311. 1892, *p.p. excl. syn.* *S. orthoptera*.

Climbing shrubs. Stems more or less prickly, branches smooth. Leaves polymorphous, oblong, ovate or ovate-lanceolate. Umbels solitary or 2 - 3 on a common peduncle, many flowered. Berries *ca* 8 mm in diam.

Gangtok (Deorali) ; 1200 - 1700 m.

Fl. & Fr. : May - Nov.

S. osmastonii Wang & Tang in Bull. Fan. Mem. Inst. Biol. 7 : 298. 1937.

Shrubs. Leaves coriaceous, with densely reticulate veins on upper surface, papillose beneath.

Sikkim, *s.l.* ; 1500 - 2100 m.

S. ovalifolia Roxb. ex D. Don, Prodr. Fl. Nepal. 49. 1825. *S. macrophylla* Roxb., Fl. Ind. 3 : 793. 1892 ; Hook.f., F.B.I. 6 : 310. 1892. KUKUR - DYOTI (Nep.).

Stout, scandent, prickly shrubs. Leaves orbicular or broadly oblong. Umbels 2 - 3 on a short peduncle. Berries 2 - 3.7 cm in diam.

Melli - Teesta ; 300 - 1000 m.

Fl. & Fr. : May - July.

S. perfoliata Lour., Fl. Cochin. 622. 1790. *S. prolifera* Roxb., Fl. Ind. 3 : 795. 1832, *p.p.* ; Hook.f., F.B.I. 6 : 312. 1892. *S. laurifolia* auct. non Willd. : Roxb., Fl. Ind. 3 : 292. 1832.

Scandent, prickly herbs. Leaves ovate-lanceolate. Umbels solitary.

Yoksum, Gangtok ; 500 - 2000 m.

Fl. & Fr. : April.

S. wallichii Kunth, Enum. Pl. 5 : 246. 1850.

Twiners. Petiole-wings oblong to sub lanceolate, up to 4 mm wide. Peduncles stout and long. Umbels 35 - 80-flowered.

Mamring ; 1200 - 1500 m.

Fl. & Fr. : Feb. - May.

S. wightii A.D.C., Mongr. Phan. 11 : 174. 1878 ; Hook.f., F.B.I. 6 : 310. 1892.

Large climbing shrubs. Branches sparsely prickly. Leaves orbicular or broadly ovate or elliptic. Umbels solitary or 2 - 3 on a common peduncle, many flowered. Berries ca 12 mm in diam.

S. zeylanica L., Sp. Pl. 1029. 1753 ; Hook. f., F.B.I. 6 : 309. 1892.

Shrubs with slender and more or less 4 - angled branches. Leaves elliptic or broadly oblong or ovate- oblong. Umbels 20 - 30 - flowered. Berries ca 12 mm in diam.

Sikkim, s.l. ; up to 1500 m.

PONTEDERIACAE

(R.C. Srivastava)

Eichhornia crassipes (Mart.) Solms in A.DC., Mongr. Phan. 4 : 527. 1883. *Pontederia crassipes* Mart., Nov. Gen. Sp. 9, t. 1823.

Floating aquatic herbs. Petioles swollen, spongy. Flowers purple - white or bluish.

Gangtok, Phodong, Luing Busty area ; up to 1800 m.

Fl. & Fr. : Almost throughout the year.

MONOCHORIA C. Presl.

Monochoria vaginalis (Burm.f.,) C. Presl., Rel. Haenk 1 : 128. 1827 ; Hook. f., F.B.I. 6 : 363. 1892. *Pontederia vaginalis* Burm.f., Fl. Ind. 80. 1768.

Herbs. Rootstock short. Leaves linear to ovate - cordate. Flowers blue, sprinkled with red.

On way to Rumtek, Ranipool ; up to 1800 m.

Fl. & Fr. : Aug. - Sept.

In marshy places, rice fields.

XYRIDACEAE

(R.C. Srivastava)

XYRIS L.

Xyris indica L., Sp. Pl. 62. 1753 ; Hook. f., F.B.I. 6 : 365 . 1892.

Perennial, tufted herbs. Leaves 6 - 8 mm wide Scapes stout, deeply grooved. Spikes ovoid or globose ; bracts orbicular or cuneately obovate, dark reddish brown.

Sikkim (*cf.* Hooker 1892).

Fl. & Fr. : June - Aug.

X. pauciflora Willd., Phytogr. 1. 2, t. 1, f. 1. 1974 ; Hook. f., F.B.I. 6 : 365. 1892.

Perennial tufted herbs. Leaves narrowly linear, shorter than scape. Spikes globose or ovoid, bracts broadly oblong.

Sikkim, *s.l.*

Fl. & Fr. : June - Aug.

X. schoenoides Mart. in Wall., Pl. Asiat. Rar. 3 : 30. 1832 ; Hook. f., F. B.I. 6 : 365. 1892.

Perennial tufted herbs, up to 70 cm tall, with linear, acuminate or subacute leaves. Heads reddish brown ; corolla yellow.

Sikkim, *s.l.* ; up to 1800 - 2700 m.

Fl. & Fr. : June - Sept.

COMMELINACEAE

(N. R. Mandal)

AMISCHOPHACELUS Rolla Rao & Kammathy

Amischophacelus axillaris (L.) Rolla Rao & Kammathy in J. Linn. Soc. Bot. 59 : 306. 1966. *Commelina axillaris* L., Sp. Pl. 42. 1753. *Cyanotis axillaris* (L.) J.A. & J.H. Schultes, Syst. Veg. 7 (2) : 1154. 1830 ; Hook. f., F.B.I. 6 : 388. 1892.

Annual herbs with slender branches. Leaves sessile. Inflorescence axillary.

Sikkim, *s.l.* ; ca 500 m.

Fl. & Fr. : Aug. - Nov.

AMISCHOTOLYPE Hassk.

Amischotolype hookeri (Hassk.) Hara, Fl. E. Himal. 399. 1966. *Forrestia hookeri* Hassk. in Flora 47 : 629. 1864 ; Hook. f., F.B.I. 6 : 384. 1892.

Annual herb. Leaves petioled, elliptic, oblanceolate, acuminate. Flowers small. Capsules oblanceolate.

Sikkim, *s.l.* ; 500 - 800 m.

Fl. & Fr. : July - Sept.

A. mollissima (Bl.) Hassk. in Flora 46 : 392. 1864. *Campelia mollisima* Bl., Enum. Pl. Jav. 1 : 7. 1827. *Forrestia mollis* Hassk. in Flora 47 : 630. 1864. *F. glabrata* Hassk., l.c. *F. marginata* var. *rostrata* Clarke in DC., Mongr. Phan. 3 : 237. 1881. *F. mollissima* (Bl.) Kds., Exk. Fl. Jav. 1 : 282. 1911.

Stout annual herbs. Leaves elliptic to ovate, oblanceolate. Flowers 6 - 12, subsessile, in capitate cymes, white or pink.

Dikchu, Sakyong valley ; 915 - 1520 m.

Fl. & Fr. : July - Aug.

COMMELINA L.

Commelina appendiculata Clarke in Commel. & Syst. Beng. 113. 1874 ; Hook. f., F.B.I. 6 : 374. 1892.

Diffusely branched annual herb, up to 90 cm tall. Leaves sessile, linear-lanceolate. Cymes bifid. Petals blue or white.

Sikkim, s.l. : 1000 m.

Fl. & Fr. : July - Sept.

C. benghalensis L., Sp. Pl. 41. 1753 ; Hook. f., F.B.I. 6 : 370. 1892.

Scandent or creeping decumbent herbs. Leaves elliptic-ovate, obtuse. Flowers blue.

Sikkim, s.l. ; 900 - 1600 m.

Fl. & Fr. : July - Oct.

C. diffusa Burm. f., Fl. Ind. 18. t. 7. f. 2. 1768. *C. communis* Dalz. & Gibbs., Bombay Fl. 252. 1861. *Commelina nudiflora* sensu Hook. f., F.B.I. 6 : 369. 1892, non L. 1753.

Diffused, glabrous herbs, rooting at the lower nodes. Leaves sessile, lanceolate or ovate-lanceolate. Cymes 1 - 3 - flowered. Flowers pale blue or white.

Fl. & Fr. : July - Aug.

C. hasskarlii Clarke, Commel. & Syst. Beng. 13. t. 3. 1874 & in DC., Mongr. Phan. 3 : 157. 1881 ; Hook. f., F.B.I. 6 : 370. 1892.

Suberect annual herbs. Leaves sessile, narrowly lanceolate, subacute. Cymes bifid, 1 - 2 - flowered. Petals blue.

Burtuk Basti ; 1000 - 1500 m.

Fl. & Fr. : Aug. - Nov.

C. paludosa Bl., Enum. Pl. Jav. 1 : 2. 1827 ; Rolla Rao & Kammathy in J. Bombay Nat. Hist. Soc. 59 : 60. 1962. *C. obligua* Buch. ex D. Don, Prodr. Fl. Nepal. 45. 1825, non Vahl 1806 ; Hook. f., F.B.I. 6 : 372. 1892. *Commelina obligua* var. *viscida* Clarke in DC., Mongr. 3 : 178. 1881.

Annual herbs. Leaves very variable. Cymes usually simple. Petals blue.

Sikkim, s.l. ; ca 1800 m.

Fl. & Fr. : Aug. - Oct.

C. sikkimensis Clarke, Commel. & Syst. Beng. t. f. 1874. in DC., Mongr. Phan. 3 : 147. 1881 ; Hook. f., F.B.I. 6 : 369. 1892.

Succulent trailing herbs. Leaves lanceolate. Inflorescence a bifid cyme, 6 - 12 - flowered. Petals blue.

Gangtok ; ca 1700 - 1830 m.

Fl. & Fr. : Sept. - May.

C. suffruticosa Bl., Enum. Pl. Jav. 1 : 3. 1827 ; Hook. f., F.B.I. 6 : 374. 1892.

Annual herbs, with stout, branched, glabrous stem. Leaves lanceolate.

Sikkim, s.l. ; ca 1800 m.

Fl. & Fr. : Aug. - Nov.

CYANOTIS D. Don

Cyanotis cristata (L.) D. Don, Prodr. Fl. Nepal. 46. 1825 ; Hook f., F.B.I. 6 : 385. 1892. *Tradescantia cristata* L., Sp. Pl. 42. 1753.

Creeping, reddish purple herbs, up to 45 cm long. Leaves distant, sessile, ovate-oblong-lanceolate or obtuse. Inflorescence a scorpioid cyme. Petals purple.

Dikchu valley ; ca 915 m.

Fl. & Fr. : Sept. - Nov.

C. vaga (Lour.) J.A. & J.H. Schult. in R. & S., Syst. Veg. 7 : 1153. 1830. *Tradescantia vaga* Lour., Fl. Cochinch. 193. 1790. *Cyanotis barbata* D. Don, Prodr. Fl. Nepal. 46. 1825 ; Hook. f., F.B.I. 6 : 385. 1892.

Annual decumbent herbs, up to 50 cm long. Leaves sessile, oblong or linear-lanceolate, acute, falcate or straight. Flowers subsessile, small.

Sichey busty ; 915 - 1700 m.

Fl. & Fr. : July - Sept.

FLOSCOPA Lour.

Floscopia scandens Lour., Fl. Cochinch. 193. 1790 ; Hook. f., F.B.I. 6 : 390. 1892.

Annual herbs, erect or subscendent. Leaves subsessile, elliptic-lanceolate, cuneate. Inflorescence axillary or terminal. Petals white, lilac or pink.

East Sikkim Dist. ; 900 - 1800 m.

Fl. & Fr. : Aug. - Nov.

MURDANNIA Royle

Murdannia divergens (Clarke) Buck. in Engl. & Pantl., Pflanzenfam. ed. 2, 15 a, 173. 1930. *Aneilena herbaceum* var. *divergens* Clarke in J.Linn.Soc. 6 : 448. 1871 & Cyrt. Beng. t. 16. 1847 & DC., Mongr. Phan. 3 : 203. 1881 ; Hook. f., F.B.I. 6 : 376. 1892.

Erect annual herbs. Leaves sessile, ovate, acute, inflorescence panicled. Petals blue.

Sikkim, s.l. ; ca 600 - 1800 m.

Fl. & Fr. : July - Sept.

M. edulis (Stokes) Faden in Taxon 29 : 77. 1980. *Commelina edulis* Stokes, Bot. Materia Med. 1 : 184. 1812. *Aneilema scapiflorum* (Roxb.) Kostel, Alleg. Med. Pharm. Fl. 1 : 127. 1831 ; Hook. f., F.B.I. 6 : 375. 1892. *Commelina scapiflora* Roxb., Fl. Ind. 1 : 178. 1820.

Annual herbs. Radical leaves lanceolate ; caudine leaves reduced to sheaths. Flowers blue-mauve.

Sikkim, s.l. ; ca 600 - 1800 m.

Fl. & Fr. : May - July.

M. japonica (Thunb.) Faden in Taxon 26 : 142. 1977. *Commelina japonica* Thunb. in Trans. Linn. Soc. London 2 : 332. 1794. *Aneilema lineolatum* Kunth, Enum. Pl. 4 : 69. 1843 ; Hook. f., F.B.I. 6 : 376. 1892. *Commelina elata* Vahl, Enum. Pl. 2 : 178. 1806. *M. elata* (Vahl) Brueck in Eng. & Prantl., Pflanzenfam. ed. 2. 15 : 173. 1930.

Annual herbs. Stems stout. Leaves oblong, acuminate. Flowers blue.

Sikkim, s.l. ; ca 600 - 1800 m.

Fl. & Fr. : July - Sept.

M. nudiflora (L.) Brenan in Kew Bull. 1952. 189. 1952. *Commelina nudiflora* L., Sp. Pl. 41. 1753, p.p. *Tradescantia malabarica* L., Sp. Pl., ed. 2, 412. 1762. *Aneilema nudiflorum* (L.) R. Br., Prodr. 271. 1810 ; Hook. f., F.B.I. 6 : 278. 1892. *Aneilema compressa* Dalz. in Hook., Kew J. Bot. 3 : 318. 1851. *A. nudiflorum* var. *compressa* (Dalz.) Clarke in DC., Mongr. Phan. 3 : 211. 1831. *Murdannia malabarica* (L.) Brueck. in Engl. & Prantl., Pflanzenfam., ed. 2, 15 a : 173. 1930. *Aneilema malabaricum* (L.) Narayanswamy & Biswas in Indian For. Rec. (n.s.), 3 : 5. 1941. *M. malabarica* (L.) Brueck. var. *compressa* (Dalz.) Sant. & Jain in Ind. Forester 92 : 643. 1966.

Erect or decumbent herbs, 10 - 25 cm long. Leaves sessile, linear to oblanceolate. Petals purplish.

East Sikkim Distt. Siche Basti.

Fl. & Fr. : Sept. - Nov.

M. vaginata (L.) Brueck. in Engl. & Prantl., Pflanzenfam. ed. 2, 15 a : 173. 1930. *Commelina vaginata* L., Mant. Pl. 2 : 77. 1767. *Aneilema vaginatum* (L.) R. Br., Prodr. 271. 1810 ; Hook. f., F.B.I. 6 : 381. 1892.

Annual decumbent herbs. Leaves 9 - 20 x 0.5 - 1 cm, acute. Stamens 6, 2 fertile.

Gangtok ; 515 - 1800 m.

Fl. & Fr. : July - Sept.

POLLIA Thunb.

Pollia hasskarlii Rolla Rao in Notes Roy. Bot. Gard. Edinb. 25 : 188. 1964. *Pollia aclisia* Hassk. in Miq., Pl. Jungh. 1 : 145. 1852, *nom. illegit.* ; Hook. f., F.B.I. 6 : 367. 1892.

Annual, stout, erect, herbs. Leaves lanceolate or oblanceolate. Inflorescence an erect panicle. Petals white.

East Sikkim Distt. ; ca 915 m.

Fl. & Fr. : Sept. - Nov.

P. subumbellata Clarke in J. Linn. Soc. Bot. 11 : 451. 1870 ; Hook. f., F.B.I. 6 : 368. 1892. *Aclisia subumbellata* (Clarke) Brueck. in Engl. & Prantl., Pflanzenfam. 2, 15 a : 176. 1930.

Annual suberect herb, 20 - 40 cm tall. Leaves elliptic-lanceolate, caudate - acuminate. Petals white. Mature fruits deep glossy blue.

East Sikkim Distt. ; *ca* 915 m.

Fl. & Fr. : Sept. - Nov.

RHOPALEPHORA Hassk.

Rhopalephora scaberrima (Bl.) Faden in Phytologia 37 : 480. 1977. *Commelina scaberrima* Bl., Enum. Pl. Jav. 1 : 4 : 1827. *Aneilema scaberrimum* (Bl.) Kunth, Enum. Pl. 4 : 69. 1843 ; Hook.f., F.B.I. 6 : 382. 1892. *Dictyospermum scaberrimum* (Bl.) Morton ex Pagr. in Phytologia 29 : 338. 1976.

Erect annual herbs, 0.5 - 1 cm tall. Leaves lanceolate, acuminate.

East Sikkim Distt. ; *ca* 915 m.

Fl. & Fr. : Sept. - Nov.

STREPTOLIRION Edgew.

Streptolirion volubile Edgew. in Proc. Linn. Soc. London 1 : 254. 1854 ; Hook.f., F.B.I. 6 : 389. 1892. *Tradescantia cordifolia* Griff., Priv. J. 208. 1847, *non* Swartz 1788. *S. cordifolium* (Griff.) O. Ktze., Rev. Gen. Pl. 2 : 722. 1891.

An annual stem twiner. Leaves deeply bilobed. Flowers white.

Gangtok, Rabong - la ; 1000 - 2750 m.

Fl. & Fr. : Aug. - Oct.

JUNCACEAE

(N.R. Mandal)

Juncus L.

Juncus allioides Franch. in Nouv. Arch. Mus. Hist. Nat. Paris 10 : 99. 1887.

Tufted annual herbs. Stems slender, more or less leafy from base upwards. Leaves tubular, filiform or rarely more or less flat.

Sikkim, *s.l.*

Fl. & Fr. : July - Oct.

J. benghalensis Kunth, Enum. Pl. 3 : 360. 1841. *J. membranceus* *sensu* Hook f., F.B.I. 6 : 397. 1892, *p.p.* *non* Royle *ex* D. Don 1839. *J. sphenostemon* Buchen in Engl., Bot. Jahrb. 6 : 233, t. 3. 1885.

Perennial herbs, 12.5 - 50 cm tall. Stem stoloniferous. Leaves slender, channeled or flattened above or tubular; sheaths long, auricled. Flowers shortly pedicelled. Seeds tailed.

Sikkim, s.l.; 1820 - 3950 m.

Fl. & Fr. : July - Oct.

J. bhutanensis Satake in J. Jap. Bot. 43 : 382. f. 1 - 2. 1968.

Annual or perennial herbs. Leaves slender. Capsules elliptic or oblong, rostrate at apex, 4 x 1.5 mm.

J. brachystigma Samuels. in Hand. - Mazz., Symb. Sin. 7 : 1236. 1936.

Herbs. Leaves subbasal, blunt, auricles short. Capsules narrowly ellipsoid.

Sikkim s.l.

Fl. & Fr. : July - Oct.

J. bracteatus Buchen. in Engl., Bot. Jahrb. 6 : 220. 1885 ; Hook. f., F.B.I. 6 : 397. 1892.

Herbs. Stem very slender, tufted, 15 - 20 cm long, with basal soft sheaths. Leaves subcompressed, few, short, filiform, tubular. Flowers pale or dark. Capsule beaked, ovoid, 1-celled.

Sikkim, s.l.; 3040 - 4250 m.

Fl. & Fr. : July - Oct.

J. cephalostigma Samuels. in Hand.-Mazz., Symb. Sin. 7 : 1233. 1936.

Herbs, up to 20 cm high. Leaves linear, cylindric, nonseptate. Flowers pale brown.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. chrysocarpus Buchen. in Engl., Bot. Jahrb. 6 : 201. 1885 ; Hook. f., F.B.I. 6 : 394. 1892.

Perennial herbs with tufted, very slender stems. Leaves 1 - 2, filiform, equalling or exceeding the stem. Cyme a solitary, many-flowered, pale yellow globose head, 1.2 - 1.7 cm in diam. Flowers sessile.

Sikkim, s.l.; 3040 - 3650 m.

Fl. & Fr. : July - Oct.

J. clarkei Buchen. in Engl., Bot. Jahrb. 6 : 210. 1885 ; Hook. f., F.B.I. 6 : 400. 1892.

Stoloniferous perennial herbs with broad leaves longer than stem. Cymes 1-many flowered heads.

Sikkim, s.l. ; 3040 - 4050 m.

Fl. & Fr. : July-Oct.

J. concinnus D. Don, Prodr. Fl. Nepal. 44. 1825 ; Hook. f., F.B.I. 6 : 339. 1892. *Juncus elegans* Royle ex D. Don in Trans. Linn. Soc. Lond. 18 : 321. 1840, *pro syn.* *Juncus elegans* Samuels. in Hand - Mazz., Symb. Sin. 1232. 1936. *J. luteocarpus* Satake in J. Jap. Bot. 43 : 384. f. 3-4, 1968.

Herbs with tuberous rootstock. Leaves flat, channelled, few, variable, filiform. Sheaths membranous. Flowers subsessile, white or creamy yellow.

Sikkim, s.l. ; 2128 - 3950 m.

Fl. & Fr. : July - Oct.

J. grisenbachii Buchen. in Abhandl. Nat. Verein. Bremen 3 : 295. 1872 ; Hook. f., F.B.I. 6 : 394. 1892.

Stoloniferous perennial herbs. Leaves equalling stem, terete channelled, septate. Cymes with spreading branches of many-flowered pale yellow globose heads. Seeds very pale, fusiform, with lax testa and small tails.

Sikkim, s.l. ; 3040 - 4355 m.

Fl. & Fr. : June - Oct.

J. himalensis Klotz. & Garcke in Die Bot. Ehreb. Der. Reis. Prim. Waldem. 60, t. 97. 1892 ; Hook. f., F.B.I. 6 : 398. 1892.

Perennial herbs. Stem 20 - 50 cm, stout, rigid, leafy below. Cymes irregularly branched, 2 to more, dark brown, 3 - 8 flowered. Flowers shortly pedicelled, brown. Capsule beaked, dark brown. Seeds long tailed.

Sikkim, s.l. ; 218 - 4256 m.

Fl. & Fr. : July - Oct.

J. inflexus L. Sp. Pl. 326. 1753. *Juncus glaucus* Ehrh. ex Sibth., Fl. Oxon. 113. 1794 ; Hook. f., F.B.I. 6 : 393. 1892.

Var. **inflexus**

Perennial herbs with rigid, glaucous, deeply striate, interrupted pithed, dark green stems.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

var. leptocarpus (Buchen.) Sanjappa in Karthik. *et al.*, Fl. Indic. Enum. Monoc. 85. 1989. *Juncus leptocarpus* Buchen. in Nachr. Ges. Wiss. Gottingen 244. 1869. *Juncus glaucus* Ehrb. var. *leptocarpus* (Buchen.) Buchen. in Engl., Bot. Jahrb. 6 : 20. 1885.

Perennial herbs.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. kingii Rendle in J. Bot. 44 : 45. 1906.

Perennial glabrous herbs. Stem terete, stoloniferous. Leaf solitary, half to three fourth of stem.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. leschenaultii Gay in Laharpe, Monogr. Junc. 137. 1827 ; Karthik. *et al.*, l.c. 84. 1989. *Juncus prismatocarpus* sensu Hook. f., F.B.I. 6 : 395. 1892, p.p. R. Br. 1810. *J. latior* auct. non Satake 1936 : Karthik. *et al.*, l.c. 85.

Very variable herbs. Stems 45 - 61 mm long, leafy, not septate, shorter than stem. Flowers green or brown, sessile.

Sikkim, s.l.

Fl. & Fr. : April - July.

J. leucanthus Royle ex D. Don in Trans. Linn. Soc. Lond. 18 : 318. 1840 ; Hook. f., F.B.I. 6 : 397. 1892.

var. leucanthus

Perennial herbs with 15 - 23 cm long stem, slender, tufted at length hollow ; basal sheaths hard, brown. Cymes 0.6 - 0.8 cm in diam. 6 - 10-flowered, pale yellow. Flowers shortly pedicelled ; style exserted. Seeds shortly tailed.

Sikkim, s.l.

Fl. & Fr. : May - July.

var. alpinus Buchen., Mongr. Junc. 396. 1890.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. leucomelas Royle ex D. Don in Trans. Linn. Soc. Lond. 18 : 319. 1840 ; Hook. f., F.B.I. 6 : 397. 1892.

Herbs with nonstoloniferous, naked, hollow stems. Leaves 1 -2, short, filiform, tubular. Cymes 3 - 8 - flowered solitary head. Anthers exserted. Seeds long - tailed.

Sikkim ; 3650 - 4870 m.

Fl. & Fr. : June - Aug.

J. membranceus Royle ex D. Don in Trans. Linn. Soc. Lond. 18 : 320. 1840 ; Hook. f., F.B.I. 6 : 397. 1892.

Perennial herbs. Stems up to 50 cm high. Cymes many flowered. Flowers white. Capsules beaked.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. minimus Buchen. in Bot. Zeit. 25 : 145. 1867 ; Hook. f., F.B.I. 6 : 400. 1892.

Perennial herbs. Rootstocks short, stout. Stems up to 10.5 cm long, tufted or one leaved, grooved. Cymes of 1 - 2 dark chestnut coloured sessile heads, 3 - 6 -flowered ; stamens included. Seeds ca 2 mm long, testa white.

Sikkim, s.l. ; 4868 - 5470 m.

Fl. & Fr. : July - Oct.

J. minticola Steud., Syn. Pl. Glum. 2 : 301. 1855. *Juncus prismatocarpus* sensu Hook.f., F.B.I. 6 : 395. 1892, p.p. non R. Br. 1810.

Herbs. Rootstock very short. Stems 45 - 60 cm tall, decumbent and rooting at the nodes. Leaves 5 - 25 x 0.15 - 0.3 cm, shorter than stem, cymes erect. Flowers green or brown, sessile. Seeds very minute, apiculate.

Sikkim, s.l.

Fl. & Fr. : May - Aug.

J. ochraceus Buchen in Abh. Nat. Ver. Brem. 3 : 292. 1872 & Engl. Bot. Jahrb. 4 : 207. t. 2. 1855 ; Hook. f., F.B.I. 6 : 394. 1892.

Densely tufted perennial herbs, 20.3 - 30.5 cm high. Stems grooved, slender. Leaves shorter or equalling the stem, channelled, wiry. Cymes compound. Capsules *ca* 3 mm long, exserted, narrowly ovoid, triquetrous, cuspidate.

Sikkim, s.l. ; 1520 - 2740 m.

Fl. & Fr. : July - Sept.

J. pseudocastaneus (Lingelsh.) Samuels. in Hand. - Mazz., Symb. Sin. 7 : 1230. 1936. *Juncus sikkimensis* var. *pseudocastaneus* Lingelsh. *apud* Limpr. f. in Fedde Repert. 12 : 316. 1922.

Herbs.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. sikkimensis Hook. f., F.B.I. 6 : 399. 1892 *et* Ic. Pl. 23 : t. 2235.

Perennial herbs. Stem stout, clothed at base with red-brown rigid sheaths. Cymes of two dark brown, unilateral, sessile, 4 - 6-flowered heads. Flowers sessile. Seeds with short thick white tails.

Sikkim, s.l. ; 3364 - 4256 m.

Fl. & Fr. : Aug. - Oct.

J. sphacelatus Decne. in Jacquem, Voy 4, Bot. 172. t. 172. 1835 ; Hook. f., F.B.I. 6 : 398. 1892.

Perennial herbs, 20 - 50 cm tall. Stems tufted, smooth. Leaves involute, obtuse or acute, unitubular, obscurely septate. Cymes of 2 - 5 -flowered heads. Capsules dark brown.

Sikkim, s.l. ; 3648 - 4864 m.

Fl. & Fr. : July - Sept.

J. thomsonii Buchen. in Bot. Zeit. 25 : 148. 1867 & in Engl., Pflriech - IV. 36. Ht. 25 : 24. 1906 ; W.W. Smith in Rec. B.S.I. 4 : 421. 1913. *Juncus leucomelas* sensu Hook. f., F.B.I. 6 : 397. 1892, *p.p. non* Royle *apud* D. Don 1840.

Perennial herbs. Stems 5 - 15 cm long. Cymes solitary. Anthers shorter than filaments. Capsules one celled.

Sherabthang ; *ca* 3900 m (*cf.* Smith 1913).

J. trichophyllus W.W. Smith in Rec. B.S.I. 6 : 103. 1914.

Annual herbs with very slender stem, tufted, 8 - 10 cm high, basal sheaths poorly developed, chestnut brown in colour. Cymes solitary, 1 - 2-flowered. Capsules obovoid-oblong, cuspidately beaked.

Sikkim, s.l.

Fl. & Fr. : July - Sept.

J. triglumis L., Sp. Pl. 328. 1753 ; Hook. f., F.B.I. 6 : 396. 1892.

Slender stiff naked stemmed herbs. Stems 5 - 15 cm long. Cymes terminal 3 - 4-flowered, dark brown.

Sikkim, s.l.

Fl. & Fr. : July - Oct.

J. uniflorus W.W. Smith in Rec. B.S.I. 6(4) : 104. 1914 ; Karthik. et al., l.c. 86. 1989.

Annual herbs, 2 - 3 (-10) cm high, densely caespitose. Stems filiform, clothed at base with few sheaths. Leaves solitary or few. Flowers solitary, terminal, blackish brown ; stamens purplish slightly longer than style. Capsules equalling the sepals, more or less beaked, dark brown, shining, 3-septate. Seeds few.

Sikkim, s.l. ; 3202 - 4256 m.

Fl. & Fr. : June - Sept.

J. wallichianus Laharpe, Monogr. Junc. 139. 1829 ; Karthik. et al., l.c. 86.

Herbs. Leaves tubular, septate. Capsules short, up to 4 mm.

Gangtok ; 1100 - 3100 m.

Fl. : May - Aug.

J. yoskisukei Goel in J. Econ. Tax. Bot. 7 : 208. 1985. *Juncus albescens* Clarke in Univ. Mus. Univ. Tokyo Bull. 2 : 161. 1971, non Fernald. 1924.

Caespitose, glabrous rushes. Stems 5 - 16 cm high. Heads single, 3 - 10-flowered. Flowers white ; stamens 6, 3-alternipetalous, outer 3 antipetalous.

Sikkim, s.l. ; 3000 - 4300 m.

Fl. & Fr. : June - July.

LUZULA DC.

Luzula campestris (L.) DC., Fl. Franc. 3 : 161. 1805 ; Hook. f., F.B.I. 6 : 401. 1892, p.p. ; Karthik. et al., Fl. Indic. Enum. Monoc. 86. 1989. *Juncus campestris* L., Sp. Pl. 329. 1753.

var. **campestris**

Perennial herbs, 15.2 - 45.7 cm high. Flowers sessile in peduncled capitate clusters.

Sikkim, s.l.

Fl. & Fr. : July - Sept.

L. chinensis N.E. Br. in J. Linn. Soc. 36 : 161. 1903 ; Mitra, Fl. Pl. E. India 2 : 93. 1985. *Luzula parivflora* Buchen., Mongr. Junc. 463. 1890, non Desv. 1808.

Herbs.

Sikkim, s.l. ; 3040 - 3344 m.

Fl. & Fr. : June - Sept.

L. effusa Buchen. in Krit. Ver. Aller. Junca. 53 : 88. 1880 ; Hook. f., F.B.I. 6 : 401. 1892.

Perennial herbs, up to 61 cm high. Leaves linear, glabrous or sparsely ciliate below. Flowers solitary, pale or dark brown. Capsules oblong, longer than the sepals, obtuse, mucronate.

Sikkim, s.l. ; 2736 - 3040 m.

Fl. & Fr. : July - Sept.

L. multiflora (Retz.) Lejeune, Fl. Env. Spa. 1 : 169. 1811. *Juncus multiflorus* Retz., Fl. Scand. Prodr. ed. 2. 82. 1795. *L. sudetica* DC., Fl. France. 6 : 306 1815 ; Kartik. et al., l.c. 87.

Herbs, up to 20 cm high. Leaves narrowly lanceolate. Flowers sessile. Capsules ellipsoid, trigonous.

Sikkim, s.l.

Fl. & Fr. : May - Sept.

L. obligantha Samuels. in Hulten. Fl. Kamt. 1 : 227. 1927.

Herbs, up to 20 cm high. Leaves long, narrow, Flowers small. Capsules equalling perianth.

Lachen ; 3000 - 3600 m.

Fl. & Fr. : May - Aug.

L. plumosa E. Meyer in Linnaea 22 : 387. 1849 ; Hook. f., F.B.I. 6 : 401. 1892.

Perennnial herbs, up to 45.5 cm high. Leaves linear, acuminate, glabrous or sparsely ciliate with very long hairs. Cymes subumbellate. Capsule oblong-obtuse or apiculate, length as sepals.

Chungthang ; 2736 - 3040 m.

Fl. & Fr. : July - Sept.

L. rostrata Muchen. ssp. **darjeelingensis** Satake in Hara, Fl. E. Himal. 404. 1966.

Pernnial herbs, caespitose, 25 - 35 cm high. Radical leaves linear, sparsely ciliate ; cauline leaves 2, much shorter and narrower. Inflorescence many flowered. Seeds ellipsoid.

Sikkim, s.l.

Fl. & Fr. : May - June.

Perennial herbs, 5 - 25.5 cm high. Leaves densely fascicled, channeled. Bracts brown with broad, white, membranous, ciliate margins and tips. Stamens 6. Capsules oblong, obtuse, shorter than sepals.

Sikkim, s.l. ; 3648 - 4409 m.

Fl. & Fr. : July - Sept.

ARECACEAE

(R.C. Srivastava)

ARECA L.

Areca catechu L., Sp. Pl. 1189. 1753. SUPARI (Nep.), BETEL NUT (Eng.).

A slender graceful palm, 12-24 m tall, with much branched spadix and ovoid orange fruits.

Majhitar - Singtam; ca 400 m. Planted.

Fl. & Fr. : April - June.

CALAMUS L.

Calamus acanthospathus Griff. in Cal. J. Nat. Hist. 5 : 39. 1844; Becc. & Hook. f., F.B.I. 6: 48. 1892.

Perennial climbers. Leaves 60 - 90 cm long; leaflets few, elliptic-lanceolate. Spadices stout, flagelliferous. Fruits obovoid to globose, shortly beaked, brown.

Sikkim (?); 400 - 1800 m.

Fl. & Fr. : March - Oct.

C. erectus Roxb., Fl. Ind. 3 : 774. 1832 : Becc. & Hook.f., F.B.I. 6 : 438. 1892.

Shrubs with erect, stout stem and remotely ciliolate margins of leaf segments. Spadices branched, elongate, ending in a flagellum. Fruits ovoid-oblong, *ca* 2.5 cm long.

Sikkim, *s.l.* : up to 1200 m.

Fl. & Fr. : Dec. - Jan.

C. flagellum Griff. in Mart., Hist. Nat. Palm. 333. t. 176. f. 9. 1849; Becc. & Hook. f., F.B.I. 6 : 439. 1892.

Shrubs. Leaves 1.8 - 2 m long ; leaflets subopposite, linear-lanceolate. Spadix 2.5 - 3 m. Fruits ovoid-oblong, cuspidately beaked, pale, with a dark band in lower half.

Sikkim *s.l.* ; up to 4000 m.

C. latifolius Roxb., Fl. Ind. ed 2, 3 : 775. 1832; Becc. & Hook.f., F.B.I. 6 : 455. 1892. KORAK - BET (Nep.).

Large climbers. Leaves 3 - 4.5 m long ; leaflets few, large, elliptic-lanceolate. Spadix decomound. Fruits subglobose, pale.

Sikkim, *s.l.* ; *ca* 600 m.

Fl. & Fr. : April - Oct.

C. leptospadix Griff. in Calc. J. Nat. Hist. 5 : 49. 1844; Becc. & Hook. f., F.B.I. 6 : 441. 1892.

Shrubs with very long spinous sheath terminating with a coiled flagellum. Flowers small. Fruits small, globose, mucronate.

Sikkim, *s.l.* ; up to 1200 m.

Fl. & Fr. : April - Nov.

CARYOTA L.

Caryota urens L., Fl. Zeyl. 187. 1757; Becc. & Hook. f., F.B.I. 6 : 422. 1892. BANSUPARI (Nep.).

Trunk, up to 10 m high, annulate. Leaflets cuniform, obliquely truncate, acutely serrate-toothed. Fruits reddish.

Namprikdong (in the forest along river) ; up to 800 m.

Fl. : Feb.

COCCUS L.

Coccus nucifera L., Sp. Pl. 1188. 1753; Hook. f., F.B.I. 6 : 482. 1893. NARIYAL (Nep.).

A large palm, up to 12 m high. Leaves up to 3 m long; leaflets up to 1 m long, coriaceous, flaccid ; petioles stout. Fruit a drupe, 12 - 30 cm long, obovoid, greenish or yellowish.

Majhitar - Singtam, Jorethang ; *ca* 400 m. Planted.

Fl. & Fr. : March - June.

DAEMONOROPS Bl.

Daemonorops jenkinsiana. (Griff.) Mart., Hist. Nat. Palm. 327. 1850; Becc. & Hook. f., F.B.I. 6 : 462. 1893. *Calamus jenkinsianus* Griff. in Calc. J. Nat. Hist. 5 : 81. 1850.

Stout, spiny climbers. Leaves pinnate, upper serrate. Fruits subspherical, shortly apiculate.

Sikkim, *s.l.*

Fl. & Fr. : Nov. - Feb.

PHOENIX L.

Phoenix acaulis Roxb. ex Buch.-Ham. in Trans. Linn. Soc. London 15 : 87. 1826; Becc. & Hook. f., F.B.I. 6 : 426. 1892.

Tall palms. Leaves 1 - 2 m long, subglaucous; lower leaflets spiny; upper ones scattered and interruptedly fascicled, linear. Fruits ovoid, *ca* 1.5 cm, orange-red.

Sikkim, *s.l.* ; up to 1600 m.

Fl. & Fr. : April - Nov.

P. lourierii Kunth, Enum. Pl. 3 : 257. 1841. **P. humilis** Royle ex Becc. & Hook. f., F.B.I. 6 : 426. 1892.

Plants with short tufted, rarely elongate stem. Leaves subglaucous ; leaflets interruptedly scattered. Fruits oblong with broad groove.

Jorethang : ca 500 m.

Fl. & Fr. : May - June.

P. rupicola T. Anders. in J. Linn. Soc. Bot 11 : 13. 1870 : Becc. & Hook. f., F.B.I. 6 : 425. 1892.

Tall palms ; trunk naked. Leaves up to 3 m long ; leaflets ca 45 cm long, bifarious (not fascicled). Fruits oblong.

Sikkim, s.l. ; up to 450 m.

Fl. & Fr. : Sept. - Nov.

P. sylvestris (L.) Roxb., Fl. Ind. 3 : 787. 1832 ; Becc. & Hook. f., F.B.I. 6 : 425. 1892. *Elate sylvestris* L., Sp. Pl. 1189. 1753.

Tall palms, 7.5 - 12 m high, clothed with persistent bases of petioles. Leaves 3 - 4.5 m long, glabrous; leaflets 30 - 60 cm long, fascicled. Fruits orange-yellow.

Jorethang ; ca 500 m.

PINANGA Bl.

Pinnaga gracilis (Roxb.) Bl., Rumphia 2 : 77. 1839 ; Becc. & Hook. f., F.B.I. 6 : 407. 1892. *Areca gracilis* Roxb., Fl. Ind. ed. 2, 3 : 619. 1832.

Stems slender, gregarious, 6 - 8 mm in diam., up to 6m tall, erect, thickened upwards. Leaves sparingly pinnate ; petiole scurfy. Spadix, white or scarlet, reflexed ; fruiting spadix pendulous. Fruits ellipsoidly fusiform, scarlet or orange.

Sikkim, s.l. : up to 900 m.

Fl. & Fr. : Nov. - April.

PLECTOCOMIA Mart. & Bl.

Plectocomia himalayana Griff. in Calc. J. Nat. Hist. 5 : 97. 1854 ; Becc. & Hook. f., F.B.I. 6 : 478. 1892.

Climbing shrubs. Stems ca 2.5 cm in diam. Leaves 1.8 - 2.4 cm long ; leaflets concolourous, tips filiform. Spadix and spikelets scurfy tomentose. Fruits depressed globose, ca 12 mm in diam.

Sikkim, s.l. ; 1200 - 2100 m (*cf. Mitra 1958*).

TRCHYCARPUS Wendl.

Trachycarpus martianus (Wall.) Wendl. in Bull. Soc. Bot. France 8 : 429. 1862 ; Becc. & Hook.f., F.B.I. 6 : 436. 1892. *Chamaerops martiana* Wall., Asiat. Rar. 3 : 5, t. 211. 1832.

Tall palms, 6 - 15 m, trunk naked for most part except beneath the crown. Leaves 1.2 - 1.5 m in diam., cut about half way down into linear 2 - lobed segments, margin denticulate. Flowers yellow. Fruits (drupes) bluish.

Gangtok ; up to 1650 m. Planted (?).

WALLICHIA Roxb.

Wallichia densiflora Mart., Hist. Palm. 3 : 190. 1838 ; Becc. & Hook. f., F.B.I. 6 : 419. 1892. *W. oblongifolia* Griff. in Calc. J. Nat. Hist. 5 : 486. 1844.

Apparently stemless or short-stemmed palm. Leaves 30 - 60 cm long ; leaflets oblong or linear-oblong, sinuately lobed, acutely toothed, white beneath.

Teesta valley, Singtam (on hill-slopes) ; up to 1400 m.

Fl. & Fr. : Not seen.

W. disticha T. Anders. in J. Linn. Soc. Bot. 11 : 6. 1969 ; Becc. & Hook. f., F.B.I. 6 : 419. 1892.

A tall palm. Stems 3 - 4.5 m high, 12.5 - 15 cm in diam., naked, annualate. Leaves 1.8 - 3 m, alternate, erect ; leaflets fascicled, linear, narrowed to base, praemorse denticulate.

Sikkim Himalaya (*cf.* Mitra 1958 ; Hook. f. 1892).

PANDANACEAE

(B. Mitra)

PANDANUS Parkinson

Pandanus nepalensis St. John in Bot. Mag. Tokyo 85 : 254, f. 7 - 8. 1972. *P. furcatus* var. *indica* Kurz in J. Bot. 5 : 102. 1867, p.p. ; Burkhill in Rec. B.S.I. 4 : 135. 1910. *P. furcatus* auct. non Roxb. : Hook. f., F.B.I. 6 : 484. 1893, p.p. BORKUNG, TARIKA (Nep.).

Medium sized trees. Leaves up to 1 m long, up to 4 cm wide, keeled, spiny along margins and keel. Fruits cone like drupes, orange when ripe.

Gangtok, Singtam, Rabangla, Gylshing, Sankhalang, Jorethang ; up to 1752 m. Common.

Fl. : Aug. - Sept. ; *Fr.* : Almost throughout the year.

Uses : Young leaves from the upper part of the stem are employed as a native medicine for antidote to snake poison. Leaves are also used for making mats and for thatching purpose.

P. sikkimensis St. John, recorded from Teesta, also refers probably to the same common species found almost throughout the State from *ca* 400 - 1500 m (- 1752 m), **Pandus unguifer** Hook. f. (syn. *P. minor* Hook.f.) is not seen in Sikkim. It is restricted to Mungpoo district of West Bengal only (R.C.S.)

ACORACEAE

(R.C. Srivastava)

ACORUS L.

- **Acorus calamus** L., Sp. Pl. 324. 1753 ; Hook. f., F.B.I. 6 : 555. 1893.
BOJO (Nep.).

Herbs. Leaves with prominent midrib. Berries redish.

Yoksum, Dikchu, near Rumtek, Phodong ; 1700 - 2300 m.

Fl. & *Fr.* : Aug. - Oct.

A. gramineus Aiton, Hort. Kew. 1 : 474. 1789 ; Hook. f., F.B.I. 6 : 556. 1893

Differs from *A. calamus* in having a very slender rhizome (*ca* 0.5 cm in diam. ; leaves without prominent midrib and longer spadix.

Sikkim, *s.l.*

Fl. & *Fr.* : Aug. - Oct.

ARACEAE

(R.C. Srivastava)

ALOCASIA (Schott) D. Don *nom. cons.*

Alocasia fallax Schott in Bonplandia 7 : 28. 1859 ; Hook. f., F.B.I. 6 : 527. 1893.

Herbs. Leaves broadly ovate or orbicular -ovate, sagittate. Spathes 12.5 - 15 cm long, yellow.

Rangpo, near Jorethang, Phensang ; 600 - 1200 m.

Fl. & *Fr.* : Aug.

A. macrorrhizos (L.) G. Don in R. Sweet, Hort. Brit. ed. 3, 631. 1839 ; Hook f., F.B.I. 6 : 526. 1893. *Arum macrorhizon* L., Sp. Pl. 965. 1753.

Herbs with massive stems. Leaves ovate, up to 125 x 75 cm, basal lobes rounded, deeply cordate, glossy, not peltate.

Rangpo - Teesta, Gangtok ; 600 - 1800 m.

A. navicularis (Koch. & Bouche) Koch. & Bouche in Index Sem. Hort. Berol. 1855. App. 2. 1855 ; Hook f., F.B.I. 6 : 527. 1893. *Colocasia navicularis* Koch. & Bouche in Index Sem. Hort. Berol. 1853 ; 13. 1853.

Herbs. Leaves broadly oblong, sagittate. Spathe 10-12.5 cm long. Spathe tube ovate-globose. Spadix ca 7.5 cm long, appendage equal to or longer than the inflorescence.

Jorethang ; 450 m.

Fl. & Fr. : July - Aug.

A. odora (Roxb.) Koch. & Bouche in Index Sem. Hort. Berol. App. 5. 1855. *Arum odorum* Roxb., Fl. Ind. 3 : 499. 1832.

Herbs. Leaves ovate, cordate. Spathe tube ovoid. Spadix white.

Melli-Teesta (on way to Peshok) ; ca 400 m.

Fl. : Aug.

AMORPHOPHALLUS Bl. ex Decne. nom. cons.

Amorphophallus bulbifer (Schott) Bl., Rumphia 1 : 148. 1837; Hook.f., F.B.I. 6 : 515. 1893. *Arum bulbiferum* Roxb. *apud* Sims in Curtis Bot. Mag. 46. t. 2072. 1819. *Pythonium bulbiferum* Schott in Schott & Endl., Melet. Bot. 18. 1832.

Tuberous herbs, up to 60 cm tall. Leaf solitary, many lobed, 30 - 45 cm in diam., ultimately bulbiferous at base of petiolules ; leaflets ovate or lanceolate. Spathe erect, pale pink or yellowish, appendage white or flesh-coloured.

Rangpo - Teesta, N. of Jorethang, Central Pandim - Singtam Road ; 300 - 1500 m.

Fl. & Fr. : May - Nov.

Notes : Tubers edible.

A. napalensis Bogner & Mayo, Avoideava 8(1) : 9. 1985.

Tuberous herbs. Leaves 30 - 45 cm (or more) in diam. ; leaflets ovate or oblong-lanceolate. Appendage clothed below with depressed neuters and all over above with tubercles.

Yoksum, Legship, Rinchung pong - Pemiongsti ; 700 - 1800 m.

Fl. & Fr. : May - Dec.

ARIOPSIS Nimmo

Ariopsis peltata Nimmo in Graham. Cat. Pl. Bombay & Civin. (Add. & Corr.) 252. 1839 ; Hook. f., F.B.I. 6 : 519. 1893.

Perennial herbs. Tubers small, clustered. Leaves ovate, 2.5 - 15 cm in diam. Spathe *ca* 2.5 cm long, nodding, violet with a green dorsal ridge, paler inside. Spadix decurved ; male flowers dark purple ; female flowers greenish.

Dikchu to Gangtok ; 900 - 2100 m.

Fl. & Fr. : May - June.

ARISAEMA Mart.

(R.C. Srivastava & P. Basu)

Arisaema concinnum Schott in Bonplandia 7 : 27. 1859 ; Hook. f., F.B.I. 6 : 505. 1893.

Herbs with globose tubers. Leaf solitary; leaflets 7 - 11, subsessile, narrowly lanceolate or oblanceolate. Spathe green on purple, striped with white.

Lachen, Kabi, Pemiongsti, above Dentam, Yoksum, Chungthang, Gnagtok ; up to 1800 m.

Fl. & Fr. : May - June.

A. consanguineum Schott, l.c. 27 ; Hook. f., l.c. 505

Herbs with depressed globose tubers. Leaf solitary ; leaflets 10 - 20, usually 15, narrowly linear-lanceolate. Spathe green. Peduncle mottled red-brown.

Chungthang ; 1500 - 2900 m.

Fl. & Fr. : July - Oct.

A. costatum (Wall.) Mart. ex Schott in Melet. Bot. 1 : 17. 1832 ; Hook. f., l.c. 501. *Arum costatum* Wall., Tent. Fl. Nep. 28 : t. 19. 1824. GURBO (Nep.).

Herbs with depressed tubers. Leaf solitary, leaflets dark green. Spatha reddish purple with striped with white.

Changu ; 2000 - 3000 m.

Fl. : May - June.

A. echinatum (Wall.) Schott in Melet. 1 : 7. 1832 ; Hook. f., l.c. 506. *Arum echinatum* Wall., Pl. Asiat. Rar. 2 : 30. 1831.

Herbs with small globose tubers. Leaf solitary ; leaflets 7 - 9, narrowly lanceolate. Spatha greenish, striped with white and brown or purple within ; tip of spadix appendage echinate.

Lachung ; 2700 - 3300 m. (*cf.* Hook. f. 1893).

Fl. : June.

A. erubescens (Wall.) Schott, l.c. 17 ; Hook. f., l.c. 506. *Arum erubescens* Wall., Pl. Asiat. Rar. 2 : 30. t. 135. 1831.

Herbs. Leaf solitary ; leaflets 9 - 12, narrowly lanceolate. Peduncle rose-coloured, streaked with red at base. Spatha white, striped with reddish purple.

Sikkim, s.l. ; 1800 - 2400 m.

Fl. : May - June.

A. flavum (Forssk.) Schott in Prodr. 40. 1860 ; Hook. f., l.c. 503. *Arum flavum* Forssk., Fl. Aeg. Arab. 157. 1775. MANE (Nep.).

Herbs. Leaves 2 ; leaflets 9 - 11, oblong or lanceolate. Spatha 2.5 - 5 cm long, yellow.

Sikkim, s.l. ; 1700 - 3600 m.

Fl. & Fr. : May - Aug.

A. galeatum Schott in Syn. Aroid. 26. 1856 ; Hook. f., F.B.I. 6 : 499. 1893.

Herbs. Leaf solitary ; leaflets ovate or cordate-acuminate, crenulate. Spatha green, striped with white.

Sikkim, s.l. ; 1800 - 2100 m.

Fl. : May - June.

A. griffithii Schott in Syn. Aroid. 26 : 1856 ; Hook. f., l.c. 499. *A. hookerianum* Schott in Oester. Bot. Wochenbl. 7 : 334. 1857. *Arisaema griffithii* Schott var. *verrucosum* (Schott) Hara, Fl. E. Himal. 2 : 340. f. 6

B. 1971. *Arisaema verrucosum* Schott in Oester. Bot. Woehenbl. 7 : 341. 1857 ; Hook. f., l.c. GURBO (Nep.).

Herbs. Leaves 2, trifoliolate ; leaflets rhomboid-orbicular. Spathe very large, dark purple, limb much broader than the tube, deeply 2 - lobed.

Lachen, Yumthang, Yakla, Dentam, Phedang to Bakhim, Tsoka, Jalook, Karponang - Changu - Thegu ; 2300 - 3900 m.

Fl. & Fr. : April - Aug.

Notes : Verrucosity on petioles and peduncles are quite variable with intermediate forms and hence can not be considered as a stable character for separating species.

A. intermedium Bl. in Rumphia 1 : 102 . 1836 ; Hook. f., l.c. 500.

var. intermedium

Herbs. Leaves 1 or 2 ; leaflets dimidiate-ovate or cordate, median elliptic. Limb of spathe lanceolate, incurved.

Sikkim, s.l. ; 2400 - 2700 m. '

Fl. : July.

var. biflagellatum (Hara) Hara, Fl. E. Himal. 2 : 332. 1971. *Arisaema biflagellatum* Hara in J. Jap. Bot. 36 : 77. 1961.

Herbs. Spathe-blade yellowish green, with a longer tail, 8 - 16 cm long. Appendage greenish.

Sikkim, s.l. ; 2400 - 2700 m.

Fl. : June.

A. jacquemontii Bl. in Rumphia 1 : 95. 1835 ; Hook. f., l.c. 505. *A. exile* Schott in Bonplandia 7 : 26. 1859 ; Hook. f., F.B.I. 6 : 506. 1893.

Herbs. Leaves 1 or 2 ; leaflets 5 - 7, elliptic or oblong-lanceolate. Spathe-limb oblong or ovate-lanceolate, striped with white, incurved and narrowed into a long drooping tail.

Changu, Bikbari, Thangsing, Dzongri, Legship, Lachung, Lachen; ca 4000 m.

Fl. : July - Aug.

A. nepenthoides (Wall.) Mart. in Flora 458. 1831 ; Hook. f., l.c. 504. *Arum nepenthoides* Wall., Tent. Fl. Nep. 26. t. 18. 1824. GURBU (Nep.).

Herbs ; Leaves 2 ; leaflets 5, elliptic or oblanceolate. Spathe-limb elliptic-ovate, base dilated into 2 broad rounded recurved auricles.

Chungthang to Lachen, Lachung, Dantam, below Bakhim, Damthang, Tendong ; up to 2700 m.

Fl. & Fr. : April - Oct.

A. ochraceum Schott in Bonplandia 7 : 27. 1959 ; Hook f., F.B.I. 6 : 507. 1813.

Herbs. Leaves radiatisect. Spathe-limb yellowish with brown stripes.

Sikkim, s.l. ; ca 3000 m.

A propinquum Schott in Oester. Bot. Wochenbl. 7 : 333. 1857 ; Hook f., l.c. 501. *A. sikkimensis* Stapf ex Chatterjee in Bull. Bot. Soc. Bengal 3 : 18. 1949. *A. wallichianum* var. *sikkimense* (Stapf. ex Chatterjee) Hara in J. Bot. 48 : 21. 1965. *A. costatum* var. *sikkimense* (Stapf.). Hara in J. Jap. Bot. 36 : 76. 1961. *A. ostiolatum* Hara in J. Jap Bot. 36 : 75. f. 1. 1961.

Herbs. Tubers subdepressed, ca 2 cm in diam. Leaf solitary, trisect ; segments ovate-elliptic. Spathe suberect.

Below Zalep la, Phedang, Changu, Jamling thang, Prek-chu-Thanshing, Dzongri ; 3500 - 4200 m.

Fl. : May - June.

A. speciosum (Wall.) Mart. in Flora 14 : 458. 1831 ; Hook.f., F.B.I. 6 : 500. 1893. *Arum speciosum* Wall., Tent. Fl. Nepal. 29, t. 20. 1824. MONGJING (Nep.).

var. **speciosum**

Herbs. Leaf solitary, trifoliate; leaflets elliptic or ovate, acuminate at apex, distinctly petiolulate, green with red margins. Spathe dark black-purple.

Lachen, Yoksum, Rate-chu, Gangtok ; 1750 - 3000 m.

Fl. & Fr. : March - July.

var. **mirabile** (Schott) Engl. in DC., Monogr. Phan. 2 : 540. 1879. *Arisaema mirabile* Schott in Oester. Bot. Wochenbl. 7 : 366. 1857.

Spadix appendage in the lower part conspicuously thickened, fusiform, curved, distinctly rugose-exasperate, sometimes forming small reticula.

Bakhim - Yoksum, Tendong Peak ; 2100 - 3100 m.

Fl. : June - July.

A. tortuosum (Wall.) Schott, Melet. 1 : 17. 1832. *Arum tortuosum* Wall., Pl. Asiat. Rar. 2 : 10, t. 114. 1830.

Herbs, up to 1.5 m tall. Leaves (1-) 2 (-3), pedately 5 - 17-foliate, leaflets rhombic-ovate, oblong to lanceolate, abruptly acuminate at apex, cuneate at base. Spathe green, sometimes dark purple.

Temi - Damthang, near Singhik, Nampok, Pemiangsti, Gangtok, Saramsa ; 1400 - 1800 m.

Fl. : April - June.

A. utile Hook. in Schott, Prodr. Aroid 30 : Bot. Mag. t. 6474.

Herbs. Leaves solitary ; leaflets sessile, rhombic-ovate or obovate, pale green with undulate, golden margins. Spathe-limb less than 10 cm broad, red-brown, coarsely reticulate.

Karponang - Changu, Lachen, Gamrothang ; 2700 - 3900 m.

Fl. & *Fr.* : May - June.

Notes : Plants are smaller than *A. griffithii* but similar in habit.

COLOCASIA Schott *nom. cons.*

Colocasia affinis Schott in Bonplandia 7 : 28. 1859 ; Hook. f., F.B.I. 6 : 523. 1893.

Herbs with globose tubers. Leaves 1 - 3, peltate, ovate-rounded, glaucous beneath. Spathe 6 - 10 cm long.

Legship - Tendong, Silak ; up to 2000 m.

Fl. & *Fr.* : June - Nov.

C. esculenta (L.) Schott in Schott & Endl., Melet. Bot. 18. 1832. *Arum esculentum* L., Sp. Pl. 965. 1753. *Arum colocasia* L., Sp. Pl. 965. 1753. *Colocasia antiquorum* Schott in Schott & Endl., Melet. Bot. 18. 1832 ; Hook. f., F.B.I. 6 : 523. 1893.

Herbs with hard, often tuberous rhizomes. Leaves peltate, ovate or subtriangular. Spathe more than 15 cm long.

Legship - Tendony, Jorethang, Yoksum ; 300 - 1200 m.

Fl. & *Fr.* : June - Nov.

C. fallax Schott in Bonplandia 7 : 28. 1859 ; Hook.f., F.B.I. 6 : 524. 1893.

Herbs with tuberous rhizomes. Leaves orbicular-ovate, 10 - 24 x 7.5 cm. Spathe much shorter than the limb.

Sikkim, s.l. ; 400 - 2000 m.

Fl. & Fr. : May - Oct.

LASIA Lour.

Lasia spinosa (L.) Thwaites, Enum. Pl. Zeyl. 336. 1864. *Draconitum spinosum* L., Sp. Pl. 967. 1753. *Pothos lasia* Roxb., Fl. Ind. 1 : 458. 1820. *Pothos heterophyllus* Roxb., Fl. Ind. 1 : 457. 1820. *Lasia heterophylla* (Roxb.) Schott in Schott & Endl., Melet. Bot. 21. 1832. Hook.f., F.B.I. 6 : 550. 1892.

Prickly plants with hastate-sagittate, deeply pedately pinnatifid leaves. Spathe very long, thick, twisted, dull purplish brown, up to 60 cm is diam.

Sikkim, s.l. ; up to 500 m.

Fl. & Fr. : Jan. - Nov.

POTHOS L.

Pothos cathcartii Schott, Aroid. 1 : 22, t. 44 - 45. 1857 ; Hook.f., F.B.I. 6 : 552. 1893.

Root-climbers with coriaceous bracts. Spathe *ca* 2.5 cm broad. Berries obovoid or oblong, 12 - 18 mm long, scarlet.

Dikchu - Singhik, Toong, Rabongla - Temi, Bhusuk ; 300 - 1500 m.

Fl. & Fr. : March - Dec.

Notes : On tree trunks.

P. roxburghii Schott, Aroid. 22. t. 37. 1858 ; Prodr. 565. 1860.

Root - climbers. Leaves obovate or lanceolate. Peduncles 4 - 6 mm long. Sapdx large, erect, inclined or deflexed.

Sikkim Himalaya ; 300 - 1200 m (*cf.* Hook. f. 1893).

REMUSATIA Schott

Remusatia hookeriana Schott in Oester. Bot. Wochenbl. 7 : 133. 1857 ; Hook.f., F.B.I. 6 : 522. 1893.

Herbs with much branched bulbiferous shoots. Leaves *ca* 10 - 15 cm long.

Dantam - Pamianchi, Jorethang, Namchi, Yoksum - Bakhim, Phamtom, Lachen, Lachung, Gangtok, Penlong - la ; 1500 - 2400 m.

Fl. & Fr. : April - Nov.

R. pumila (D. Don) H. Li & A. Hary, Acta Bot. Yunn., Suppl. 5. 28. 1992. *Gonathanthus pumilus* (D. Don.) Engler & Krause in Engler, Pflzenreich IV - 23E. Ht. 71 : 19, t. 5a - m. 1920 *Caladium pumilum* D. Don, Prodr. Fl. Nepal. 21. 1825. *G. sarmentosus* Klotzsch. & Otto in Link & Klotzsch., Icon. Pl. Rar. Hort. Roy. Bot. Berol. 1 : 33. t. 14. 1841. Hook. f., F.B.I. 6 : 522. 1893.

Terrestrial or epiphytic herbs. Leaves oblong-ovate, green below, yellow above, deep purple on sides. Spathe 12 - 20 cm long.

Toong -Chungthang, Namchi - Jorethang, Namthang, Yumthang, Sedonchen - Rongli, Kibek, Gangtok ; 1200 - 2400 m.

Fl. & Fr. : May - Nov.

Abundant on moist shady slopes.

RHAPHIDOPHORA Hassk.

Rhaphidophora callophyllum Schott in Bonplandia 5 : 45. 1857.

R. lancifolia Schott in Bonplandia 5 : 45. 1857 ; Hook.f., F.B.I. 6 : 545. 1893.

Climbers. Leaves falcately-lanceolate or ovate-lanceolate, 15 - 25 x 3.7 - 7.5 cm, acuminate or caudate, bases unequal sided; petioles 10 - 17.5 cm long, channelled to the knee. Spathe red inside.

Sikkim, s.l. ; 900 - 1500 m.

Fl. & Fr. : Jan.

R. decursiva (Roxb.) Schott in Bonplandia 5 : 45. 1857 ; Hook.f., F.B.I. 6 : 546. 1893. *Pothos decursivus* Roxb., Fl. Ind. 1 : 456. 1820. *R. exima* Schott in Bonpland. 5 : 45. 1857 ; Hook.f., F.B.I. 6 : 587. 1893.

Huge climbers. Leaves oblong, with 5 - 15 unequal lobes. Spathe oblong-obovate, beaked, cream-yellow.

Deorali, Temi, Yoksum - Bakhim. 1300 - 2000 m.

Fl. & Fr. : July - Nov.

R. glauca (Wall.) Schott in Bonplandia 5 : 45. 1875 ; Hook. f., F.B.I. 6 : 547. 1893. *Pothos glaucus* Wall., Pl. Asiat. Rar 2 : 45 t. 156. 1832.

Climbers. Leaves oblique with 3 - 5 unequal lobes. Spathe oblong or oblong-lanceolate, pale yellow.

Yoksum, W. of Singtam, Deorali, Kabi ; 400 - 1800 m.

Fl. & Fr. : July - Nov.

R. grandis Schott in Oester. Bot. Wochenbl. 349. 1858.

Climbers. Leaves up to 1 x 0.5 m ; leaflet pairs greenish beneath; petioles grooved above. Spathe orange-yellow inside and outside.

Rathong - Chu valley, above Yoksum, W. of Singtam, Gangtok - Kabi ; 600 - 2100 m

Fl. & Fr. : May - Feb.

R. hookeri Schott in Bonplandia 5 : 45. 1857 ; Hook. f., F.B.I. 6 : 540. 1893.

Climbers. Leaves oblong, cuspidately caudate-acuminate, base unequal sided. Spathe green, yellow inside.

Below Rumtek Monastery, Singtam - Gangtok ; up to 1250 m.

Fl. & Fr. : March - June.

SAUROMATUM Schott

Sauromatum brevipes (Hook.f.) N.E. Brown in Gard. Chorn. Ser. 34 : 93. 1903. *Typhonium brevipes* Hook. f., F.B.I. 6 : 511. 1893.

Tuberous herbs with purple spotted stem, purplish green spathe and purple spadix appendage.

Sikkim Himalaya ; 2400 - 2800 m.

Fl. & Fr. : April - June.

SCINDAPSUS Schott

Scindapsus officinalis (Roxb.) Schott in Schott & Endl., Melet. Bot. 21. 1832 ; Hook. f., F.B.I. 6 : 541. 1893. *Pothos officinalis* Roxb., Fl. Ind. 1 : 452. 1820.

Climbers. Leaves ovate, elliptic-ovate or nearly orbicular, 12.5 - 35 x 6 - 15 cm. Spathe up to 15 cm long, green, yellow within ; spadix equalling spathe, greenish yellow.

Sikkim, s.l. ; 300 m.

TYPHONIUM Schott

Typhonium diversifolium Wall. ex Schott, Aroid. 13 : t. 20 . 1855 ; Hook. f., F.B.I. 6 : 510. 1893.

var. **diversifolium**

Tuberous herbs. Spathe erect, green striped , purple tinged-dark purple, spadix - appendage, Female flowers purple.

Lachen ; 2400 - 4300 m.

Fl. & Fr. : June - Sept.

var. **microspadix** Engl., Pflanzenr. 73 : 111. 1920.

Differs from var. *diversifolium* in its smaller inflorescence.

Lingtoo, Yumthang ; 3650 - 3950 m.

Fl. & Fr. : July - Sept.

T. trilobatum (L.) Schott in Wien. Zeitschr. 3 : 72. 1829 ; Hook. f., F.B.I. 6 : 509. 1893. *Arum trilobatum* L., Sp. Pl. 965. 1753.

Herbs with subglobose tubers. Leaves 12.5 - 30 cm in diam., lobed ; lobes ovate. Spathe 7.5 - 30 cm long ; limb 3.7 - 10 cm broad, red-purple inside. Appendage bright red.

Sikkim, s.l.; ca 450 m.

Fl. & Fr. : July - Aug.

JUNCAGINACEAE

(R.C. Srivastava)

TRIGLOCHIN L.

Triglochin maritima L., Sp. Pl. 339. 1753 ; Hook. f., F.B.I. 6 : 563. 1893.

Scapigerous, marshy herbs. Leaves 1 - 3 mm wide, slightly flattened at the tip. Flowers small, brown-green. Fruits oblong - ovoid.

Llonakh, Gaigong to Kareng, Lachen, Thangu, Makula, Lungnak ; 2600 - 4600 m.

Fl. & Fr. : June - Aug.

T. palustris L., Sp. Pl. 38. 1753 ; Hook. f., F.B.I. 6 : 563. 1893.

Scapigerous, herbs. Leaves flaccid, up to 1 mm wide. Perianth green with purplish margins. Fruits narrowly clavate.

Lachen, above Lambi, Tangla, Thangshing ; 2900 - 4700 m.

Fl. : April - Aug.

POTAMOGETONACEAE

(R.C. Srivastava)

POTAMOGETON L.

Potamogeton filiformis Persoon, Syn. 1 : 152. 1805 ; Karthik. et al., l.c. 284.

Herbs, up to 40 cm high. Stems slender, branched at base. Leaves submerged, consisting of 2 tubes, blunt apex. Flower-whorls distant.

Chamgong, Lhonak ; 3600 - 4600 m.

Fl. : Sept.

P. nodosus Poir. in Lam., Encycl. Suppl. 4 : 535. 1816. **P. indicus sensu** Hook. f., F.B.I. 6 : 565. 1893.

Herbs. Stems terete. Leaves oblong-elliptic or elliptic-lanceolate, coriaceous ; petioles stout. Spikes densely flowered.

Lachung ; ca 2700 m.

Fl. : Aug.

P. octandrus Poir., Encycl., Suppl. 4 : 534. 1816. **P. javanicus** Hassk. in Act. Sci. Ind. Nersl. 1, 8 : 26. 1856 ; Hook. f., F.B.I. 6 : 566. 1892.

Herbs, up to 60 cm high. Stem slender. Leaves ovate-oblong, entire. Flowers minute, in interrupted whorls.

Thangu ; ca 4200 m.

Fl. & Fr. : April - June.

P. pectinatus L., Sp. Pl. 127. 1753 ; Hook. f., F.B.I. 6 : 567. 1893.

Herbs. Stem filiform, dichotomously branched. Leaves filiform. Flowers interruptedly whorled.

Lungmathang ; 3600 - 5100 m.

Fl. & Fr. : Oct.

In river bed.

SPARGANIACEAE**SPARGANIUM L.**

Sparganium fallax Graebner in Allg. Bot. Zeitsch. 4 : 32. 1898.
S. simplex sensu Hook f., F.B.I. 6 : 490. 1893, *non* Huds. 1778.

Aquatic perennial herbs. Leaves linear, spongy below. Inflorescence monoecious. Flowers dark brown.

Chaten ; *ca* 2670 m.

ERIOCAULACEAE

(P. Basu)

ERIOCAULON L.

Eriocaulon alpestre Hook. f. & Thoms. *ex* Koern. in Mus. Bot. Lugd. Bot. 3 : 163. 1867 ; Hook. f., F.B.I. 6 : 578. 1893.

Herbs. Leaves subulate. Seeds yellowish, glabrate.

Lachen valley, Lachung, Chola-range ; 1500 - 3000 m.

E. luzulaefolium Mart. in Wall., Pl. Asiat. Rar. 3 : 28. 1832 ; Hook. f., F.B.I. 6 : 582. 1893.

Herbs. Heads truncate. Seeds yellow.

Subtropical zone of E. Sikkim District ; up to 1700 m. Common.

E. nepalense Prescott *ex* Bong in Mem. Acad. Imp. Sci. St. Petersb. ser. B., Sci. Match. 1 : 610. 1830 ; Hook. f., F.B.I. 6 : 581. 1893.

Herbs. Leaves broadly ligulate. Seeds ciliate.

Gangtok ; *ca* 1500 m.

E. oryzetorum Mart. in Wall., Pl. Asiat. Rar. 3 : 28. 1832 ; Hook. f., F.B.I. 6 : 579. 1893.

Herbs with distinct stem. Leaves ensiform. Seeds minute, yellow.

Sikkim, *s.l.* ; 600 - 800 m.

E. seaboldianum Steud., Syn. Pl. Cyp. 2 : 272. 1855 ; Hook. f., l.c. 6 : 577. 1893.

Herbs with distinct stem. Leaves linear, flat. Seeds red-brown.

Below Gangtok ; 1200 - 1750 m.

E. truncatum Ham. ex Mart. in Wall., Pl. Asiat. Rar. 3 : 20. 1832 ; Hook. f., l.c. 6 : 578. 1893.

Herbs with distinct stem. Leaves subulate. Seeds pale yellow.

Alpine Sikkim.

CYPERACEAE

(R.C. Srivastava)

BLYSMUS Panz. ex J.A. Schultes

Blysmus compressus (L.) Panz. ex Link var. *sikkimensis* (Clarke) Karthik. in Karthik. et al., Fl. Ind. Enum. Monoct. 32 - 33. 1989. *Scirpus caricis* Retz. var. *sikkimensis* Clarke in Hook. f., F.B.I. 6 : 661. 1893.

Herbs. Stems 10 - 40 cm tall. Spikelets distichous. Nuts sessile, ellipsoid.

Lachen, Chamnago, Yumthang, Lhonak, Kupup, Muguthang, Chola ; 2700 - 4500 m.

Fl. & Fr. : July.

BULBOSTYLIS Kunth

Bulbostylis barbata (Rottb.) Clarke in Hook. f., F.B.I. 6 : 651. 1893. *Scirpus barabtus* Rottb., Progr. 27. 1772. & Descr. Ic. Pl. 52. t. 17. f. 4. 1773. *Stenophyllum barbata* (Rottb.) T. Cooke, Fl. Pres. Bombay 2 : 887. 1908 (3 : 401. 1967. Repr. ed). *Fimbristylis barbata* (Rottb.) Benth., Fl. Austr. 7 : 321. 1878.

Annual tufted herbs, up to 25 cm tall. Leaves in basal clusters, almost capillary, sheathed. Inflorescence terminal, capitate.

Jorethang - Ronbong bridge, Rangpo - Teesta ; 500 - 3000 m.

Fl. & Fr. : Sept.

B. densa (Ball.) Hand. - Mazz. in Karsten & Schenk, Vegetationsb. 20(7) : 16. 1930 ; Kern in Reinwardtia 6 : 51. 1961. *Scirpus densus* Wall. in Roxb., Fl. Ind. 1 : 231. 1820. *Isolepis trifida* Nees in Wight, Contr. Bot. Ind. 108. 1834. *Bulbostylis capillaris* Kunth var. *trifida* (Nees) Clarke in Hook. f., F.B.I. 6 : 652. 1893. *Bulbostylis trifida* (Nees) Nelmes in Kew Bull. 1950 : 209. 1950. *Fimbristylis capillaris* var. *trifida* (Nees) Mitra, Fl. Pl. E. India 1 : 108. 1958.

Annual, tufted herbs, up to 40 cm tall. Leaves in basal clusters, almost capillary, sheathed. Inflorescence anthelate.

Lachen, Lachung, Katao, Zemu, Gangtok, Pangthang, Dzongri, Hilay - Murhay, Chunghang, Kabi, Phodong, Thangu, Penthang, Yumthang, Thanka La ; 300 - 4100 m.

Fl. & Fr. : July - Sept.

CAREX L.

Carex alopecuroides D. Don *ex* Tilloch & Taylor in Phil. Mag. 62 : 455. 1823 ; Mabberley in Taxon 29 : 603. 1980. *Carex japonica* Thunb. var. *alopecuroides* (D. Don *ex* Tilloch & Taylor) Clarke in Hook. f., F.B.I. 6: 737. 1894. MONGSHER (Nep.).

Perennials, up to 80 cm tall. Stems trigonous, narrowly winged. Leaves clustered at base, one sometimes higher up also. Inflorescence racemose, 7 - 10 cm long. Utricles yellowish green. Nuts ellipsoid, triquetrous, deep brown.

Lachung, Gangtok, Tumlong ; 1450 - 1800 m. in rock crevices.

Fl. & Fr. : Aug. - Sept.

C. alpina Swartz in Liljeb. Svensk. Fl. ed. 2 : 26. 1796 ; Clarke in Hook. f., F.B.I. 6 : 730. 1894.

Herb, 10 - 65 cm tall. Leaves much shorter than the stems. Spikes 3 - 4, cuboid. Utricles ovoid-ellipsoid, yellow to pale brown.

Yumcho - la, Yumthang ; ca 4500 m.

Fl. & Fr. : April - June.

C. atrata L., Sp. Pl. 1387. 1753 ; Clarke in Hook. f., F.B.I. 6 : 731. 1894.

var. *atrata*

Herbs, up to 60 cm tall. Leaves equaling the stem. Spikes 3 - 6, 18 - 25 mm long. Utricle large ovoid or ellipsoid, inflated, yellow brown, beakless. Nuts not filling the utricle.

North-East Sikkim, Lhonak, Samdong, Lachen ; 3300 - 5100 m.

Fl & Fr. : July - Sept.

var. *glacialis* Boott, Illustr. 3 : 114. t. 365. 1862. *Carex duthiei* Clarke var. *glacialis* (Boott) Clarke in Hook. f., l.c. 732.

Herbs, up to 15 cm long. Leaves much shorter than stems. Spikes very small, ca 8 x 4 mm. Utricles small, slightly inflated.

Momay, Kankola, Yumthang Dzongri ; 3900 - 5100 m.

Fl. & Fr. : July - Aug.

var. **pullatta** Boott, Illustr. 3 : 114 . t. 364. 1862. *Carex duthiei* Clarke in Hook.f., l.c. 731. *Carex atrata* L. ssp. *pullata* (Boott) Kukenth. in Engl., Pflanzenr. 38 : 400. 1909.

Herbs. Lower spikes very shortly peduncled. Utricles narrow oblong. Nuts small stalked, nearly filling the utricles.

Changu, Bikbari, Lachen, Lhonak, Dzongri, Kupup ; 3300 - 4500 m.

Fl. & Fr. : July - Aug.

C. atrofusca Schkuhr., Riedgr. 1 : 106, f. 82. 1801. *C. atrofusca* Schkuhr. ssp. *minor* (Boott) T. Koyama in Ohashi, Fl. Pl. E. Himal. 3 : 122. 1975. *Carex ustulata* Wahlenb. var. *minor* Boott, Illustr. 1 : 71. t. 194. f. 1. 1858. *Carex ustulata* sensu Clarke in Hook.f., F.B.I. 6 : 734. 1894, non Wahlenb. 1803. *Carex atrofusca* var. *angustifructus* Kukenth. in Engl., Pflanzenr. 38 : 554. 1909.

Herbs, 10 - 40 cm tall. Spikes 2 - 5, terminal, male or bisexual. Utricles ellipsoid, shortly beaked.

Muguthang, Giagong plains, Chemathang to Thangshing, Momay, Zemu, Lhonak, Yumthang ; 3300 - 5300 m.

Fl. & Fr. : July.

C. baccans Nees in Wight, Contrib. Bot. Ind. 122. 1834 ; Clarke in Hook.f., F.B.I. 6 : 722. 1894. HARKATE (Nep.).

Perennials, up to 20 m tall. Leaves coriaceous, scattered all along and exceeding the stem. Panicles 30 - 85 cm long. Utricles globose, bright red on maturity.

Pakyong, Gangtok, Rimbi, Namchi, Deorali, Mamring, Chalisay ; 1200 - 2400 m.

Fl. & Fr. : Sept. - March.

C. bigelowii Torr. ex Schweinitz. in Ann. Lyc. New York 1 : 67. 1824. *Carex rigida* Gooden in Trans. Linn. Soc. 2 : 193. t. 22, f. 10. 1794, non Schrenk 1789.

Herbs, up to 20 cm tall. Leaves curved, flat. Spikes 3 - 4 together. Utricles ellipsoid.

Lachen ; ca 3600 m.

Fl & Fr. : July - Aug.

C. capillacea Boott, Illustr. 1 : 44. t. 110. 1858 ; Clarke in Hook. f., F.B.I. 6 : 713. 1894.

Herbs, 10 - 30 cm tall, tufted. Stems and leaves capillary. Spikes 6 - 8 x 2.5 - 3 mm, dense, Utricles ovoid, pyramidal.

Lachen, Yakla ; 2700 - 3600 m.

Fl. & Fr. : July - Aug.

C. continua Clarke in Hook. f., F.B.I. 6 : 717. 1894.

Perennials, 20 - 40 cm tall. Leaves many, much longer than stem. Inflorescence 10 - 20 x 3.5 cm long, interrupted at base. Utricles oblong-ellipsoid, trigonous, beak short, deeply bifid.

Rangeet river side, Teesta river side ; up to 500 m.

Fl. & Fr. : Sept. - Oct.

C. cruciata Wahlenb. in Kongl. Vetensk. Acad. Handl. Stockh. 24 : 149. 1803 ; Clarke in Hook. f., F.B.I. 6 : 715. 1894 ; Nelmes in Reinwardtia 1 : 277. 1951. *Carex condensata* Nees in Wight, Contrib. Bot. Ind. 123. 1834 ; Clarke in Hook. f., l.c. 716.

Perennials, up to 15 cm tall. Stems trigonous. Leaves coriaceous, all along and exceeding the stem. Panicles up to 50 cm long. Male glumes more than 3 mm. Utricles subsessile, inflated, ellipsoid, ovoid-oblong, subtrigonous, blackish brown.

Gangtok, Alichu near Busuk, Pakyong, Lachen, Namchi, on way to Katau ; up to 3300 m.

Fl. & Fr. : April - June (-Oct.).

On hill side walls.

var. *argocarpus* Clarke in Hook. f., F.B.I. 6 : 717. 1894.

Herbs. Secondary panicles pyramidal, dense. Utricles white, inflated, conspicuous.

Rangeet valley, Bakhim to Yoksum, Gangtok ; 600 - 2200 m.

Fl. & Fr. : Aug. - Oct.

C. cruenta Nees in Wight, Contrib. Bot. Ind. 128. 1834 ; Clarke in Hook. f., F.B.I. 6 : 734. 1894. *Carex ferruginea* sensu Clarke in Hook. f., l.c. 738 non Scop. 1772.

Herbs, up to 60 cm tall. Inflorescence ca 30 cm long. Spikes 5 - 12, remote ; terminal spike male. Utricles ellipsoid.

Zemu, Lachen ; 3900 - 4500 m.

C. daltonii Boott, Illustr. 1 : 5, t. 16. 1858 ; Clarke in Hook. f., F.B.I. 6 : 727. 1894.

Herbs, up to 70 cm tall. Leaves many, longer than stem. Inflorescence up to 45 cm long. Spikes *ca* 5 x 0.4 cm, pale yellow, terminal one often male. Utricles ellipsoid, trigonous ; beak oblong-linear, *ca* 2/3 of the utricles.

Chungthang; up to 3000 m.

Fl. & Fr. : April - June.

C. decora Boott, Illustr. 1 : 5. t. 15. 1858 ; Clarke in Hook. f., F.B.I. 6 : 725. 1894.

Herbs, 60 - 90 cm tall. Leaves much longer than stem. Inflorescence up to 45 cm long. Spikes 5, linear, purple-red, terminal one often wholly male. Utricles narrow-ellipsoid, trigonous, beaked.

Bakhim, Laghep, Tsomgo, Chola ; 2700 - 3600 m.

Fl. & Fr. : May - Nov.

C. filicina Nees in Wight, Contrib. Bot. Ind. 23. 1834, incl. var. *meiogyna* (Nees) Strachey. *Carex nilagirica* Steud., Syn. Cyp. 207. 1855. *Carex filicina* var. *minor* Boott, Illustr. 3 : 106. t. 317. & 318. 1862 ; Clarke in Hook. f., l.c.

Herbs, 20 - 40 cm tall. Leaves subbasal, much longer than stems, narrow. Inflorescence elongate, cylindric spikes many, up to 20 cm long. Spikelets 6 - 8 mm long. Utricles *ca* 3 mm long, brown.

Karponang, Phodonchen, Bakhim, Lachen, Tsoka, Tendong R. F., Lungthang, Mainam R.F., Gangtok, Pentong, Lachung - Yumthang, Panthong, Kyangosla, Chungthang, Aritar ; 1500 - 3700 m.

Fl. & Fr. : July - Aug.

C. finitima Boott, Illustr. 1 : 44. t. 112. 1858 ; Clarke in Hook. f., F.B.I. 6 : 736. 1894. *Carex finitima* var. *attenuata* Clarke in Hook. f. l.c. 736.

Perennials, up to 40 cm tall. Stems trigonous. Leaves clustered at base, shorter than stem. Inflorescence up to 22 cm long. Utricles ellipsoid, trigonous. Nuts ellipsoid, trigonous, brown.

Zemu, Karponang, Lachen ; 3000 - 3600 m.

Fl. & Fr. : May.

C. foliosa D. Don *ex* Tilloch & Taylor

Herbs. Leaves 5 - 6.5 mm wide, culms stout, up to 55 cm high, ridged, acutely trigonous. Inflorescence 8-10 cm long. Utricles *ca* 3.5 mm x 1.2 mm, hispid upwards.

Lachen ; 2150 - 2900 m.

Fr. : Aug. - Sept.

C. fragilis Boott, Illustr. 1 : 7, t. 21. 1858 ; Clarke in Hook. *f.*, F.B.I. 6 : 728. 1894.

Herbs, 20 - 35 cm tall. Leaves narrow, equaling the stem. Spikes *ca* 2.5 cm long. Utricles oblong obovoid ; beak linear.

Lachen, Lachung ; 2700 - 3300 m.

C. fucata Boott *ex* Clarke in Hook. *f.*, F.B.I. 6 : 710. 1894.

Herbs. Spikes dense, blackish. Utricles ellipsoid, nerves not distinct.

Dzongri, Toong, Phedang, Mamaichu, Lachen ; 2700 - 3600 m.

Fl. & Fr. : Aug. - Oct.

C. fusiformis Nees in Wight., Contrib. Bot. Ind. 128. 1834 ; Clarke in Hook. *f.*, F.B.I. 6 : 736. 1894.

Herbs, up to 45 cm tall. Leaves flat, up to 30 cm long. Spikes 6 - 12, lax. Utricles ellipsoid, beak oblong-linear.

Dzongri, Toong, Phedong, Mamichu Lake, Lachen ; 3000 - 3600 m.

Fr. & Fr. : Aug. - Oct.

C. gracilenta Boott *apud* Strachey, Cat. Pl. Kumaon 73. 1854. *Carex alpina* var. *gracilenta* (Boott) Clarke in Hook. *f.*, F.B.I. 6 : 730. 1894.

Very slender herbs. Leaves *ca* 1.5 mm wide. Spikes small.

Lachen, Kupup, Eumtso - la, Dzongri, Thangshing, Phedang, Thangu, Yumthang ; 3300 - 4200 m.

Fl. & Fr. : June - Sept.

C. haematostoma Nees in Wight, Contrib. Bot. Ind. 125. 1834 ; Clarke in Hook. *f.*, F.B.I. 6 : 744. 1894. *C. nakaoana* T.Koyama in Acta Phytotax. Geobot. 15 : 112. 1954.

Herbs. Stem up to 65 cm tall. Leaves flat, up to 50 cm long. Spikes purple, 1.2 - 6.2 x 3 - 0.6 cm. Utricles compressed, ovoid.

Dzalep, Ningbil, along Choktsering-chu, Lachen, Yumthang, Dzongri, Thangu, Thangshing ; 3300 - 5100 m.

Fl. & Fr. : May - Aug.

C. inaequalis Boott ex Clarke in Hook. f., F.B.I. 6 : 726. 1894
emend. Kukenth. in Engl., Pflanzenr. 38 : 545. 1909.

Herbs, 30 - 40 cm tall. Leaves narrow, equalling the stem. Inflorescence *ca* 15 cm long. Spikes linear. Utricles small, narrow, ellipsoid, trigonous ; beak oblong-linear, 2/3 of utricles.

Lachen ; 2100 - 3300 m.

Fl. & Fr. : April - July.

C. inanis Kunth, Enum. Pl. 2 : 522. 1837 ; Clarke in Hook.f., F.B.I. 6 : 743. 1894. *Carex schlagintweitiana* Boeck., Cyp. Nov. 1 : 48. 1888 ; Clarke in Hook.f., *l.c.* *Carex setigera* D. Don var. *schlagintweitiana* (Boeck.) Kukenth. in Engl., Pflanznr. 38 : 419. 1909.

Herbs, up to 37 cm tall. Leaves narrow, exceeding the stem. Spikes *ca* 16 mm long, terminal male. Utricles small, ovoid, trigonous, hispid beaked.

Zemu, Lachen, Lachung, Yumthang, Yangmaquala, Karponag, Tumbok, Yoksum to Mintok-khola ; 2400 - 3000 m.

Fl. & Fr. : June - Nov.

C. inclinis Boott ex Clarke in Hook.f., F.B.I. 6 : 728. 1894.

Herbs, 10 - 25 cm tall. Leaves many, up to 30 cm long. Spikes solitary, males on top, females at base. Utricles narrow, ellipsoid, trigonous.

Lachen ; 2100 - 3900 m.

Fl. & Fr. : June - Aug.

C. indica L., Mant. Pl. 2 : 574. 1771 ; Clarke in Hook.f., F.B.I. 6 : 744. 1894.

Perennials up to 10 cm tall. Leaves coriaceous, all along and exceeding the stem, with dark brown sheath. Panicles up to 30 cm long. Utricles subsessile, ellipsoid. Nuts obovoid, subglobose.

Sikkim, s.l. ; ca 1500 m.

Fl. & Fr. : Aug.

C. insignis Boott, Illustr. 1 : 5, t. 10. 1858 ; Clarke in Hook.f., F.B.I. 6 : 725. 1894.

Perennials, up to 50 cm tall. Stems triquetrous, covered by red sheaths. Leaves evenly scattered throughout the stem. Spikes long, linear, lax, darkpurple. Utricles ellipsoid, trigonous.

Gangtok, Sakyong, on way to Bakhim, Pentong, Kewzing, Bitu ; 1500 - 2100 m.

Fl. & Fr. : July-Aug.

C. laeta Boott, Illustr. 1 : 69, t. 188. 1858 ; Clarke in Hook.f., F.B.I. 6 : 745. 1894.

Herbs, up to 15 cm tall, caespitose. Spikes slender 2 - 3 distant, terminal male. Utricles obovoid-ellipsoid, trigonous.

Tsomgo, Tungu ; 3600 - 3900 m.

Fl. & Fr. : June - Sept. Endemic.

C. lehmannii Drejer, Symb. Caricol. 13. t. 2. 1844 ; Clarke in Hook.f., F.B.I. 6 : 730. 1894.

Stout herbs. Lowest spike 2.5 - 7.5 cm distant ; lowest bract much overtopping the inflorescence. Utricles up to 2 mm.

Zemu, Changu, Chola, Yumthang, Dzongri ; 3600 - 3900 m.

Fl. & Fr. : July - Aug.

C. longipes D. Don ex Tilloch & Taylor in Phil. Mag. 62 : 455. 1823 ; Clarke in Hook.f., F.B.I. 6 : 704. 1894. *C. nepalensis* Spr., Syst. 3 : 811. 1826. *C. longipes* var. *nepalensis* (Spr.) Boott, Ill. 4. : 190. 1867 ; Clarke in Hook.f., l.c. 704.

Herbs, 15-60 cm long. Leaves shorter than stem. Inflorescence very long, lax. Spikes greenish yellow. Utricles flattened, ovoid or ellipsoid, beaked.

Lachung to Yumthang, Lachen ; 2100 - 2700 m.

Fl. & Fr. : May - Aug.

C. microglochin Wahlenb. in Handl. Kong. A cad. Stockh. 24 : 140. 1803 ; Clarke in Hook.f., F.B.I. 6 : 711. 1894.

Herbs, up to 10 cm high. Leaves almost equalling the culms, subbasal, inrolled. Inflorescence consisting of a single androgynous spike. Utricles narrowly lanceolate, finally deflexed.

Lhonak, Gamothang ; 3600 - 4100 m.

Fl. & Fr. : May - Sept.

C. monopleura Kretz. in Not. Syst. Herb. Inst. Bot. Acad. Sci. URSS 7 : 35. 1937 ; Nelmes in Kew Bull. 1950 ; 208. 1950. *C. rochebrunii* Franch. & Sav., Enum. Fl. Japon. 2 : 555. 1878. *C. remota* var. *rochebrunii* (Franch & Sav.) Clarke in Hook.f., F.B.I. 6 : 706. 1894.

Herbs, 20 - 30 cm long. Leaves up to 20 x 0.3 cm. Spikes 6 - 9, oblong. Utricles *ca* 4 mm long ; beak triangular.

Bakhim - Yoksum, Lachen, above Tsoka, before Yumthang ; 2700 - 3600 m.

Fl. & Fr. : June - Aug.

C. moorcroftii Falc. ex Boott in Proc. Linn. Soc. 20 : 140. 1851 ; Clarke in Hook. f., F.B.I. 6 : 733. 1894.

Herbs, up to 20 cm high. Leaves basal, shorter than culms. Spikes large, reddish black, terminal pale. Utricles large, ellipsoid, trigonous, abruptly contracted into beak, pale upwards.

Giagong to Kareng, Chholamo Gurudongmar, Yumechu, Llonakh, 4800 - 5100 m.

Fl. & Fr. : Aug.

C. munda Boott, Illustr. 1 : 7, t. 20. 1858 ; Clarke in Hook.f., F.B.I. 6 : 727. 1894.

Herbs, 25 - 45 cm tall. Leaves equalling the inflorescence. Spikes oblong-linear. Utricle ellipsoid, trigonous, greenish yellow, beaked.

Yakchey - Yumthang, Lachen, Lunguthang, on way to Katau from Lachung, Karponang, Dzongri, Leghep, Changu, Chola, Zemu ; 3000 - 3900 m.

Fl. & Fr. : July - Aug.

C. myosurus Nees in Wight, Contrib. Bot. Ind. 122. 1834 (excl. rep. Pl. Nepal) ; Clarke in Hook.f., F.B.I. 6 : 723. 1894. *C. myosurus* var. *eminens* (Nees) Clarke in Hook.f., F.B.I. 6 : 723. 1894. *Carex eminens* Nees

in Wight, Contrib. Bot. Ind. 122. 1834. *C. myosurus* var. *rotongensis* Clarke in Hook. f., F.B.I. 6 : 723. 1894.

Herbs, up to 90 cm tall. Leaves often equalling the stem. Panicles ca 30 cm long, dark reddish brown. Utricles narrowly ellipsoid - trigonous.

Deorali, Gangtok, Rishi Khola ; 1800 - 2400 m.

Fl. : July - Sept.

C. nivalis Boott in Proc. Linn. Soc. 1 : 256. 1846 ; Clarke in Hook. f., F.B.I. 6 : 732. 1894.

Herbs. Terminal spikes wholly male, or female at top. Utricles broad, much compressed.

Lachen, Samdung ; 3300 - 5100 m.

C. notha Kunth, Enum. Pl. 2 : 421. 1837 ; Clarke in Hook. f., F.B.I. 6 : 709. 1894.

Herbs. Spikes linear-cylindric, brown. Utricles ellipsoid ; beak linear, short.

Upper Sikkim, Chola ; 3300 - 3600 m.

Fl. & Fr. : July - Aug.

C. nubigena D. Don ex Tilloch & Taylor in Phil. Mag. 62 : 455. 1823 ; Clarke in Hook. f., F.B.I. 6 : 702. 1894.

Herbs. Leaves long, narrow. Inflorescence ovoid to linear, 1.2 - 12 cm long, interrupted. Spikes ovoid, androgynous. Utricles ellipsoid.

Gangtok, Mintok-Khola to Prek-chu, Kupup, Karponang, Lachen, Lachung, Changu, Tosa, Yumthang ; 1800 - 4200 m.

Fl. & Fr. : May - June.

C. obscura Nees in Wight, Contrib. Bot. Ind. 126. 1834 ; Clarke in Hook. f., F.B.I. 6 : 731. 1894.

Herbs, 30 - 60 cm tall. Leaves overtopping the stem. Spikes blackish, terminal one occasionally male. Utricles oblong, beaked, shining yellow.

Chola, Manmaichu, Tosa, Zemu ; 3600 - 3900 m.

Fl. & Fr. : July - Aug.

var. *brachycarpa* Clarke in Hook. f., F.B.I. 6 : 731. 1894.

Utricle ovoid-ellipsoid, narrowed abruptly into a short oblong beak, scabrous.

Yumthang, Dzongri, Jamlingthang, Thangsing, Kupup, Pentangla, Zemu valley, Lachen, N.E. Sikkim ; 3300 - 3600 m.

Fl. : July - Aug.

O. *obscuriceps* Kukenth. in Engl., Pflanzenr. 38 : 723. 1909. **C. *lurida*** Clarke in Hook. f., F.B.I. 6: 742. 1894, *non* Wahlenberg

Herbs. Leaves inserted near base of culm, almost equaling culms. Spikes 4 - 5, distant, purple-red. Utricle-beak 2 - fid.

Lachen ; 2400 - 3600 m.

Fl. & Fr. : May - Aug.

C. *oedorrhapha* Nelmes in Kew Bull. 1939 ; 659. 1939. **Carex *tumida*** Boott, Illustr. 1 : 66. t. 181. 1858, *non* Beilschm. 1850 ; Hook. f., F.B.I. 6 : 741. 1894.

Herbs, 30 - 45 cm tall. Leaves longer than stem. Spikes 4 - 5, suberect, terminal one male. Utricles ellipsoid.

Sikkim, *s.l.* ; 2400 - 3000 m.

C. *olivacea* Boott in Proc. Linn. Soc. 1 : 286. 1845 ; Clarke in Hook. f., F.B.I. 6 : 741. 1894.

Herbs, up to 90 cm high. Leaves basal, one on culm, exceeding culms. Inflorescence of one terminal male and 3 - 9 female spikes. Utricles armed, turgid and trigonous, abruptly contract into beak.

Sikkim. *s.l.* ; 2000 - 3050 m.

Fl. & Fr. : April - Sept.

C. *orbicularis* Boott in Proc. Linn. Soc. 1 : 254. 1854. **C. *rigida*** *sensu* Clarke in Hook.f., F.B.I. 6: 711. 1894, *non* Gooden 1894. **C. *vulgaris*** *sensu* Clarke in Hook. f., F.B.I. 6 : 711. 1894, *non* Fries 1842.

Herbs, 10 - 25 cm tall. Leaves flat, recurved. Spikes 3 - 4, cylindric, dense. Utricle ellipsoid ; beak minute, entire.

Lachen ; 2700 - 3600 m.

Fl. & Fr. : July - Aug.

C. pamirensis Clarke ex Fedtsch in J. Bot. St. Petersb. 1 : 19. 1908.
C. rostrata sensu Clarke in Hook. f., F.B.I. 6 : 740. 1894, non Stokes 1787.

Herbs, up to 90 cm tall. Spikes 3 - 6, distant, erect, upper most 1 - 3 male, lower female. Utricle ovoid to ellipsoid, inflated, suddenly narrowed into a beak.

Lachen ; 2700 - 3600 m.

C. parva Nees in Wight, Contrib. Bot. Ind. 120. 1834 ; Clarke in Hook. f., F.B.I. 6 : 712. 1894.

Herbs. Spikes brown, 6 - 18 mm long. Utricles lanceolate, acuminate, long beaked, glabrous.

Kupup, Lachen; 3600 - 4800 m.

Fl. & Fr. : July - Aug.

C. phacota Spr. in L., Syst., Veg. ed. 16, 3 : 826. 1826 ; Clarke in Hook. f., F.B.I. 6 : 708. 1894.

Herbs, up to 75 cm tall, triquetrous, Leaves equalling the stem. Inflorescence 10 - 15 cm long. Spikes 3 - 14, linear-cylindrical. Utricles ovoid. Beak short triangular.

Lachung ; 1800 - 2100 m.

C. polyccephala Boott, Illustr. 1 : 4, t. 12. 1858 ; Clarke in Hook. f., F.B.I. 6 : 725. 1894.

Robust herbs, 30 - 45 cm tall. Leaves long, exceeding the inflorescence. Inflorescence elongate-oblong, dense. Spikes large, oblong-cylindric, straw coloured. Utricles ellipsoid, trigonous.

Dikeeling, Labdong (Lepdand) ; 1500 - 3700 m.

Fl. & Fr. : April - July.

C. praeclara Nelmes in Hook., Ic. Pl. 35. t. 3403. 1940.

Herbs, up to 20 cm high. Inflorescence forming a 2.5 - 3.5 cm, dense head. Spikes 4 - 5, broadly ovoid, androgynous. Utricles with up to 0.5 mm long beak.

Lhonakh, Jonsony-la valley ; 4880 - 5030 m.

Fl. : Aug.

C. praelonga Clarke in Hook. f., F.B.I. 6 : 707. 1894.

Herbs, 60 - 90 cm tall. Spikes 4 - 9, linear. Utricle obovate, yellow-brown.

Dikeeling, West of Dzongri ; 2250 - 2700 m.

Fl. & Fr. : April - May.

Carex pseudofoetida Kukenth. in Bot. Tidsskr. 28 : 225, f. 1. 1907.

Herbs, up to 20 cm high. Leaves subbasal, deeply channelled, inrolled, shorter than culms. Inflorescence a densely congested head, ovoid to oblong. Spikes 3, androgynous. Utricles narrowly elliptic, pale golden.

Kareng - Chholamo ; 3050 - 4780 m.

Fl. & Fr. : June - Aug.

C. psychrophila Nees in Wight, Contrib. Bot. Ind. 127. 1834 ; Clarke in Hook. f., F.B.I. 6 : 732. 1894.

Herbs, up to 60 cm tall. Leaves up to 45 x 0.4 cm. Spikes cylindric, terminal one female. Utricles small, obscurely nerved.

Yumthang ; ca 3600 m.

C. pulchra Boott, Illustr. 1 : 4. t. 13. 1858 ; Clarke in Hook. f., F.B.I. 6 : 727. 1894.

Herbs, 20-40 cm tall. Leaves shorter than inflorescence. Inflorescence oblong. Spikes linear, brown. Utricles narrow, ellipsoid - trigonous.

Lachen, before Yumthang, Lachung, Zemu, Lachung, Karponang ; 2100 - 3300 m.

Fl. & Fr. : July.

C. radicalis Boott, Illustr. 1 : 56. t. 147. 1858 ; Clarke in Hook. f., F.B.I. 6 : 729. 1894.

Herbs. Culms almost capillary, 20 - 30 cm tall. Spikes 1 - 3, androgynous. Utricles ovoid, hairy, dull green.

Lachen ; 3000 - 3300 m.

Fl. & Fr. : July - Aug.

C. rara Boott in Proc. Linn. Soc. 1 : 284. 1854 & Illustr. 1 : 44. t. 109. 1858.

Herbs, up to 40 cm tall. Stem filiform, trigonous. Leaves few, filiform, clustered at base. Spikelets solitary, terminal androgynous. Utricle ovoid - oblong, beaked.

Laghep, Phedung Yumthang valley, Changu, Chola, Yakla, Legshisp ; 3000 - 3900 m.

Fl. & Fr. : July - Aug.

C. royleana Nees in Wt., Contrib. Bot. Ind. 127. 1834. *C. breviculmis sensu* Clarke in Hook. f., F.B.I. 6 : 746. 1894, *non* R. Br. 1810.

Herbs. Stem caespitose, 10 - 40 cm tall. Leaves 6 - 25 x 0.3 cm, flat. Spikes 2 - 5, oblong, terminal one male, female with 8 - 14 utricles. Utricles small ellipsoid, trigonous, sparsely pubescent.

Lachung ; 2700 - 3000 m.

C. sahnii Ghildyal & Bhattacharyya in J. Econ Tax. Bot. 7 : 447. 1985. *Carex kingiana* Clarke in Kew Bull. Addl. Ser. 8 : 75. 1908, *non* Lev. & Van. 1902 ; Ghildyal in Nayar & Sastry, Red Data Book Ind. Pl. 2 : 102. 1988.

Herbs. Spikelets linear-cylindric, androgynous, interrupted at base. Utricles oblong-ellipsoid, trigonous.

Phodong ; *ca* 1800 m. Endemic.

C. setigera D. Don in Trans. Linn. Soc. 14 : 330. 1824 ; Clarke in Hook. f., F.B.I. 6 : 743. 1894. *C. aristata* D. Don *ex* Tilloch & Taylor in Phil. Mag. 62 : 455. 1823.

Herbs. Leaves equalling the stem. Spikes 4 - 8, cylindric, distant, terminal 1 (-2 -3) male, lowest female. Utricles ovoid, hispid, suddenly narrowed into small oblong notched beak.

Chungthang, Lachen ; 2100 - 2400 m.

C. setosa Boott, Illustr. 3 : 108, t. 327 - 329. 1862 ; Clarke in Hook. f., F.B.I. 6 : 745. 1894.

Herbs with setose stem, 15 - 30 cm long. Leaves flat. Spikes 2 - 5, terminal one male. Utricles oblong-obovoid, trigonous, setose.

Chamnago, Lachen, Namdee ; 2700 - 3600 m.

C. sikkimensis Clarke in Hook. f., F.B.I. 6 : 708. 1894. *C. fucata* var. *sikkimensis* (Clarke) Kukenth. in Engl., Pflanzenr. Heft. 38 : 344. 1908.

Herbs. Leaves narrow, long. Spikes 4 - 7, linear, clavate, female at top. Utricles ellipsoid, ca 2 mm long.

Dzongri ; 3600 - 3900 m.

C. spiculata Boott var. *nobilis* Clarke in Hook. f., F.B.I. 6 : 724. 1894. *C. nobilis* Boott, Illustr. 1 : 4, tt. 9 - 11. 1858. *C. myosurus* ssp. *spiculata* (Boott) Kukenth. var. *nobilis* Kukenth. in Engl., Pflanzenr. 38 : 259. 1909.

Perennials, up to 90 cm tall. Leaves coriaceous, much exceeding the stems. Panicles up to 40 cm long. Utricles subsessile, subcoriaceous, not inflated, ellipsoid, trigonous, gradually beaked. Nuts stipitate, ellipsoid, triquetrous, dark brown.

Ryang ; up to 1200 m.

Fl. & Fr. : July - Sept.

C. stramentitia Boott ex Boeck. in Linnaea 40 : 351. 1876 ; Clarke in Hook. f., F.B.I. 6 : 717. 1894.

Perennials, up to 12 cm tall. Leaves coriaceous, all along and exceeding the stem, up to 15 mm wide. Panicles up to 35 cm long. Utricles subsessile, slightly obovoid, trigonous, abruptly beaked. Nuts ovoid-ellipsoid, triquetrous, yellowish brown.

Pakyong, Burmiak, Rangpo ; up to 1400 m.

Fl. & Fr. : Dec. - May.

C. thomsonii Boott, Illustr. 1 : 1, t. 1. 1858 ; Clarke in Hook. f., F.B.I. 6 : 703. 1894.

Perennials, up to 30 cm tall. Stems trigonous. Leaves scattered all along the stem, sheaths black. Inflorescence spicate. Glumes and utricles purplish red, streaked and dotted. Nuts obovoid, deep brown.

Great Rangeet, Little Rangeet ; 300 - 600 m.

Fl. & Fr. : April.

C. vesiculosa Boott, Illustr. 3 : 107. t. 323. 1862 ; Clarke in Hook. f., F.B.I. 6 : 717. 1894.

var. vesiculosa

Perennials, up to 1 m tall. Stem trigonous. Leaves coriaceous ; sheaths yellowish brown. Inflorescence irregularly panicled, panicles narrow, up to 40 cm long. Glumes ovate-lanceolate, yellowish brown, red

streaked. Utricles ellipsoid. Nuts stipitate, ellipsoid, triquetrous, irregularly ribbed, not inflated, yellowish brown, beak ca half the size of utricle.

Rumtek-Martam ; 1000-1800 m.

Fl. & Fr. : June - Dec.

In dry to moist, open places, among boulders.

var. **pallida** (Kukenth.) R.C. Srivastava

var. **congesta f. pallida** Kukenth. in Engl., Pflanzenr. Heft. 38. IV, 20 : 238. 1958.

Plants forming clumps on hillside walls, smaller than var. *vesiculosus*. Spikes, glumes and nuts paler.

Bitu locality on way to Kabi and Chitrey-Phamtam.

Fl. & Fr. : July - Aug.

On hill side walls at moist locations.

C. wallichiana Spr., Syst. 3 : 812. 1826. *Carex foliosa* D. Don ex Tilloch & Taylor in Phil. Mag. 62 : 455. 1823. *Carex muricata* L. var. *foliosa* (D. Don ex Tilloch & Taylor) Clarke in Hook. f., F.B.I. 6 : 703. 1894.

Perennials, up to 90 cm tall. Stems triquetrous. Leaves usually shorter than stem. Inflorescence spicate, up to 10 cm long. Glumes ovate, pale brown. Utricles ovate-lanceolate, pale greenish. Nuts ovoid.

Lachen ; 2150 - 2900 m.

Fl. & Fr. : July - Sept.

CYPERUS L.

Cyperus alulatus Kern in Reinwardtia 1 : 463. f. 1. 1952. *Cyperus iria* var. *rectangularis* Kukenth. in Engl., Pflanzenr. 101. 152. 1935. *Cyperus rectangularis* (Kukenth.) Bennet in Indian Forester 95 : 692. 1969.

Annual herbs, up to 40 cm tall. Leaves equalling to or longer than the stem. Spikelets greenish yellow. Nuts obovoid, reddish brown.

Ranipool, Tingpong.

Fl. & Fr. : Aug. - Oct.

C. castaneus Willd., Sp. Pl. 1 : 278. 1797 ; Clarke in Hook. f., F.B.I. 6 : 598. 1893.

Annual herbs, up to 15 cm tall. Anthela usually simple, often reduced to a single head ; spikelets linear-oblong ; glumes oblong-spathulate. Nuts *ca* 1 mm long.

Geyzing, Jorethang - Rongbong Bridge Rd ; *ca* 500 m.

Fl. & Fr. : Sept. - Oct.

C. compressus L., Sp. Pl. 46. 1753 *emend.* Dandy in Exell. Cat. s. Tome 357. 1844 ; Clarke in Hook. f., F.B.I. 6 : 605. 1893. *Cyperus pectiniformis* R. & S., Mant. 2 : 128. 1824. *Cyperus compressus* var. *pectiniformis* (R. & S.) Clarke in J. Linn. Bot. 21 : 97. 1814.

Herbs. up to 10 cm high. Anthela simple, lax. Spikelets linear or oblong, compressed, greenish white to green, glumes ovate, 3 - 4 mm long. Nuts broadly obovoid, shortly apiculate.

Jorethang, Deorali (Gangtok), near Rabong - la, near Singtam, Melli ; 550 - 2100 m.

Fl. : June - Oct.

C. cuspidatus Kunth in H.B.K., Nov. Gen. Sp. 1 : 204. 1815 ; Clarke in Hook. f., F.B.I. 6 : 598. 1893. *Cyperus uncinatus* *sensu* Clarke in Phillip. J. Sci. Bot. 2 : 82. 1907, *non* Poir. 1806.

Annual herbs, up to 20 cm tall. Anthela rays 1 - 3 ; spikelets linear, compressed ; glumes ovate, aristate. Nuts *ca* 1 mm long, puncticulate.

Jorethang ; up to 2700 m.

Fl. & Fr. : May - June (-Oct.)

C. difformis L., Sp. Pl. 67. 1753 ; Clarke in Hook. f., F.B.I. 6 : 599. 1893.

Annuals, up to 70 cm tall. Anthela simple. Spikelets stellately spreading ; glumes 3-nerved. Nuts obovoid, trigonous, almost equalling the glumes.

Sikkim, *s.l.* ; up to 2400 m.

C. distans L.f., Suppl. 1.3. 1781 ; Clarke in Hook. f., F.B.I. 6 : 607. 1893.

Herbs: Anthela large, compound ; spikelets 10 - 20- flowered. Nuts oblong or narrowly ellipsoid.

Singtam - Ranipool, Jorethang ; up to 900 m.

Fl. & Fr. : Aug. - Sept.

C. exaltatus Retz., Obs. Bot. 5 : 117. 1789 ; Clarke in Hook. f., F.B.I. 6 : 617. 1893.

Herbs, up to 1 m tall. Spikes long cylindric, lax ; spiklets linear, 8 - 20 - flowered, brightly coloured.

Dikeeling ; *ca* 1000 m.

Fl. & Fr. : Sept. - Oct.

C. halpan L., Sp. Pl. 45. 1753 ('*haspan*') ; Clarke in Hook. f., F.B.I. 6 : 600. 1893.

Annual herbs, up to 30 cm tall. Anthela compound or decompound ; spikelets linear-lanceolate ; glumes ovate - oblong. Nuts *ca* 0.5 mm long.

Tangpo - Teesta, Rishi - Khola, Dzongu, Tendong ; up to 1700 m.

Fl. & Fr. : April - Oct.

C. involucratus Rottboell.

Perennial herbs. Stems leafless, tufted. Basal sheaths papery. Inflorescence umbellate ; bracts leaflike. Spikelets in umbels of 5 - 9. Nuts oblong-elliptic, acute.

Gangok ; *ca* 1750 m. Cult.

C. iria L., Sp. Pl. 45. 1753 ; Clarke in Hook. f., F.B.I. 6 : 606. 1893.

Cyperus parviflorus Nees in Wight, Contrib. Bot. Ind. 87. 1834 *excl. syn.*, *non* Vahl 1806. *Cyperus iria* var. *parviflorus* (Nees) Miq., Fl. Ind. Bat. 3 : 270. 1859. *Cyperus paniciformis* Franch. & Sav., Pl. Jap. 2 : 103. 537. 1879. *Cyperus kamphoeveneri* Boeck., Cyp. Nov. 1 : 5. 1888 ; Clarke in Hook. f., F.B.I. 6 : 594. 1893.

Annual herbs, up to 60 cm tall. Stem compressed, trigonous. Anthela simple or compound ; spikelets oblong, glumes keeled. Nuts broadly striperate, ovoid-ellipsoid, triquetrous, *ca* 1.5 mm long.

Ranipool, Gangtok, Rangpo - Teesta, Jorethang, Temi, below Runetek, Singtam ; up to 1150 m.

Fl. & Fr. : June - Oct.

C. laxus Lamk., Ill. 1 : 146. 1791 *emend.* Poir. in Lamk., Encycl. 7 : 260. 1806 ; J. Raynal in Adansonia 2, 17 : 277. 1978. *Cyperus*

diffusus Vahl, Enum. Pl. 2 : 321. 1805 ; Clarke in Hook. f., F.B.I. 6 : 603. 1893.

Herbs 20 - 75 cm tall. Leaves equalling the stem, flat. Spikelets 3 - 9 together, digitate. Nuts broadly ellipsoid, triquetrous.

Ratepani - Suntaley, Rishi, near Singtam, Rangpo ; 350 - 750 m.

Fl. & Fr. : Aug. - Sept.

C. niveus Retz., Obs. Bot. 5 : 12. 1788 ; Clarke, l.c. 601.

Slender herbs, up to 40 cm tall. Spikelets 3 - 12 mm strongly compressed, linear-oblong, 20 - 40-flowered. Nuts oblong - ovoid.

Sikkim, s.l. ; up to 1800 m.

Fl. & Fr. : Feb. - Aug.

C. nutans Vahl, Enum Pl. 2 : 363. 1806 ; Clarke, l.c. 607.

Herbs, up to 1 m tall. Anthela compound. Spikelets loosely spicate.

Legship, N.W. of Singtam, Namli.

Fl. : Aug. - Oct.

C. pangorei Rottb., Descr. Ic. Rar. Nov. Pl. 31. t. 7. f. 3. 1773.

Cyperus corymbosus Rottb. var. *pangorei* (Rottb.) Clarke in J. Linn. Soc. 21 : 292. 1884 & in Hook. f., F.B.I. 6 : 612. 1893. *Cyperus tegetum* Roxb., Fl. Ind. 1 : 208. 1832 ; Clarke in Hook. f., l.c. 613.

Perennials, up to 80 cm tall. Spikelets clustered. Glumes linear, obtuse, ca 3 mm long. Nuts narrowly ovoid-oblong, trigonous, ca 1.5 mm long.

Melli ; up to 300 m.

Fl. & Fr. : June - Oct.

C. pilosus Vahl, Enum. Pl. 2 : 534. 1806 ; Clarke in Hook. f., F.B.I. 6 : 609. 1893. *Cyperus obliquus* Nees in Wight, Contrib. Bot. Ind. 86. 1834. *Cyperus pilosus* var. *obliqua* (Nees) Clarke in J. Linn. Soc. 21 : 151. 1884 & in Hook. f., l.c. 610. *Cyperus pilosus* var. *polyantha* Clarke, l.c. 148. & in Hook. f., l.c.

Perennial herbs, up to 50 cm tall. Leaves canaliculate or plicate. Anthela compound, lax ; spikes ovoid ; spikelets linear ; glumes ovate, muticus or mucronulate. Nuts ovoid, ca 1.2 mm long.

Temi, near Singtam, Phodong, Rinchingpong, Ranipool, Rangeet, valley, Jorethang ; 500 - 600 m.

Fl. & Fr. : June - Oct.

C. rotundus L., Sp. Pl. 45. 1753 ; Clarke in Hook. f., F.B.I. 6 : 614. 1893.

Perennial herbs, up to 30 cm tall. Stems triquetrous. Anthela lax ; bracts foliaceous ; spikelets linear-lanceolate, glumes ovate, keeled. Nuts oblong-obovoid, ca 1.5 mm long, black.

Gangtok, Teesta, Saramsa, near Ranipool, Jorethang, Legship, Namthang ; up to 2150 m.

Fl. & Fr. : March - June.

C. silleensis Nees in Wight, Contrib. Bot. Ind. 79. 1834 ; Clarke in Hook. f., F.B.I. 6 : 600. 1893.

Annual herbs, up to 30 mm tall, tufted. Stems trigonous. Leaves 4 - 6, shorter than to subequaling the stem. Anthela compound ; spikelets in clusters, lanceolate ; glumes ovate-lanceolate, greenish stramineous to reddish, keeled. Nuts ellipsoid, apiculate ca 0.8 mm long, brownish.

Rangpo, Teesta ; ca 350 m.

Fl. & Fr. : June - July.

C. tenuiculmis Boeck. in Linnaea 36 : 286. 1870. *Cyperus zollingeri* sensu Clarke in Hook. f., F.B.I. 6 : 613. 1893, non Steud. 1855; Kern in Reinwardtia 2 : 108. f. 6. 1952. *Cyperus zollingeri* var. *condensatus* Kukenth. in Engl., Pflanzenr. 101 : 134. 1935.

Annuals or perennials, up to 30 cm tall, tufted. Stems triquetrous, grooved. Leaves 5 - 8, half to 2/3 of stem, margins revolute. Anthela simple or compound, lax ; spikelets shortly spicate, in clusters of 2 - 15 ; glumes elliptic-lanceolate, muticus, fulvous. Nuts obovoid, triquetrous, blackish.

Gangtok (near Elephat Mansion, Deorali), Rishi, Yoksum, near Runtek.

Fl. & Fr. : July - Oct.

C. tenuispica Steud., Syn. Cyp. 11. 1855. *Cyperus flavidus* sensu Clarke in Hook. f., F.B.I. 6 : 600 - 601. 1893, non Retz. 1789.

Annual herbs, up to 4 cm tall. Anthela lax ; glumes laxly arranged, ovate-oblong, slightly keeled towards the top. Nuts *ca* 0.5 mm long, tubercled, white.

Sikkim, *s.l.* ; in rice fields.

ELEOCHARIS R. Br.

Eleocharis congesta D. Don, Prodr. Fl. Nepal. 41. 1825 ; Clarke in Hook. *f.*, F.B.I. 6 : 630. 1893. *Eleocharis afflata* Steud., Syn. Cyp. 2 : 76. 1855 ; C.B. Clarke in Hook. *f.*, *l.c.* 629. *Eleocharis subvivipara* Boeck. in Linnaea 36 : 424. 1870 ; Clarke in Hook. *f.*, *l.c.* *Eleocharis chaetaria* var. *subvivipara* (Boeck.) C.E. C. Fischer in Gamble, Fl. Pres. Madras 1648. 1931. (3 : 1145. 1967, Repr. ed.) *Eleocharis congesta* var. *subvivipara* (Boeck.) T. Koyama in J. Fac. Sci. Univ. Tokyo 3, 8 : 90. 1962.

Annuals or perennial herbs, up to 40 cm tall. Glumes ovate to lanceolate ; bristles retrorsely scabrous. Nuts *ca* 1 mm long ; style base shortly pyramidal triangular.

Ranipool, Penlong - la, Yoksum, Phedong to Kabi, Gangtok ; up to 2300 m.

Fl. & Fr. : May - Oct.

E. dulcis (Burm. *f.*) Trin. *ex* Henschel, Vita Rumph. 186. 1833. *Andropogon dulce* Burm. *f.*, Fl. Ind. 219. 1768. *Scirpus platagineus* Retz., Obs. Bot. 5 : 14. 1789. *Eleocharis plantaginea* (Retz.) R. & S., Syst. Veg. 2 : 150. 1817 ; Clarke in Hook. *f.*, F.B.I. 6 : 625. 1893.

Perennial herbs, up to 20 cm tall. Spikelets 1.5 - 6 x 3 - 6 mm; bristles retrorsely scabrous, light brown; styles 2 - 3 fid. Nuts obovate; style base triangular.

Sikkim *s.l.* ; *ca* 350 m.

Fl. & Fl. : Nov. - Dec.

E. palustris (L.) R. Br., Prodr. 224. 1810 ; Clarke in Hook. *f.*, F.B.I. 6 : 628. 1893. *Scirpus palustris* L., Sp. Pl. 70. 1753.

Perennial herbs, up to 30 cm tall. Spikes ovoid or subcylindric, densely many-flowered ; glumes membranous, ovate-lanceolate, muticus.

Lachen, Lhonak ; 3000 - 4310 m.

Fl. & Fr. : June - Aug.

E. tetraquetra Nees in Wight, Contrib. Bot. Ind. 113. 1834 ; Clarke in Hook. f., F.B.I. 6: 630. 1893.

Herbs, 20 - 90 cm tall. Spikelets oblong-ellipsoid. Nuts obovoid, trigonous.

Lachen, Lachung, Chungthang, North Rangeet ; 450 - 3300 m.

Fl. & Fr. : May - Oct.

ERIOPHORUM L.

Eriophorum comosum Wall. ex Nees in Wight, Contrib. Bot. Ind. 110. 1834 ; Clarke in Hook. f., F.B.I. 6 : 664. 1893. *Scirpus comosus* Wall. in Roxb., Fl. Ind. 1 : 234. 1820.

Perennial herbs, up to 45 cm long. Leaves almost equalling the stem. Anthella decompound or supra decompound, with many ellipsoid spikelets ; glumes keeled. Nuts sessile, ellipsoid, ca 3 mm long.

Chalisay, Geyzing, Rongli Khola, Padamchen, Rorathang, Rangpo, on way to Pakyong, Jorethang, Ratepani, Singtam, Marchak before Dikchu ; up to 1200 m.

Fl. & Fr. : Sept. - Oct.

Notes : On hillside walls along Teesta ad Rangeet valleys.

E. microstachyum Boeck. in Linnaea 38 : 399. 1874 ; Clarke in Hook. f., F.B.I. 6 : 664. 1893.

Herbs, 5 - 15 cm long, slender. Spikelets 1 - 3, subcapitate, brown.

Sikkim, s.l. ; 2400 - 4800 m.

Fl. & Fr. : Sept. - Oct.

FIMBRISTYLIS Vahl

Fimbristylis aestivalis (Retz.) Vahl, Enum. Pl. 2 : 288. 1806 ; Clarke in Hook. f., F.B.I. 6 : 637. 1893. *Scirpus aestivalis* Retz., Obs. Bot. 4 : 12. 1786. *Fimbristylis bis-umbellata* sensu C.E.C. Fischer in Gamble, Fl. Pres. Madras 1898. 1936 (3 : 1151. 1967 Repr. ed.), *non* Bub. 1850.

Annuals, up to 20 cm tall. Leaves setaceous. Spikelets solitary ; glumes ovate, keels green. Nuts elliptic or obovate.

Lachen, Yoksum, Dentam - Pamianchi, Gangtok, Ranipool ; up to 1700 m.

Fl. & Fr. : April - Aug.

F. bisumbellata (Forssk.) Bub., Dodec. 30. 1850. *Scirpus bisumbellatus* Forssk., Fl. Age.-Arab. 15. 1775. *Fimbristylis dichotoma* sensu Clarke in Hook. f., F.B.I. 6 : 635. 1893, *non* (L.) Vahl, 1806.

Annual herbs, up to 20 cm tall. Anthela compound to decomound ; spikelets solitary ; glumes keeled. Nuts biconvex, shortly stipitate.

Royang, Rumtek ; 600 - 2500 m.

Fl. & Fr. : July - Sept.

F. complanata (Retz.) Link, Hort. Bot. Berol. 1 : 292. 1827 ; Clarke in Hook. f., F.B.I. 6 : 646. 1893. *Scirpus complanatus* Retz., Obs. Bot. 5 : 14. 1789.

Annual herbs. Anthela compound or decomound ; spikelets lanceolate to oblong-ovoid ; glumes keeled. Nuts broadly ovoid, trigonous, *ca* 1 mm long, verruculose.

Lindok, above Yoksum, on way to Yumthang, ; 1800 - 2100 m.

Fl. & Fr. : July - Aug.

F. dichotoma (L.) Vahl, Enum. Pl. 2 : 287. 1806. *Scirpus dichotomus* L., Sp. Pl. 50. 1753. *Fimbristylis diphylla* Vahl, l.c. 289 ; Clarke in Hook. f., F.B.I. 6 : 636. 1893 (*excl. var. pluristriata*). *Fimbristylis diphylla* var. *annua* (R. & S.) Clarke in Hook. f., l.c. 637. *Fimbristylis annua* R. & S. var. *paucispiculata* Blatt. & Mcc. in J. Bombay Nat. Hist. Soc. 37 : 544. 1934.

Annual herbs, up to 80 cm tall. Leaves usually shorter than stem. Anthela simple to decomound ; spikelets ovoid-globose. Nuts ovoid, *ca* 1 mm, shortly stipitate.

Near Phensong, on way to Lachung, Tingbong, Gezing, Rangpo - Teesta, Gangtok, Chungthang - Lachung, Temi, Dikchu, near Pheonsong ; up to 2700 m.

Fl. & Fr. : June - Oct.

F. littoralis Gaud. in Freyc., Voy. Bot. 413. 1826. *F. miliacea* sensu Benth., Fl. Austral. 316. 1878 ; Clarke in Hook. f., F.B.I. 6 : 644. 1893, *p.p.* *omne non* Vahl 1806, *non Scirpus miliaceus* L.

Annual or perennial herbs, up to 80 cm tall. Flowering shoots many, usually leafless ; anthela lax, supradecomound ; spikelets ovoid ; glumes ovate. Nuts obovoid or suborbicular, trigonous, densely verrucose, pale brown.

Pachekhani, Ranipool, on way to Rishi ; up to 900 m.

Fl. & Fr. : June - Oct.

F. miliacea (L.) Vahl, Enum Pl. 2 : 287. 1806, *emend.* Blake in J. Arnold Arb. 35 : 216. 1954 ; Clarke in Hook. f., F.B.I. 6 : 644. 1893. *Scirpus miliaceus* L., Syst. Veg. ed. 10, 2 : 868. 1759.

Herbs. Leaves shorter to equaling culm. Anthela lax, compound or decomound. Spikelets oblong-ovoid. Nuts broadly ellipsoid to subglobose.

Phodong to Kabi, Pachekhani, up to 1800 m.

Fl. & Fr. : July - Oct.

F. schoenoides (Retz.) Vahl, Enum. Pl. 2 : 286. 1806 ; Clarke in Hook. f., F.B.I. 6 : 634. 1893. *Scirpus schoenoides* Retz., Obs. Bot. 5 : 14. 1789.

Herbs, up to 20 cm tall. Leaves 0.5 - 1 mm wide. Anthela simple or reduced to a solitary spikelet. Spikelets broadly ovoid, terete. Nuts stipitate, obovoid.

Ranipool ; ca 750 m.

Fl. & Fr. : July - Aug.

F. stolonifera Clarke in Hook. f., F.B.I. 6 : 637. 1893.

Herbs with slender creeping stolons covered with dull fibrous scales. Inflorescence simple. Spikelets up to 7.5 x 4 cm. Glumes reddish brown.

Phodong ; 1680 - 2900 m.

Fl. & Fr. : June - Sept.

F. thomsonii Boeck. in Linnaea 37 : 37. 1871 ; Clarke in Hook. f., F.B.I. 6 : 646. 1893.

Annual or perennial herbs, up to 80 cm tall. Leaves equalling to or longer than the stem. Anthela compound to supradecomound ; spikelets ellipsoid ; glumes spiral, ovate-oblong. Nuts obovoid, trigonous, brownish.

Rinchimpang ; up to 1500 m.

Fl. & Fr. : June - July.

INDOCOURTOISIA Bennet & Raiz.

Indocourtoisia cyperoides (Roxb.) Bennet & Raiz. in Ind. Forester 107 : 432. 1981. *Kyllinga cyperoides* Roxb., Fl. Ind. 1 : 187. 1820.

Courtoisia cyperoides (Roxb.) Nees in Linnaea 9 : 286. 1834 ; Clarke in Hook. f., F.B.I. 6 : 625. 1893. *Mariscus cyperoides* (Roxb.) A. Dietrich. in L., Sp. Pl. ed. 6, 2 : 438. 1833.

Herbs. Stems tufted, trigonous. Leaves shorter than to exceeding stems ; sheaths very long, pale. Inflorescence up to 11.5 x 8.5 cm. Spikelets flat, elliptic. Nuts linear-elliptic, sharply trigonous, apiculate, pale brown.

Sikkim, s.l. (cf. Kukenth. 1936) ; ca 300 m.

Fl. : Dec.

ISOLEPIS R. Br.

Isolepis setacea (L.) R. Br. 222. 1810 ; T. Koyama in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 112. 1978. *Scirpus setaceus* L., Sp. Pl. 73. 1753 ; Clarke in Hook. f., F.B.I. 6 : 654. 1893.

Herbs, 2.5 - 20 cm tall. Leaves ca 2.5 cm long, setaceous. Spikelets 1 - 3, in a quasilateral head. Nuts obovoid, trigonous.

Lachung - Yumthang, Lachen - Lodge bridge, Thangu, Zemu ; 3600 - 3900 m.

Fl. & Fr. : June - Oct.

KOBRESIA Willd.

Kobresia capillifolia (Decne.) Clarke in J. Linn. Soc. Bot. 20 : 378. 1884 & Hook. f., F.B.I. 6 : 697. 1894. *Elyna capillifolia* Decne. in Jacquem., Voy. 4 (Bot.) : 173. t. 174. 1840.

Herbs, up to 30 cm tall. Leaves linear-setaceous. Spikes linear. Nuts ellipsoid.

Chholamo, Giagong, Gurudongmar, above Dikchu ; 3900 m.

Fl. & Fr. : July.

K. clarkeana (Kukenth.) Kukenth. in Engl. Pflanzenr. 38 : 48. 1909. *Schoenoxiphium clarkeanum* Kukenth. in Bull. Herb. Boiss. 4, 2. : 49. 1904.

Herbs, tufted, up to 35 cm high. Leaves almost equalling culms. Inflorescence a linear panicle ; lateral spikes ca 7, appressed, upper overlapping. Prophylls oblong, blunt.

Cho - le - la ; ca 3800 m.

Fl. & Fr. : June.

K. curticeps (Clarke) Kukenth. in Engl., Pflanzenr. 38 : 47. 1909.
Carex curticeps Clarke in Hook. f., F.B.I. 6 : 729. 1894.

Herb with 30 - 60 cm long slender culms. Leaves linear, up to 30 cm long, flat. Inflorescence linear, 2.5 - 12 cm long ; spikes loosely arranged. Achenes narrowly elliptic or oblong, slightly beaked.

Dzongri, Chamnago, Changu, Chola, Chakung - Chu, Gnathang, Tsomgo, Lungthang, Kyangoshla ; 3300 - 3700 m.

Fl. & Fr. : June - July.

K. curvata (Boott) Clarke in Kew Bull. Addl. Ser. 8. 68. & 137. 1908. *Carex curvata* Boott, Illustr. Carex 1 : 2, t. 5. 1858 ; Clarke in Hook. f., F.B.I. 6 : 728. 1894.

Herbs, up to 36 cm tall. Leaves filiform, 2 - 16 cm long. Inflorescence linear or linear-oblong, 1.5 - 8 cm long. Nuts elliptic - ovoid, ca 2 mm long.

Kupup, Karponang, Nathula, Kyangosla, Lachung, Tsomgo, Thangu, Changu ; 3700 - 4900 m.

Fl. & Fr. : July - Aug.

K. duthiei Clarke in Hook. f., F.B.I. 6 : 697. 1894.

Herbs up to 40 cm tall. Leaves linear, flat, 1 - 16 cm long. Inflorescence erect, consisting of a single linear or linear-oblong spike. Nuts obovoid-oblong, shortly beaked.

Nathula, Bikbari, Dzongri, to Prek-chu, Jelep - La ; 3900 - 4200 m.

Fl. & Fr. : July - Aug.

K. esenbeckii (Kunth) Noltie in Edinb. J. Bot. 50 : 43. 1993. *Carex esenbeckii* Kunth, Enum Pl. 2 : 522. 1837, *non* Boott 1845. *K. seticulmis* Boeck. in Linnaea 39 : 3. 1875 ; Clarke in Hook. f., F.B.I. 6 : 695. 1894. *Carex trinervis* Nees in R. Wight, Contrib. Bot. Ind. 120. 1834, *nom. illegit.* *Kobresia trinervis* Boeck. in Linnaea 39 : 4. 1875, *nom. illegit.* *K. hookeri* var. *dioica* Clarke in Hook. f., F.B.I. 6 : 695. 1894. *K. angusta* Clarke, l.c. 965.

var. esenbeckii

Herbs, 4 - 30 cm tall. Leaves shorter than equalling the stem, linear, flat. Inflorescence erect, consisting of a single linear androgynous spike. Nuts linear, beaked.

Chakung - Chu, Tosa, Sherabthang, Lachen, Changu, Namdee, Llonakh, Tsomgo Telepla, Momay ; 3300 - 3900 m.

Fl. & Fr. : May - July.

Notes : A polymorphic species.

var. *fissiglumis* (Clarke) Noltie in Edinb. J. Bot. Soc. 50 (1) : 43. 1993. *K. fissiglumis* Clarke in Hook. f., F.B.I. 6 : 696. 1894.

Differs from var. *esenbeckii* in its unisexual spikes in having prophylls open to the base, smooth keel and female glumes being oblong, truncate, with distinct hyaline apex.

Bikbari to Chaunrikiang, west of Chemathang, Thegu ; 4250 - 4300 m.

K. gammiei Clarke in Kew Bull. Addl. ser. 8 : 68. 1908.

Herbs, up to 20 cm long. Leaves almost equalling the stem. Inflorescence 4 - 5 cm long. Spikes 10 - 12 mm long. Styles 3-fid.

Tongsup near Dzongri ; to Gamothang, Jamlingthang Bikbari, Monlepcha, Thangshing to Lam Pokhri ; 3400 - 3800 m.

Fl. & Fr. : July - Aug.

K. laxa Nees in Wt., Contrib. Bot. Ind. 119. 1834 ; Clarke in Hook. f., F.B.I. 6 : 698. 1894.

var. *laxa*

Herbs, 3 - 35 cm tall. Leaves linear, flat, up to 30 cm long. Inflorescence bright red, lax, paniculate ; glumes of lowest spike leafy. Prophylls linear. Nuts narrowly oblong or elliptic-oblong, ca 3 mm long.

Lachen ; 3400 - 3800 m.

Fl. & Fr. : July - Aug.

var. *curvirostris* (Clarke) R.C. Srivastava. *Kobresia curvirostris* Clarke in Hook. f., F.B.I. 6 : 699. 1894.

Herbs, up to 15 cm tall, slender. Leaves as stem, ca 2 mm wide. Spikes densely panicled. Panicles ca 2.5 x 0.8 cm. Nuts narrowly oblong ; beak cylindric, exert, curved.

Thangu ; ca 4200 m.

***K. nepalensis* (Nees)** Kukenth. in Engl., Pflanzenr. 38 : 40. 1909. *Uncinia nepalensis* Nees in Wt., Contrib. Bot. Ind. 129. 1834. *Carex linearis* Boott, Illustr. 1 : 51, t. 136. 1858 ; Clarke in Hook. f., F.B.I. 6 : 712. 1894. *K. nepalensis* var. *elachista* (Clarke) Kukenth. in Engl., Pflanzenr. 38 : 40. 1909. *Carex linearis* var. *elachista* Clarke in Hook. f., F.B.I. 6 : 713. 1894.

var. nepalensis

Herbs, up to 40 cm tall. Leaves filiform, equaling or longer than stem. Inflorescence linear, up to 7 cm long. Spikes androgynous, male spike terminal. Nuts oblong or lanceolate, beaked.

Chakung - chu, Shaerbthang, Jalep - la, Nathu - la, Lachen ; 3900 - 5700 m.

Fl. & Fr. : July - Aug.

var. vaginosa (Clarke) R.C. Srivastava *K. vaginosa* Clarke in Hook. f., F.B.I. 6 : 695. 1894.

Herbs, up to 20 m tall. Stems sheathed up to almost three - fourth. Leaves filiform. Spikes dioecious. Nuts linear - oblong.

Momay, near Bikbari ; 4300 - 4900 m.

Fl. & Fr. : July - Sept.

K. prainii Kükenth. in Bull. Herb. Boiss. 4, 2 : 50. 1904 & in Engl., Pflanzenr. 38 : 40. 1909. *K. utriculata* Clarke in Kew Bull. Add. Ser. 8 : 67. 1908.

Herbs, up to 20 cm tall. Leaves linear, almost equaling culms. Nuts oblong-elliptic, minutely apiculate, not beaked.

Makula, Muguthang ; 3350 - 5270 m.

Fl. & Fr. : May - Sept.

K. pygmaea (Clarke) Clarke in Hook.f., F.B.I. 6 : 696. 1894. *Hemicarex pygmaea* Clarke in J. Linn. Soc. Bot. 20 : 383. 1883.

Herbs, 5 - 60 mm tall, base covered with brown lamina bearing outer sheaths. Leaves up to 6 cm long. Inflorescence erect, consisting of a single unbranched spike, unisexual or androgynous spikes. Nuts oblong or obovate, slightly beaked.

Kareng-Cholamo, Llonakh, Momay, Kongra-la, near Tebhla Pass, Muguthang ; 3900 - 4500 m.

Fl. & Fr. : May - Sept.

K. royleana (Nees) Boeck. in Linnaea 39 : 8. 1875 ; Clarke in Hook. f., F.B.I. 6 : 698. 1894. *Trilepis royleana* Nees in Edinb. New Phil. J. 17 : 267. 1834 & in Wight, Contrib. Bot. Ind. 119. 1834.

var. royleana

Herbs, up to 40 cm tall. Leaves linear, flat, 3 - 40 cm long. Inflorescence erect, linear-oblong, consisting of many congested, elliptic or oblong spikes. Nuts oblong, *ca* 2.2 mm long.

Dzongri, Samdung ; 3500 - 4800 m.

Fl. & Fr. : July - Sept.

var. *humilis* (C. A. Mey) Kukenth. in Engl., Pflanzenr. 38 : 46. 1909. *Elyna humilis* C.A. Mey *apud* Trautv. in Acta Horti Petrop. 1 : 21. 1871. *Kobresia persica* Kukenth. & Bornm. in Oest. Bot. Z. 47 : 133. t. 2. 1897.

Samding, Kareng - Chholamo ; *ca* 4900 m.

Fl. & Fr. : July - Sept.

K. schoenoides (C. A. Mey) Steud., Sy. Cyp. 246. 1855 ; Clarke in Hook. f., F.B.I. 6 : 697. 1894. *Elyna schoenoides* C.A. Mey in Ledeb. Fl. Alt. 4 : 235. 1853.

Stout herbs, up to 16 cm tall, base covered with coriaceous laminaless outer sheaths. Leaves linear-filiform, 6 - 16 cm long. Inflorescence erect of a single oblong, 1 - 3 x 0.3 x 0.5 cm spike. Nuts obovoid, *ca* 2 x 1 mm, slightly beaked, trigonous.

Jalep- La, Nathu-La, Kupup, Kareng-Cholamo, Pa to Yongdi, Ynorth-Donkiala, Llonakh ; 3900 - 5700 m.

Fl. & Fr. : May - June.

K. sikkimensis Kukenth. in Engl., Pflanzenr. 38 : 47. 1909 ; Clarke in Kew Bull. Addl. Ser. 8 : 145. 1908.

Herbs, up to 60 cm tall. Leaves linear, flat, upto 32 x 0.3 cm. Inflorescence oblong, 4 - 6 cm long, lax. Spikes loosely arranged, unisexual ; sterile rachillas included in the prophyll.

Dzongri, Gamothang ; 3800 - 4600 m.

Fl. & Fr. : July - Aug.

K. uncinoides (Boott) Clarke in Hook. f., F.B.I. 6 : 698. 1894. *Carex uncinoides* Boott, Illustr. 1 : 8. t. 23. 1858.

Herbs, up to 35 cm tall. Leaves linear, flat, up to 21 cm long. Inflorescence erect, oblong, 1 - 7 x 1 cm ; spikes closely arranged. Nuts oblong or oblanceolate, slightly beaked, trigonous.

Sherabthang, Lachung, Lachen, Changu, Dzongri, Chakung-Chu, Chola, Megu, Yakla, Jalepla, Kupup, Nathula, Tsomgo ; 3300 - 4300 m.

Fl. & Fr. : May - July (-Oct.)

K. vidua (Boott ex Clarke) Kukenth. in Engl., Pflanzenr. 38 : 40. 1909. *Carex vidua* Boott ex Clarke in Hook. f., F.B.I. 6 : 713. 1894.

Herbs, 14 - 20 cm tall. Leaves equal to or shorter than stem, linear. Inflorescence dioecious consisting of a single spike. Nuts oblong, beaked.

Lachen, Thangshing to Lam Pokhri ; 3900 - 4400 m.

Fl. & Fr. : July.

KYLLINGA Rottb.

Kyllinga brevifolia Rottb., Descr. & Ic. 13. t. 4. f. 3. 1773 ; Clarke in Hook. f., F.B.I. 6 : 588. 1893. *Cyperus brevifolius* (Rottb.) Hassk., Cat. Hort. Bogor 24. 1844.

Perennial herbs with long, creeping, slender rhizomes. Culms arranged in a single row along rhizome. Leaves radical and subradical, shorter than culms. Inflorescence a single globose head (rarely 2 or 3), greenish or brownish. Nuts ellipsoid, obtuse.

Singtam, Namli, Lachung, Lachen, Ranipool, Busuk, Gangtok, Saramsa, Yoksum ; up to 2400 m.

Fl. & Fr. : June - Sept.

K. nemorallis (J. R. & G. Forst.) Dandy ex Hutch. & Dalziel., Fl. W. Trop. Afr. 2, 2 : 487. 1836. *Thryocephalon nomorale* J. R. & G. Forst., Charact. Gen. Pl. 129. t. 65. 1776. *Kyllinga monocephala* Rottb., Descr. Ic. Nov. Pl. 13. f. 4. 1773, *nom. illeg.* ; Clarke in Hook. f., F.B.I. 6 : 588. 1893.

Perennial herbs, laxly tufted with long creeping rhizomes. Culms 10 - 45 cm tall, 1 - 1.5 mm thick. Leaves many, equal to or shorter than culms. Inflorescence a single head, bearing a globose, central spike and 2 - 3 small lateral spikes. Nuts yellowish brown.

Saramsha, near Jorethang, Tashiding, Melli R.F., Singtam ; up to 1400 m.

Fl. & Fr. : July - Sept.

K. odorata Vahl ssp. *cylindrica* (Nees) T. Koyama in Gard. Bull. Singapore 30 : 161. 1977. *K. cylindrica* Nees in Wight, Contrib. Bot. India 91. 1834 ; Clarke in Hook. f., F.B.I. 6 : 588. 1893.

Herbs. Stems not thickened at base, 10 - 30 cm tall. Leaves much shorter than stem. Heads of 3 spikes, medium cylindric, lateral much shorter, subovoid, straw coloured. Nuts yellow-brown, black on maturity.

Saramsha ; Rango - Teesta, 1100 - 1700 m.

Fl. & Fr. : June - July.

LIPOCARPHA R. Br.

Lipocarpha chinensis (Obs.) Kern in Blumea Suppl. 4 : 167. 1958.

Scirpus chinensis Obs., Daqb. Ostind. Resa 220. 1757. *Hypaelyptum argenteum* Vahl, Enum. Pl. 2 : 283. 1806. *Lipocarpha argentea* (Vahl) R. Br. ex Nees in Linnaea 9 : 287. 1835 ; Clarke in Hook. f., F.B.I. 6 : 667. 1893.

Herbs. Rhizomes lacking. Stem 5 - 40 cm tall, obtusely trigonous. Leaves equalling to or shorter than stems. Inflorescence of a single head. Spikelets 1 - 8, pale or fuscous. Nuts sessile, obovoid or ellipsoid, obtuse.

Lachen, Phodong to Kabi, near Rani pool, Phensong ; up to 1800 m.

Fl. & Fr. : July - Aug.

MARISCUS Vahl

Mariscus compactus (Retz.) Boldingh., Zakf. Landb. Java 77. 1916.

Cyperus compactus Retz., Obs. Bot. 5 : 10. 1789. *Cyperus dilutus* Vahl, Enum Pl. 2 : 357. 1806. *Mariscus microcephalus* Presl., Rel. Haenk. 1 : 182. 1830 ; Clarke in Hook. f., F.B.I. 6 : 624. 1893. *Mariscus dilutus* (Vahl) in Wight, Contrib. Bot. Ind. 90. 1834.

Perennial herbs, up to 50 cm tall. Stems spongy, trigonous or subterete, 2 - 5 mm thick. Leaves 5 - 7. Anthela compound or decumbent, greenish when young, bright reddish brown with age.

Namthang, Saramsa ; up to 1400 m.

Fl. & Fr. : May - June.

M. cyperinus (Retz.) Vahl, Enum. Pl. 2 : 377. 1806 ; Clarke in Hook. f., F.B.I. 6 : 621. 1893. *Kyllinga cyperina* Retz., Obs. Bot. 6 : 21. 1791. *Cyperus cyperinus* (Retz.) Valck.-Sur., Gesl. Cyp. Mal. Arch. 154. t. 6, f. 10. 1898.

var. **cyperinus**

Perennial herbs. Culms 10 - 60 cm tall, 1 - 3 mm thick, triquetrous. Leaves several at base. Anthela simple open but with short rays, sometimes nearly head-like.

Lower Siche Basti, Gangtok, on way to Rumtek, Saramsa, Ronglikhola, Melli, Chuchachen, Dentam, Namli ; up to 1700 m.

Fl. & Fr. : May - Sept.

var. *bengalensis* Clarke in Hook. f., F.B.I. 6 : 621. 1893. *Cyperus cyperinus* var. *bengalensis* (Clarke) Kukenth. in Engl., Pflanzenr. 101 : 520. 1936.

Umbels contracted into 1 head ; spikes shortly cylindric or ovoid ; spikelets linear, bearing 2 - 4 nuts in fruits, spreading at right angles.

Sikkim, s.l. ; up to 1200 m.

Fl. & Fr. : May - July.

var. *laxatus* (Clarke) Kukenth. in Engl., Pflanzenr. 101 : 519. 1936. *Cyperus umbellatus* Clarke var. *laxatus* Clarke in J. Linn. Soc. 21 : 201. 1888. *Mariscus tenuifolius* Schrad. ex Nees, Fl. Bras. 2 : 46. 1842 ; Clarke in Hook. f., F.B.I. 6 : 622. 1893. *Mariscus cyperinus* ssp. *laxatus* (Clarke) T. Koyama in Gdns. Bull. Singapore 30 : 157. 1977.

Herbs with very narrow leaves. Umbels simple, contracted. Spikes solitary, cylindric or ovoid, lax ; spikelets bearing usually 2 nuts.

Gangtok, Dzongu, Chungthang ; up to 1800 m.

Fl. & Fr. : May - June.

M. dubius (Rottb.) C.E.C. Fischer in Gamble, Fl. Pres. Madras 1644. 1931 (3 : 1142. 1967. Repr. ed.). *Cyperus dubius* Rottb., Descr. & Ic. Pl. 21. t. 4. f. 5. 1773. *Mariscus dregeanus* Kunth, Enum. Pl. 2 : 120. 1837 ; Clarke in Hook. f., F.B.I. 6 : 620. 1893.

Herbs up to 30 cm tall, without stolons. Leaves often equalling stem. Umbels contracted in 1 very dense ovoid head ; spikelets ovoid, bearing 2 - 6 nuts.

Gangtok ; ca 1800 m.

Fl. & Fr. : June - Aug.

M. hookerianus Clarke in Hook. f., F.B.I. 6 : 623. 1893. *Cyperus sikkimensis* Kukenth. in Engl., Pflanzenr. 101 : 467. 1936, non *Cyperus hookerianus* Arn. nec *C. hookeri* Boeck.

Herbs with horizontal, woody rhizomes. Culms up to 50 cm tall. Inflorescence of 7 cylindric spikes. Spikelets oblong, bearing 4, trigonous, dull white, nuts.

Sikkim (cf. Hook. f. 1893).

Fl. : July.

M. paniceus (Rottb.) Vahl, Enum. Pl. 2 : 378. 1806 ; Clarke in Hook. f., F.B.I. 6 : 620. 1893. *Kyllinga panicea* Rottb., Descr. Ic. 15, t. 4. f. 1. 1773. *Cyperus paniceus* (Rottb.) Boeck. in Linnaea 36 : 381. 1870.

Perennial, tufted herbs. Culms up to 45 cm tall, 1 - 2 mm thick, trigonous. Leaves 3 - 7, shorter than the culm. Anthela simple, contracted, greenish.

Dzongu, Namchi, Melli ; up to 1200 m.

Fl. & Fr. : May - Dec.

M. squarrosus (L.) Clarke in Hook. f., F.B.I. 6 : 623. 1893 ; Hooper in Saldanha & Nicolson, Fl. Hassan Dt. 690. 1976. *Cyperus squarrosus* L. in Torner, Cent. Pl. 2 : 6. 1756. *Cyperus aristatus* Rottb., Descr. Pl. Rar. Progr. 22. 1772. Clarke in Hook. f., l.c. 606. *Mariscus aristatus* (Rottb.) Tang & Wang in Fl. Rep. Pop. Sin. 11 : 178. t. 59, f. 12 - 13. 1963 ; T. Koyama in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 116. 1978.

Small, annual herbs. Culms 3 - 10 cm tall, 0.5 - 1.5 mm thick, triquetrous. Anthela simple, open or often reduced to a single hemispherical spike.

Lachung, Lachen, Keodom, Chungthang ; 2400 - 2700 m.

Fl. & Fr. : May - Oct.

M. sumatrensis (Retz.) Raynal in Adansonia 15 : 110. 1975. *Kyllinga sumatrensis* Retz., Obs. Bot. 4 : 13. 1786. *Scirpus cyperoides* L., Mant. Pl. 2 : 181. 1771. *Mariscus sieberianus* Nees ex Clarke in Hook. f., F.B.I. 6 : 622. 1893. *Cyperus cyperoides* (L.) O. Ktze., Rev. Gen. Pl. 3 : 333. 1898.

var. sumatrensis

Perennial herbs, not stoloniferous. Culms 1 - many together, erect, triquetrous, up to 80 cm tall, 1 - 3 mm wide. Leaves many, equal to or shorter than culm. Anthela open, simple.

Tashiding, Chunchachen, Rumtek, Burtuk, Basti, Phodong, Aho, Ronglikhola, Nam Nam road (Deorali), Sada, Yoksum, Temi, Singtam ; 400 - 1800 m.

Fl. & Fr. : March - Dec.

var. evolutior (Clarke) Karthik. in Karthik. *et al.*, l.c. 64. 1989. *Mariscus sieberianus* var. *evolutior* (Clarke) Hook. f., F.B.I. 6 : 622. 1893.

Cyperus cypereoides var. *evolutior* (Clarke) Kukenth. in Sinensis 3 : 80. 1932 & Engl., Pflanzenr. 101 : 516. 1926.

Herbs with linear spikelets bearing 2 - 4 nuts.

Sikkim, s.l.

Fl. & Fr. : March - Dec.

PYCREUS P. Beauv.

Pycrus diaphanus (Schrad. ex R. & S.) Hooper & T. Koyama in J. Jap. Bot. 51 : 316. 1976. *Cyperus diaphanus* Schrad. ex R. & S., Mant. 2 : 477. 1824. *Cyperus latespicatus* Boeck. in Flora 42 : 433. 1859. *Pycrus latespicatus* (Boeck.) Clarke in Hook. f., F.B.I. 6 : 590. 1893. *Cyperus diaphanus* var. *latespicatus* (Boeck.) Kern in Steenis, Fl. Males. 1, 7 : 653. 1974.

Herbs. Stems 10 - 30 cm tall. Leaves half in length to stem. Inflorescence simple. Spikelets spicate, ca 4 mm wide, straw coloured or reddish brown. Nuts obovoid, compressed, black.

Gangtok (New Cantt.) ; 1100 - 2100 m.

Fl. & Fr. : Sept. - Oct.

P. flavidus (Retz.) T. Koyama in J. Jap. Bot. 51 : 316. 1976. *Cyperus flavidus* Retz., Obs. Bot. 5 : 13. 1789. *Cyperus globosus* All., Fl. Pedem 49. 1789, non Forssk. 1775. *Cyperus capillaris* Koen. ex Roxb., Fl. Ind. 1 : 198. 1820. *Pycrus globosus* (All.) Reichb., Fl. Germ. Excurs. 2 : 140. 1830. *Pycrus capillaris* (Koen. ex Roxb.) Nees ex Clarke in Hook. f., F.B.I. 6 : 591. 1893.

var. **flavidus**

Herbs, up to 80 cm tall. Leaves 1 - 4, shorter than or equalling the stems. Anthela simple. Spikelets many, linear, 15 - 50 -flowered. Nuts obovoid, lenticular, apiculate, blackish brown

Rinkinpung, up to 2000 m.

Fl. & Fr. : June - Sept.

var. **erectus** (Clarke) Karthik. in Karthik. et. al., Fl Ind. Enum. Monocot. 65. 1989. *Pycrus globosus* var. *erectus* Clarke in J. Linn. Soc. 35 : 15. 1898 & 36 : 205. 1903. *Cyperus globosus* var. *erectus* (Clarke) Kukenth. in Engl., Pflanzenr. 101 : 356. 1936. *Cyperus flavidus* var. *erectus* (Clarke) Korla in Bull. Bot. Surv. Ind. 9 : 236. (1967) 1968.

Anthela simple or compound. Spikelets cinnamomeus to purplish with age, 20 - 40-flowered.

Sikkim, s.l. ; up to 1200 m.

Fl. & Fr. : April - May.

P. polystachyos (Rottb.) B. Beauv., Fl. Oware 2 : 48, t. 86, f. 2. 1807 ; Clarke in Hook. f., F.B.I. 6 : 592 1893.

Herbs, up to 60 cm high, tufted. Stems leafy below. Leaves flat, much shorter than the stem. Inflorescence a dense compound head. Spikes with 4 - 11 linear spikelets.

Sikkim, s.l. (BM).

P. pumilus (L.) Nees ex Clarke in Hook. f., F.B.I. 6 : 591. 1893. *Cyperus pumilus* L., Cent. Pl. 2 : 6. 1756.

Herbs, up to 25 cm tall. Leaves shorter than stem. Anthela simple or subcompound. Spikelets 10 - 40 in subglobose heads ; 8 - 20-flowered. Nuts ovoid, apiculate, blackish brown.

Phensong, N. W. of Singtam, Tashiding, up to 1400 m.

Fl. & Fr. : June - Nov.

P. sanguinolentus (Vahl) Nees ex Clarke in Hook. f., F.B.I. 6 : 590. 1893. *Cyperus sanguinolentus* Vahl, Enum Pl. 2 : 351. 1806.

ssp. **sanguinolentus**

Herbs. Stems 10 - 50 cm long. Leaves equaling or much shorter than stems. Inflorescence simple. Spikelets shortly spicate or clustered, linear or oblong. Nuts ovoid, greyish brown.

Lindok, Pangthang, Rishi, Lindok, Kupup, Lachen, Phadonchen ; up to 3200 m.

Fl. & Fr. : July - Aug.

ssp. **melanocephalus** (Miq.) Karthik. *et al.*, Fl. Ind. Enum. Monocot. 66. 1989. *Cyperus melanocephalus* Miq., Fl. Ind. Bat. 3 : 259. 1859. *Cyperus sanguinolentus* f. *melanocephalus* (Miq.) Kuekenth. in Engl., Pflanzenr. 101 : 386. 1936. *Cyperus sanguinolentus* ssp. *melanocephalus* (Miq.) Kern in Reinwardtia 3 : 55. 1954.

Herbs, up to 25 cm high. Leaves 1 - 2 mm broad. Anthela capitately contracted ; bracts erect or suberect. Nuts ca 1.5 mm long.

Rishi ; 1000 - 1700 m.

Fl. & Fr. : May - Sept. (-Dec.).

P. stramineus Clarke in Hook. f., F.B.I. 6 : 589. 1893. *Cyperus stramineus* Nees in Wight, Contrib. Bot. Ind. 74. 1834., *non* Desf. 1820. *Cyperus substramineus* Kukenth. in Engl., Pflanzenr. 101 : 398. 1936.

Herbs, 5 - 30 cm tall. Inflorescence reduced to 1 compound head, of 5 - 12 straw coloured. Spikelets 15 - 40 (-70) - flowered. Nuts obovoid, blackish brown.

Sikkim, s.l. : up to 1700 m.

Fl. & Fr. : Aug. - Oct.

P. sulcinux (Clarke) Clarke in Hook. f., F.B.I. 6 : 593. 1893. *Cyperus sulcinux* Clarke in J. Linn. Soc. 21 : 56. 1884.

Herbs up to 40 cm tall. Anthela simple, lax, spikelets 40 - 50-flowered. Nuts subsessile, linear-oblong, blackish brown.

Legship, Phensang ; up to 1500 m.

Fl. & Fr. : July - Nov.

P. uniloides (R. Br.) Urban, Symb. Antill. 2 : 164. 1900. *Cyperus uniloides* R. Br., Prodr. 216. 1810.

Herbs, up to 90 cm high. Leaves shorter than culms. Inflorescence simple umbel. Spikes of 3 - 20 spikelets. Spikelets up to 22 - flowered, yellow or brownish yellow.

Melli - Teesta ; up to 2600 m.

Fr. : Aug.

RHYNCHOSPORA Vahl *nom. cons.*

Rynchospora corymbosa (L.) Britt. in Trans. New York Acad. Sci. 11 : 84. 1892. *Scirpus corybosus* L., Gent. Pl. 2 : 7. 1756. *Rynchospora aurea* Vahl, Enum. Pl. 2: 229. 1806 ; Clarke in Hook. f., F.B.I. 6 : 671. 1893.

Robust herbs, 60 - 90 cm tall, without stolons. Leaves 30 - 60 x 1.2 - 2.5 cm. Panicles *ca* 30 cm, composed of 3 - 5 corymbs. Spikelets usually solitary, reddish brown, *ca* 8 mm long. Perianth-bristles 6, equalling nuts in length. Nuts *ca* 2.5 mm, smooth, reticulated or transversely wavy ; beak conical, 4 - 5 mm long, grooved, pale greenish.

Sikkim, s.l. (*fide* Hook. f. 1893).

Fl. & Fr. : June - Nov.

R. rugosa (Vahl) Gale in Rhodora 46 : 275, t. 835. f. 1 a & b. 1944.
Schoenus rugosus Vahl, Ecolog. Amer. 5. 1798. *Rhynchospora glauca*
sensu Clarke in Hook. f., F.B.I. 6: 671. 1893, *non* Vahl 1806. *Rhynchospora*
brownii R. & S., Syst. Veg. 2 : 86. 1817. *Rhynchospora rugosa* ssp.
brownii (R. & S) T. Koyama in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 118.
1978.

var. **rugosa**

Perennial herbs, up to 80 cm tall. Leaves canaliculate, rigid. Panicles narrow, consisting of 3 - 4 corymbiform anthelas. Spikelets ovate - lanceolate, ferruginous. Nuts subsessile, obovoid.

Sikkim, *s.l.* ; 1000 - 1500 m.

Fl. & Fr. : June - Aug.

var. **griffithii** (Boeck.) Verma & Venna Chandra in Bull. Bot. Surv. India 22 : 128. (1980) 1982. *Rhynchospora griffithii* Boeck. in Linnaea 38 : 404. 1874 ; Clarke in Hook. f., F.B.I. 6 : 672. 1893. *Rhynchospora griffithii* var. *levisetis* Clarke in Hook. f., l.c. 6: 672. 1893.

Perianth-bristles twice longer than the nut-beak. Nuts narrowly oblong, pale brown.

Lachen, Lachung ; *ca* 3000 m.

Fl. & Fr. : July - Nov.

var. **sikkimensis** (Clarke) Verma & Veena Chandra in Bull. Bot. Surv. India 22 : 130. 1980. *Rhynchospora sikkimensis* Clarke in Hook. f., F.B.I. 6 : 672. 1893.

Herbs. Panicles narrow, spikelets many, bearing 3 - 6 nuts. Perianth bristles 6, scarious, twice longer than nuts. Nuts narrowly obovoid, truncate ; beak longer than the nuts.

Khetchipheri Lake ; 2400 m.

Fl. & Fr. : July - Oct.

SCHOENOPLECTUS (Reichb.) Palla

Schoenoplectus grossus (L.f.) Palla in Allg. Bot. Z. 17 : Biebl. 3. 1911. *Scirpus grossus* L.f., Suppl. 104. 1781 ; Clarke in Hook. f., F.B.I. 6 : 659. 1893. *Scirpus kysoor* Roxb., Fl. Ind. 1 : 230. 1832. *Scirpus grossus* var. *kysoor* (Roxb.) Clarke in Hook. f., l.c. 660.

Perennials, up to 2 m tall. Stems triquetrous below, 3-winged upwards. Leaves clustered at base. Anthela terminal. Spikelets ovoid-oblong. Nuts oblong-obvoid, triquetrous, *ca* 1.2 mm long.

Sikkim, s.l. ; up to 600 m.

S. juncoidea (Roxb.) Palla in Bot. Jahrb. Syst. 10 : 299. 1888.
Scirpus juncoides Roxb., Fl. Ind. 1 : 218. 1820. *Scirpus erectus* sensu Clarke in Hook. f., F.B.I. 6 : 656. 1893, *non* Poir. 1804.

Herbs. Culms up to 80 cm tall, *ca* 3 mm thick, sheathed at base. Leaves lacking. Inflorescence pseudolateral, head with 1 - 9 spikelets.

Lachen, Phodong, between Phodong - Kabi ; 2400 m.

Fl. & Fr. : March - June.

S. mucronatus (L.) Palla in Verh. K.K. Zool. Bot. Ges. Wien 38 : 657. 1893. *Scirpus mucronatus* L., Sp. Pl. 50. 1753. *Scirpus triangulatus* Roxb., Fl. Ind. 1 : 219. 1820. *Schoenoplectus triangulatus* (Roxb.) Sojak in Cas. Nar. Muz. (Prague) 141 : 62. 1972.

Perennial herbs with creeping rhizomes. Culms up to 100 cm tall. Leaves lacking. Inflorescences pseudolateral, usually capitate.

Near Kabi, Gangtok, Kabi - Phodong, Lachen, Yoksum ; 2400 - 2700 m.

Fl. & Fr. : March - June.

S. supinus (L.) Palla in Bot. Jahrb. 10 : 299. 1889 ; J. Raynal in Adansonia 2, 16 : 145. 1976. *Scirpus supinus* L., Sp. Pl. 49. 1753 ; Clarke in Hook. f., F.B.I. 6 : 655. 1893.

Herbs. Stems sessile. Spikelets in a single lateral head. Nuts obovoid, triquetrous.

Sikkim, s.l. ; up to 900 m. .

SCIRPUS L.

Scirpus ternatanus Reinw. ex Miq., Fl. Ind. Bat. 3 : 307. 1856 ('ternatensis'). *Scirpus chinensis* Munro in Seem., Bot. Voy. Herald 423. 1857 ; Clarke in Hook. f., F.B.I. 6 : 662. 1893.

Perennial herbs. Stems terete below, trigonous above. Leaves scattered all along the stem. Anthela decomound, spikelets in clusters of 4 - 15, ovoid - oblong ; glumes ovate to suborbicular, brown. Nuts broadly obovoid.

Nampok, Lachen, Lachung, Lachen ; 1800 m.

Fl. & Fr. : April - June.

SCLERIA P. Berg.

Scleria levis Retz., Obs. Bot. 4 : 13. 1786. *Scleria hebecarpa* Nees in Wight, Contrib. Bot. Ind. 117. 1834 ; Clarke in Hook. f., F.B.I. 6 : 689. 1894. *Scleria pubescens* Steud., Syn. Pl. Glum. 2 : 168. 1855. *Scleria hebecarpa* var. *pubescens* (Steud.) Clarke in Hook. f., l.c.

Perennials, up to 1 m tall. Stem triquetrous. Leaves 80 x 0.8 cm. Inflorescence paniculate, of a terminal and 1 - 2 lateral partial panicles, spikelets unisexual, pale brown. Nuts subglobose, apiculate, glossy white (rarely bluish black).

Teesta & Rangeet valleys ; 300 - 900 m.

Fl. & Fr. : April - Oct.

S. lithosperma (L.) Swartz, Prodr. 18. 1788 ; Clarke in Hook. f., F.B.I. 6 : 685. 1894. *Scirpus lithospermus* L., Sp. Pl. 51. 1753.

Perennials, up to 70 cm tall. Leaves scattered on stem, aggregated in middle, 1 - 5 mm wide. Panicles up to 30 cm long, narrow, of 1 lateral and 2 - 3 lateral partial panicles. Spikelets bisexual, solitary or 2 - 3 together. Nuts oblong-ovoid, trigonous, glossy-ferrugineous.

Sikkim, s.l. ; up to 900 m.

Fl. & Fr. : July - Aug.

S. parvula Steud., Syn. Pl. Glum. 2 : 174. 1855 ; Kern in Blumea 11 : 202. 1961. *Scleria tessellata* sensu Clarke in Hook. f., F.B.I. 6 : 686. 1894, p.p. non Willd. 1805.

Annual, tufted herbs, up to 1 m tall. Leaves up to 50 x 0.5 cm. Panicles up to 4 cm long, of 1 terminal and 1 - 3 lateral panicles. Spikelets unisexual. Nuts obovoid, apiculate, glossy-white.

Lachung ; ca 2700 m.

Fl. & Fr. : May - Oct.

S. terrestris (L.) Fassett in Rhodora 26 : 159. 1924. *Zizania terrestris* L., Sp. Pl. 991. 1753. *Scleria elata* Thw., Enum. Pl. Zeyl. 353. 1864 ; Clarke in Hook. f., l.c. 690. *S. elata* var. *decolorans* Clarke in Hook. f., l.c. 690. *S. elata* var. *latior* Clarke in Hook. f., l.c. 690.

Perennial herbs, up to 1 m tall. Leaves up to 75 x 0.3 cms. Panicles up to 15 cm long, of 1 terminal and 2 - 5 lateral partial panicles. Spikelets unisexual, stramineous to purplish. Nuts ovoid, apiculate, glossy-white, blackish on ripening

Rangpo, Dzongu ; up to 1800 m.

Fl. & Fr. : July - Dec.

POACEAE

Subfamily : BAMBUSOIDEAE

(S. Kumar)

BAMBUSA Schreb. *nom. cons.*

Bambusa arundinacea (Retz.) Willd., Sp. Pl. 2 : 245. 1799 ; Gamble in Hook. f., F.B.I. 7 : 395. 1896. *Bambos arundinacea* Retz., Obs. Bot. 5 : 24. 1789.

Tall, thorny, perennials. Branches bright green, shiny, curved, with 2 - 3 recurved spines on nodes ; culms large, broad ; sheaths thickly clothed with golden hairs. Leaves linear-lanceolate, with swollen petioles.

Cultivated in the lower hills (*cf.* Varmah & Bahadur 1980).

B. paflida Munro in Trans. Linn. Soc. 26 : 97. 1868 ; Gamble in Hook. f., F.B.I. 7 : 389. 1896. PUSHEE (Lep.) ; BONGSHING (Bhut.).

Culms olive green. Nodes not so conspicuous ; nodes and internodes thin-walled. Culm-sheaths truncate at top. Leaves glabrous above, white-hirsute beneath, subulate tipped.

Teesta valley, Great Rangeet valley, Pedong ; 1500 - 1800 m.

B. tulda Roxb., Fl. Ind. 2 : 193. 1832 ; Gamble in Hook. f., F.B.I. 7 : 387. 1896. *B. teres* Buch.-Ham. ex Munro in Trans. Linn. Soc. 26 : 95. 1868. *B. nutans* Wall. ex Munro, l.c. 92 ; Hook. f., l.c. 387. *B. longispicula* Gamble in Brandis, Indian Trees 668. 1906. PAOSHIDING YING (Lep.) ; MALLOO, MAHLA (Nep.).

Culms straight, nodes not swollen ; internodes thick-walled with a white ring below nodes ; culm-sheaths dorsally clothed with black hairs, auricled at base, auricles with dense bristles.

Notes : Cultivated up to 1500 m for ornamental purposes. It fetches good prices to growers.

CHIMONOBAMBUSA Makino

Chimonobambusa hookeriana (Munro) Nakai in J. Arn. Arb. 6 : 151. 1925. *Arundinaria hookeriana* Munro in Trans. Linn. Soc. London 26 : 29. 1868 ; Gamble in Hook. f., F.B.I. 7 : 382. 1896. PRONG, PRAONG (Lep.) ; SINGHANI (Nep.).

Culms slender, caespitose, glaucous green ; nodes ringed by the base of sheath ; internodes thick-walled with a dark-bluish ring at top. Leaves linear.

Yoksum, Chungthang ; 1800 - 2200 m.

Notes : The parent plant dies after maturity of seeds and formation of suckers from the root.

DENDROCALAMUS Nees

Dendrocalamus hamiltonii Nees & Arn. ex Munro in Trans. Linn. Soc. 26 : 151. 1868 ; Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 84. t. 74. 1896 *et* in Hook. f., F.B.I. 7 : 405. 1896. PAO (Lep.) ; TAMA (Nep.).

Culms with many branches at angles, curved downwards, young culms appressedly pubescent ; nodes with scars ; internodes thick-walled ; culm - sheaths glabrous, truncate at apex. Panicles furrowed on one side, bearing half verticillate semiglobular purple heads.

Yoksum, Rate - chu, Gangtok ; 1000 - 2000 m.

Notes : Widely used for matwork, building and baskets.

D. hookeri Munro in Trans. Linn. Soc. 26 : 151. 1868 ; Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 83. t. 73. 1896 *et* in Hook. f., F.B.I. 7 : 405. 1896. PATU (Lep.) ; TILI BANS (Nep.).

Culms caespitose with long curved branches, internodes dark green, thick-walled ; culm sheaths densely covered with black brown hairs outside. Leaves oblong - lanceolate, cuspidate, with prominent yellow midrib. Panicles large, bearing globular heads.

Notes : Cultivated at 600 - 1500 m for making baskets and water-buckets.

D. patallaris Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 86. t. 75. 1896 *et* in Hook. f., F.B.I. 7 : 406. 1896. PAGJIOK, PAGJIOK - PAO (Lep.).

Culms striate, dull green ; nodes ringed ; internodes thin-walled ; culm-sheaths long cylindric, sparsely strigose, with wings and fringed margins. Panicles compound, bearing distant distichous heads.

Sikkim, *s.l.* ; *ca* 1200 m.

Notes : Ornamental.

D. sikkimensis Gamble in Hook., Ic. Pl. t. 1770. 1888 *et* in Ann. Roy. Bot. Gard. Calc. 7 : 82. t. 72. 1896 *et* in Hook. f., F.B.I. 7 : 405. 1896. PUGRIANG (Lep.).

Culms caespitose, naked below, dark green ; nodes ringed, internodes rough ; culm-sheaths densely covered with a dark golden brown felt. Panicles large, bearing red-brown, globose heads.

Sikkim, *s.l.* ; 1200 - 1800 m.

Notes : It is the largest, beautiful bamboo of Sikkim, used for making 'Chunga' for carrying water or milk by local inhabitants.

DREPANOSTACHYUM Keng f.

Drepanostachyum intermedium (Munro) Keng. f. in J. Bomb. Res. 2 : 28. 1983. *Arundinaria intermedia* Munro in Trans. Linn. Soc. 16 : 28. 1868 ; Gamble, Ann. Roy. Bot. Gard. Calc. 7 : 15. 1896 *et* in Hook. f., F.B.I. 7 : 381. 1896. *Chimonobambusa intermedia* (Munro) Nakai in J. Arn. Arb. 6 : 151. 1925. PARMIOK (Lep.) ; NIGALO, TITI NIGALA (Nep.).

Small shrubs. Culms slender, greyish green ; nodes swollen with a prominent ring and white line below ; culm-sheaths glabrous, striate, ciliate below. Leaves linear-oblong-lanceolate. Panicles whorled ; racemose, borne on leafless culms.

Sikkim *s. l.* ; 1200 - 2100 m.

D. gracilis (Blanchard) S. Kumar *comb. nov.* *Arundinaria gracilis* Blanchard in Revne Horticole 490. 1886 ; Camus, Les. Bambusees 38. 1912 ; Varmah & Bahadur in Ind. For. Rec. 6 : 1. 1980.

Small caespitose shrubs. Culms green. Branches turning yellow at maturity. Culm-sheath with an imperfect blade and divaricate panicle.

Sikkim *s.l.* (*cf.* Varmah & Bahadur 1980).

D. jainianum (Das & Pal) R. Majumdar in Bull. Bot. Surv. India 25 : 235. t. 1. (1983) 1895. *Chimonobambusa jainiana* Das & Pal in J. Econ. Tax. Bot. 4 : 1023 - 24. t. 1. 1983.

Culms caespitose, light brown, striate with tubercle based hairs. Nodes with a peltate hairy disc. Leaves oblong-lanceolate, acuminate, hairy along prominent midrib on lower surface. Panicles loose, drooping.

Sikkim, *s.l.*

Fl. & Fr. : March - April.

D. polystachyum (Kurz ex Gamble) R. Majumdar in Fl. Ind. Enum. Monocot. 277. 1989. *Arundinaria polystachya* Kurz ex Gamble in Ann. Roy. Bot. Gard. Calc. 7 : t. 5. 1896 *et* in Hook. f., F.B.I. 7 : 378. 1896. *Chimonobambusa polystachya* (Kurz ex Gamble) Nakai in J. Arn. Arb. 6 : 151. 1925.

Small hairy shrubs. Culms soft ; culm - sheaths lacking. Leaves lanceolate, acuminate, tip twisted. Panicles large, terminal or axillary.

Sikkim, s.l. ; 900 - 1500 m.

D. suberectum (Munro) R. Majumdar in Bull. Bot. Surv. India 25 : 236. 1985. *Arndinaria suberecta* Munro in Trans. Linn. Soc. London 26 : 32. 1868 ; Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 25. t. 24. 1896 *et* in Hook. f., F.B.I. 7 : 385. 1896.

Culms caespitose, olive green. Nodes smooth, slightly swollen. Internodes thick walled. Branchlets whorled. Culm-sheaths recurved, subulate. Leaves linear, subsessile.

Sikkim, s.l.

HIMALAYACALAMUS Keng f.

Himalayacalamus falconeri (Hook. f. ex Munro) Keng f. in J. Bomb. Res. 2 : 24. 1983. *Thamnocalamus falconeri* Hook. f. ex Munro in Trans. Linn. Soc. 26 : 34. 1868. *Aurndinaria falconeri* (Hook. f. ex Munro) Benth. in Benth. & Hook., Gen. Pl. 3 : 1208. 1883 ; Gamble in Ann. Roy. Bot. Gard. Calc. 1896 *et* in Hook. f., F.B.I. 7 : 383. 1896. *Drepanostachyum falconeri* (Hook. f. ex Munro) J. Campbell, Gen. Himal. Bamb. 23. 1983.

Tall shrubs. Young culms smooth, old ones scurvy white with yellow green stripes ; nodes swollen. Culm-sheaths oblong, truncate tipped, with persistent base. Leaves soft, oblong-lanceolate, glabrous. Panicles leafy.

Sikkim, s.l. ; 1900 - 2700 m.

MELOCALAMUS Benth.

Melocalamus indicus R. Majumdar in Bull. Bot. Surv. India 25 : 236. t. 1. (1983) 1985.

Evergreen, scandent bamboo. Culms almost solid spreading and arching over tall trees, hanging downwards with tufts of green leaves and branched tips ; nodes with a woody ring ; internodes white scurfy. Culm-sheaths crustaceous, falling at maturity. Panicles with globular culsters of spikelets.

Sikkim, s.l.

PHYLLOSTACHYS Sieb. & Zucc.

Phyllostachys assamica Gamble ex Brandis, Indian Trees 667. 1906. *P. bambusoides* Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 27, t. 27. 1896 *et* in Hook. f., F.B.I. 7 : 386. 1896, *non* Sieb. & Zucc.

Culms caespitose, yellow, flattened on one side. Branches semi - verticillate. Culm-sheaths thin, papery, striate, rounded above into a short neck. Leaves having black hairy petiole, glaucous beneath with white stiff hairs below.

Gangtok.

SCHIZOSTACHYUM Nees

Schizostachyum munroi S. Kumar & P. Singh in J. Ind. Bot. Soc. 1992. *Cephalostachyum capitatum* Munro in Trans. Linn. Soc. 26 : 139. 1868. *Schizostachyum capitatum* (Munro) R. Majumdar in Karthik. *et al.*, Fl. Ind. Enum. Monocot. 281. 1989, *non* Rupr. 1840. PAYONG (Lep.) ; GOBIA, GOPE (Nep.).

Semiscandent shrubs. Culm-sheaths pale brown, appressed pubescent. Leaves ovate-lanceolate. Inflorescence dense terminal or axillary, globular heads. Caryopsis beaked, brownish, shiny.

Sikkim, s.l. ; 600 - 2400 m.

S. polymorphum (Munro) R. Majumdar in Karthik. *et al.*, Fl. Ind. Enum. Monocot. 242. 1989. *Pseudostachyum polymorphum* Munro in Trans. Linn. Soc. 26 : 142. t. 1. 1868 ; Gamble in Ann. Roy. Bot. Gard. Calc. 7 : *et* in Hook. f., F.B.I. 7 : 409. 1896. PARPHIOK, PAPHOK (Lep.) ; FILING (Nep.).

Shrubs with creeping rhizomes. Culms subscandent ; nodes simple ; internodes thin-walled ; culm-sheaths truncate with strigose brownish hairs and very small auricles with a tuft of bristles. Panicles large with wiry angled rachis. Caryopsis globose, crustaceous.

Sikkim, s.l.

S. sharmae S. Kumar & P. Singh in J. Ind. Bot. Soc. 1992. *Schizostachyum latifolium* (Munro) R. Majumdar in Karthik. *et al.*, Fl. Ind. Enum. Monocot. 281. 1989, *non* Gamble. *Cephalostachyum latifolium* Munro in Trans. Linn. Soc. 26 : 140. 1868 ; Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 106. 1898 *et* in Hook. f., F.B.I. 7 : 412. 1896.

Semiscandent shrubs with thin darkgreen, rough, culms having a whitish ring below nodes. Culm-sheaths thin, papery, straw-coloured with

rounded tip, auricled. Leaves unequal at base. Inflorescence a terminal globular head composed of clusteres of spikelets. Caryopsis broadly ovoid, stalked, curved above.

Sikkim, s.l. (*cf.* Varmah & Bahadur 1980).

THAMNOCLAMUS Munro

Thamnochalamus aristatus (Gamble) E.G. Camus, Les. Bambus 54. 1913. *Arundinaria aristata* Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 18. t. 17. 1896 *et in* Hook. f., Fl. Brit. Ind. 7 : 382. 1896. BHESHA (Bhut.) ; BABAIN (Lep.) ; RATONIGALA (Nep.).

Shrubs with glabrous yellow culms. Branchlets reddish. Nodes prominent. Culm-sheaths loose, glabrous with ciliate margin and few stiff bristles at mouth, fascicled. Leaves on joints of branchlets. Inflorescence drooping.

Sikkim, s.l. ; 2200 - 3300 m.

YUSHANIA P.C. Keng

Yushania maling (Gamble) R. Majumdar in Fl. Ind. Enum. Monocot. 283. 1989. *Arundinaria maling* Gamble in Kew Bull. 1912 ; 139. 1912. *Fargesia maling* (Gamble) J. Campbell, Gen. Himal. Bamb. 40. 18 1985. PHIUNG, MHEEM (Bhut.); PHYEUM MIKNU (Lep.) ; MALING, MALINGO, KAORE-MALING, PARENT (Nep.).

Gregarious shrubs. Culms thick walled, hollow, indistinctly ribbed, green, with a ring of long bristles at nodes. Culm-sheaths straw-coloured, prominently striate, with ciliate margins, covered with brown haris on the outer surface and auricled on both sides. Caryopsis oblong, dorsally furrowed.

Sikkim, s.l. ; 1800 - 3600 m.

Y. microphylla (Griff.) R. Majumdar in Fl. Ind. Enum. Monocot. 283. 1989. *Bambusa microphylla* Griff., Journ. 1 : 242. 259. 1847. *Arundinaria microphylla* (Griff.) Munro in Trans. Linn. Soc. 26 : 32. 1868 ; Gamble in Ann. Roy. Bot. Gard. Calc. 7 : 22. t. 118. 1896 *et in* Hook. f., F.B.I. 7 : 383. 1896.

Gregarious caespitose shrubs with densely tufted culms. Internodes very long, glabrous. Nodes prominent. Leaves linear-lanceolate.

Sikkim, s.l. ; 2400 - 3100 m.

Y. pantlingii (Gamble) R. Majumdar in Fl. Ind. Enum. Monocot. 283. 1989. *Arundinaria pantlingii* Gamble in Ann. Roy. Bot. Gard. Calc.

7 : 129. t. 118. 1896 *et in* Hook. f., F.B.I. 7 : 380. 1896. *Semiarundinaria pantlingii* (Gamble) Nakai in J. Arn. Arb. 6 : 151. 1925. *Fargesia pantlingii* (Gamble) J. Campbell, Gen. Himal. Bamb. 41. 1985.

Culms smooth. Nodes with a hairy ring. Culm-sheaths smooth, striate, auricled. Panicles on leafless branches. Caryopsis oblong, grooved, dark brown.

Chunghang, Lachung ; 1800 - 2300 m.

Y. racemosa (Munro) R. Majumdar in Karthik. *et al.*, Fl. Ind. Enum. Monocot. 283. 1989. *Arundinaria racemosa* Munro in Trans. Linn. Soc. 26 : 17. 1868 *emend.* Gamble in Kew Bull. 1912. 138. 1912 ; Gamble in Ann. Roy. Bot. Gard. Calcutta 7 : 9, t. 8. 1896 *et in* Hook. f., F.B.I. 7 : 379. 1896. *Fargesia racemosa* (Munro) Yi in J. Bomb. Res. 2 : 39. 1983.

Gregarious shrubs. Culms thin-walled, erect, with a ring on nodes ; internodes bluish green, scabrous. Culm-sheaths striate, covered with scattered black hairs. Leaves linear-lanceolate with setaceous tip.

Dzongri, Chola valley ; 3000 - 3600 m.

Subfam. : PANICOIDEAE

(S. Kumar)

ALLOTEROPSIS C.B. Presl.

Alloteropsis semialata (R. Br.) Hitch. in Contrib. U.S. Nat. Herb. 12 : 210. 1909 ; Bor, Grass. Ind. 276. 1960. *Panicum semialatum* R. Br., Prodr. 192. 1810. *Axonopus semialatus* (R. Br.) Hook. f., F.B.I. 7: 64. 1896.

Small grass. Culms densely tufted. Leaf blades linear, finely acuminate. Spikes subdigitate. Spikelets pale brown, awned.

Sikkim, s.l. ; up to 1950 m.

Fl. & Fr. : May - Sept.

APLUDA L.

Apluda mutica L., Sp. Pl. 82. 1753 ; Bor, l. c. 93. *Apluda varia* Hara in DC., Monogr. Phan. 196. 1889 ; Hook. f., F.B.I. 7: 150. 1896.

Perennial grass. Leaf-blades linear. Panicles false, with single node ; racemes comprising of 1 sessile and 2 awnless pedicelled spikelets enclosed in a spathe.

Jorethang, Teesta valley, Rangpo ; 800 - 2500 m.

Fl. & Fr. : Aug. - Nov.

APOCOPSIS Nees

Apocopsis palaeacea (Trin.) Hochr. in Bull. N. Y. Bot. Gard. 6 : 262. 1910 ; Bor, l. c. 96. *Ischaemum palaeaceum* Trin. in Mem. Acad. Sci. Peters. ser. 6, 2 : 293. 1832. *Apocopsis royleanus* Nees in Ann. Mag. Nat. Hist. 7 : 142. 1841 ; Hook. f., F.B.I. 7 : 142. 1896.

Perennial, erect grass. Leaf blades narrowly lanceolate, acuminate. Peduncles long, slender. Spikelets 1 - 2, with rusty villous rachis. Pedicelled spikelets represented by pedicels only.

Sikkim, s. l. ; 1700 m.

Fl. & Fr. : July - Dec.

ARTHRAXON P. Beauv.

Arthraxon castratus (Griff.) Narayanswami ex Bor, Fl. Assam 5 : 376. 1940 et Grass. Ind. 99. 1960. *Andropogon castratus* Griff., Notul. 3 : 89. 1851. *Anthraxon rufus* (Steud.) Hochst. in Flora 39 : 188. 1856 ; Hook. f., F.B.I. 7 : 144. 1896.

Perennial erect grass. Leaf-blades oblong or ovate-lanceolate, with cordate amplexicaul base. Peduncle of inflorescence densely whitish pubescent. Racemes with shortly ciliate joints. Sessile spikelets compressed.

Reshi ; 500 m.

Fl. & Fr. : Dec.

A. hispidus (Thunb.) Makino in Bot. Mag. Tokyo 26 : 214. 1912 ; Bor, Grass. Ind. 99. 1960. *Phalaris hispida* Thunb., Fl. Jap. 44. 1784. *A. ciliaris* P. Beauv., Ess. Agrost. 111. 152. t. 11, f. 6. 1812 ; Hook. f., F.B.I. 7 : 146. 1896. *A. nudus* (Steud.) Hochst. in Flora 39 : 188. 1856 ; Bor, l. c. ; Hook. f., l. c. 101. *A. submuticus* (Steud.) Bor, l. c. 144 ; Bor, l. c. 102. *A. breviaristatus* Hack. in DC., Mongr. Phan. 6 : 350. 1889 ; Hook. f., l. c. ; Bor., l. c. 99. *A. inermis* Hook. f., l. c. ; Bor, l. c. 100. *A. quartinianus* (A. Rich.) Nash. in N. Amer. Flora 17 : 99. 1912 ; Bor, l. c. 102. *A. hookeri* (Hack.) Henr. in Blumea 4 : 526. 1941 ; Bor, l. c. 99.

Annual grass with densely tufted trailing culms. Leaf-blades sparsely hairy. Spikelets, solitary, lanceolate, without a lemma in lower floret. Pedicelled spikelet represented by a vestigeal lower glume.

Gangtok, Richingpong to Gaysing, Little Rangeet, up to 1900 m.

Fl. & Fr. : June - Sept.

A. lancifolius (Trin.) Hochst. in Flora 39 : 188. 1856 ; Bor, *l. c.* 100. 1960. *Andropogon lancifolius* Trin. in Mem. Acad. Sci. Petersb. 62 : 171. 1832. *Anthraxon microphyllus sensu* Hook. *f.*, F.B.I. 7 : 146. 1896 *p. p.*, *non* (Trin.) Hochst. 1856.

Annual grass with decumbent slender culms. Leaf-blades with a cordate, amplexicaul base. Paired spikelets only in upper part of inflorescence ; lower glume of sessile spikelets with indistinct nerves.

Gangtok, Penengla, Yak - la, Richingpong to Gyalsing ; 900 - 1800 m.

A. microphyllus (Trin.) Hochst. in Flora 39 : 188. 1856 ; Hook. *f.*, F.B.I. 7 : 147. 1896. *p. p.* ; Bor, *l.c.* ed. 2. 1. 1973. *Andropogon microphyllus* Trin. in Mem. Acad. Sci. St. Petersb. 6, 2 : 275. 1833. *Anthraxon sikkimensis* Bor in Kew Bull. 1951 : 447. 1952 *et Grass. Ind.* 102. 1960.

Annual grass. Culms slender, glabrous ; nodes pubescent. Leaf-blades with hairy margins. Peduncles gilbarous. Spikelets paired throughout.

Lachung ; 2000 - 3000 m.

Fl. & Fr. : July - Oct.

BOTHRIOCOCHLOA O. Kuntze

Bothriochloa bladhii (Retz.) S. T. Blake in Proc. R. Soc. Queensland 80 : 62. 1969. *Andropogon bladhii* Retz., Obs. Bot. 2 : 27. 1781. *Bothriochloa intermedia* (R. Br.) A. Camus in Ann. Soc. Linn. Lyon. 1930. n. s. 76 : 164. 1931 ; Bor, Grass. Ind. 108. 1960. *B. glabra* (Roxb.) A. Camus, *l.c.* ; Bor, *l.c.* ; 107. *B. odorta* (Donna Lisboa) A. Camus, *l.c.* ; Bor, *l.c.* 109. *Andropogon intermedium* R. Br., Prodr. 202. 1810 ; Hook. *f.*, F.B.I. 7 : 175. 1896. *A. odoratus* Donna Lisboa in J. Bombay Nat. Hist. Soc. 4 : 123. 1889. *Dichanthium bladhii* (Retz.) W. D. Clayton in Kew Bull. 32 : 3. 1977. *D. odoratum* (Donna Lisboa) Jain & Deshpande in Bull. Bot. Surv. Ind. 20 : 134. 1979.

Tufted perennials with tall robust culms. Leaf-blades linear. Racemes in irregular whorls. Peduncle simple or branched. Rachis pubescent. Pedicelled spikelets usually pitted. Sessile spikelets narrowly elliptic.

Gangtok ; 500 - 1800 m.

Fl. & Fr. : Aug. - Jan.

B. ischaemum (L.) Keng in Cotrib. Biol. Lab. Sci. Soc. China Bot. 10 : 201. 1936 ; Bor, l.c. 108. 1960. *Andropogon ischaemum* L., Sp. Pl. 1047. 1753 ; Hook. f., F.B.I. 7 : 171. 1896.

Tufted perennials with mostly basal leaves. Inflorescence in subdigitate racemes. Spikelets oblong-lanceolate, sessile.

Sikkim s.l.

BRACHIARIA Griseb.

Brachiaria ramosa (L.) Stapf in Prain, Fl. Trop. Afr. 9 : 542. 1919 ; Bor, l.c. 284. 1960. *Panicum ramosum* L., Mant. 1 : 29. 1767 ; Hook. f., F.B.I. 7 : 36. 1896.

Annuals. Leaf-blades broadly linear. Rachis triquetrous. Spikelets adaxial. Lower glumes 5 - 7 - nerved.

Singtam ; up to 1800 m.

Fl. & Fr. : July - Oct.

B. reptans (L.) Gard. et Hubb. in Hook., Ic. Pl. sub. t. 3363. 1938 ; Bor, l.c. 285. 1960. *Panicum reptans* L., Syst. Nat. ed. 10. 870. 1759. *Panicum prostratum* Lam., Ill. 1 : 171. 1791 ; Hook. f., F.B.I. 7 : 33. 1896.

Annuals. Culms decumbent, rooting at nodes. Leaf-blades glabrous, ovate-lanceolate, with cordate amplexicaul base and thickened margins. Spikelets glabrous, paired. Lower glume truncate. Caryopsis flattened, oval.

South Sikkim District ; 200 - 1800 m.

Fl. & Fr. : June - Oct.

Notes : A good fodder grass with edible grains.

B. villosa (Lam.) A. Camus in Lecomte, Fl. Gen. de l' Indo - Chine 7 : 433. 1922 ; Bor, l.c. 286. 1960. *Panicum villosum* Lam., Ill. 1 : 173. 1791, p.p. ; Hook. f., F.B.I. 7 : 34. 1896, p.p.

Annuals. Culms rooting at base, shortly hairy below. Leaves pubescent or villous at both the surfaces. Spikelets solitary, pubescent. with lower glumes acute or obtuse but not truncate.

Sikkim, s. l. ; up to 2400 m.

Fl. & Fr. : July - Sept.

CAPILLIPEDIUM Stapf

Capillipedium assimile (Steud.) A. Camus in Lecomte, Fl. Gen. de l' Indo - Chine 7 : 314. 1922 ; Bor, Grass. Ind. 110. 1960. *Andropogon assimilis* Steud. in Zoll. Syst. Verz. 58. 1854 ; Hook. f., F. B. I. 7 : 179. 1896. *Dichanthium assimile* (Steud.) Deshpande in Fasc. Fl. India 15 : 6. 1894.

Perennials. Culms decumbent, nodes glabrous. Leaflets hispidulus, linear-lanceolate. Panicles branched, bearded in the axis. Racemes jointed, filiform. Sessile spikelets shorter than pedicelled spikelets.

Lower Burtuk Basti, Baluakhani, on way to Rongli, Jorethang, Deorali ; 600 - 2100 m.

Fl. & Fr. : Sept. - Dec.

C. parviflorum (R. Br.) Stapf in Prain, Fl. Trop. Afr. 9 : 169. 1917 ; Bor. l.c. 112. 1960. *Holous parviflorus* R. Br., Prodr. 199. 1810. *Andropogon micranthus* Kunth, Rev. Gram. 1 : 165. 1829 ; Hook. f., F. B. I. 7 : 178. 1896. *Dichanthium parviflorum* (R. Br.) DC. in Amer. J. Bot. 54 : 386. 1967.

Rumtek, Baluakhani ; up to 2000 m.

Fl. & Fr. : Jan. - March.

CHYSOPOGON Trin.

Chrysopogon aciculatus (Retz.) Trin., Fund. Agrost. 188. 1822 ; Bor, l. c. 115. 1960. *Andropogon aciculatus* Retz., Obs. Bot. 5 : 22. 1789 ; Hook. f., F.B.I. 7 : 188. 1896.

Grass with decumbent culms. Leaves linear. Panicles erect with slender filiform branches. Spikelets compressed, sessile. Pedicels of pedicelled spikelets glabrous.

Ranipul, Rangpo, Molli, Chhuzachen ; up to 1700 m.

C. gryllus (L.) Trin., Fund. Agrost. 188. 1822 ; Bor, Grass. Ind. 117. 1960. *Andropogon gryllus* L., Cent. Pl. 2 : 33 : 1756 ; Hook. f., F.B.I. 7 : 187. 1896.

Grass wth erect culms. Panicles lax, branches whorled, densely bearded. Lower glume of sessile spikelet cartilaginous. Pedicelled spikelet linear-lanceolate.

Luing Basti ; 800 - 2900 m.

C. lancearius (Hook. f.) Haines, Bot. Bihar & Orissa 1036. 1924 (3 : 103. 1961 rep. ed.) ; Bor, l. c. 118. 1960. *Andropogon lancearius* Hook. f., F.B.I. 7 : 190. 1896.

Perennials with glabrous nodes. Leaflets lanceolate, acuminate, minutely spinulose at margins. Panicles bearing solitary, erect racemes. Pedicels of pedicelled spikelets hairy.

Sikkim, s.l. ; 500 - 800 m.

COIX L.

Coix lacryma-jobi L., Sp. Pl. 972. 1753 ; Hook. f., F.B.I. 7 : 100. 1896 ; Bor, l. c. 264. 1960.

var. *lacryma-jobi* SAONKRU (Hindi) ; GARGAR (Beng.).

Annuals. Culms shiny. Leaflets cordate at base, acuminate at apex, sheath broad. Female spikelets surrounded by a hardened bract forming head like structure. Male spikelets loosely imbricate on a slender rachis.

Burtuk, Riang, Tumlong ; 800 - 1500 m.

Fl. & Fr. : May - Oct.

var. **ma-yuen** (Romanet) Stapf ex Hook. f., F.B.I. 7 : 100. 1896 ; Bor, l.c. 265. 1960. *Coix ma-yuen* Romanet in Bull. Soc. Accl. Paris, 3, 8 : 442. 1881.

Annuals with soft striate involucre. Fruits crustaceous or chartaceous.

Sikkim s.l. ; up to 1500 m.

Fl. & Fr. : July - Sept.

Notes : It is commonly cultivated for preparation of a beer from the soft shelled fruits. The stem and leaves are used as fodder.

CYMBOPOGON Spr.

Cymbopogon flexuosus (Nees ex Steud.) Wats. in Atkins., Gaz. N. W. Prov. Ind. 392. 1881 ; Bor, Grass. Ind. 127. 1960. *Andropogon flexuosus* Nees ex Steud., Syn. Pl. Glum. 1 : 388. 1854. *A. nardus* L. var. *flexuosus* Hack. in DC., Monogr. Phan. 6 : 603. 1889 ; Hook. f., F.B.I. 7 : 207. 1896. *C. flexuosus* (Nees ex Steud.) Wats. var. *sikkimensis* Bor in J. Bombay Nat. Hist. Soc. 52 : 162. 1954 et l.c.

Aromatic perennials. Culms stout, glabrous. Leaves linear, glaucous, green with purple tinge, coarsely hairy on upper surface. Inflorescence nodding. Sessile spikelets oblong, pedicelled spikelets acuminate.

Ratepani, Ranipul, Rangpo ; 900 - 1100 m.

C. khasianus (Hack.) Stapf ex Bor, Ind. For. Rec. (Bot.) 1 : 3. 92. 1983 et Grass. Ind. 128. 1960 ; Soenarko in Reinwardtia 9 : 346. 1977. *Andropogon naruds* var. *khasianus* Hack. in DC., Mongr. Phan. 608. 1889 ; Hook. f., F.B.I. 7 : 206. 1896.

Perennials ; nodes tomentose ; internodes, shiny, glabrous. Leaves linear, acuminate at apex. Panicles decompound, with noded axis, elliptic acute spatheole. Sessile spikelets purplish with lower glume ungrooved. Pedicelled spikelets elliptic, acute.

Sikkim s. l. ; 100 - 1500 m.

C. martinii (Roxb.) Wats. in Atkins., Gaz. Ind. 10 : 392. 1882 ; Bor, l. c. 129. 1960. *Andropogon martinii* Roxb., Fl. Ind. 1 : 280. 1820. *A. schoenanthus* L. var. *martinii* (Roxb.) Hook. f., F.B.I. 7 : 204. 1896. *C. motia* Gupta in Proc. Ind. Acad. Sci. B. 71 : 92. 1970.

Tufted perennials with woody stout rootstock. Culms with swollen nodes. Leaves tapering to form a filiform tip, cordate at base. Pseudopanicles with long narrowly elliptic spatheole, green, turning reddish on maturity. Lower glume of sessile spikelets with V-shaped groove in lower half. Pedicelled spikelets obtuse.

Singtam ; 900 - 1200 m.

Fl. & Fr. : Sept. - Nov.

Notes : An important essential oil yielding grass.

C. microtheca (Hook.) A. Camus in Rev. Bot. Appl. 1 : 284. 1921 ; Bor, l. c. 129. 1960. *Andropogon microtheca* Hook. f., F.B.I. 7 : 208. 1896.

Perennials, with erect smooth culms. Leaves glabrous, sheaths auricled. Inflorescence glabrous, branched with spatheoles, dilated at tip, joints of racemes densely ciliate on edges. Spikelets smaller than other species.

Luing Basti, Nan Dunga, between Tarkhola and Rangpo, Teesta valley ; 400 - 1200 m.

C. pendulus (Nees ex Steud.) Wats. in Atkins., Gaz., N. W. Prov. Ind. 392. 1882 ; Bor, l. c. 131. 1960. *Andropogon pendulus* Nees ex Steud., Syn. Pl. Glum. 1 : 388. 1854. *A. nardus* var. *grandis* Hack. in DC., Monogr. Phan. 6 : 605. 1889, p. p. ; Hook. f., F.B.I. 7 : 206. 1896, p.p.

Robust perennials. Culms tall stout. Leaves linear, acuminate with a pleasant lemon odour. Panicles pendulous, decompound, subtended by spathes. Lower glume of sessile spikelets with a concave groove from base to apex.

Teesta valley.

CYRTOCOCCUM Stapf

Cyrtococcum accrescens (Trin.) Stapf in Hook., Ic. Pl. sub. t. 3096. 1922 ; Bor, l. c. 291. 1960. *Panicum accrescens* Trin., Sp. Gram. Ic. 1 : t. 88. 1828.

Perennials, with tall culms arising from a branching, decumbent base. Leaves linear-lanceolate, tapering to a sharp point. Panicle lax, glabrous, branches capillary. Spikelets distinctly gibbous ; tip of flower glume acute, upper glume boat shaped, pedicelled-spikelets with cupulate tip.

Ratong to Yuksom.

Fl. & Fr. : July - Oct.

C. longipes (Wt. & Arn. ex Hook. f.) A. Camus in Bull. Mus. Hist. Nat. Paris 27 : 1921 ; Bor, l. c. 291. 1960. *Panicum longipes* Wt. & Arn. ex Hook. f., F.B.I. 7 : 58. 1896.

Perennials. Culms glabrous. Leaves linearlanceolate, finely pubescent beneath. Panicle fastigiately branched ; branches capillary. Spikelets acuminate, 3-nerved.

Rishap. *Fl. & Fr.* : June - Nov.

C. oxyphyllum (Steud.) Stapf in Hook., Ic. Pl. sub. t. 3096. 1922 ; Bor, l. c. 291. 1960. *Panicum oxyphyllum* Steud., Syn. Pl. Glum. 1 : 65. 1854. *P. pilipes* Nees & Arn. ex Buese in Miq., Pl. Jungh. 3 : 376. 1854 ; Hook. f. F.B.I. 7 : 57. 1896.

Small perennials with decumbent creeping culms. Leaves oblong-lanceolate. Panicles lax, with pilose branches. Pedicels shorter than spikelets.

Sikkim, s. l.

Fl. & Fr. : June - Dec.

C. patens (L.) A. Camus in Bull. Mus. Hist. Nat. Paris 27 : 18. 1928 ; Bor, l. c. 292. 1960. *Panicum patens* L., Sp. Pl. 58. 1753 ; Hook. f., F.B.I. 7 : 57. 1896.

Perennials. Culms branched, glabrous. Leaves acuminate, with hispid margins. Panicle lax, flexuous, spreading. Spikelets orbicular; lower glume obtuse or subacute.

Sikkim, s. l.

DICHANTHIUM Willemet.

Dichanthium annulatum (Forssk.) Stapf in Prain, Fl. Trop. Africa 9 : 178. 1979 ; Bor, l. c. 133. 1960. *Andropogon annulatum* Forssk., Fl. Aegypt.-Arab. 173. 1775 ; Hook. f., F.B.I. 7 : 196. 1896.

Densely tufted perennials. Culms hairy, noded. Leaves linear, acuminate, hairy. Racemes subdigitate. Caryopsis elliptic, light brown.

Sikkim, s. l.; up to 1100 m.

Fl. & Fr.: Sept. - Dec.

D. caricosum (L.) A. Camus in Bull. Mus. Hist. Nat. Paris 27 : 549. 1921 ; Bor, Grass. Ind. 134. 1960. *Andropogon caricosus* L., Sp. Pl. ed. 2. 1480. 1763 ; Hook. f., F. B. I. 7 : 196. 1896.

Perennials. Culms glabrous, noded. Leaves linear-lanceolate. Racemes with solid joints and pedicels. Spikelets obovate, awned. Caryopsis linear or fusiform.

Sikkim, s. l.

Fl. & Fr.: July - Dec.

DIGITARIA Haller nom. cons.

Digitaria ciliaris (Retz.) Koel., Descr. Gram. 27. 1802. *Panicum ciliare* Retz., Obs. Bot. 4 : 16. 1786. *Digitaria biformis* Willd., Enum. Pl. Hort. Berol. 92. 1809 ; Bor, l. c. 299. 1960. *D. adescendens* (Kunth) Henr. in Blumea 1 : 92. 1934 ; Bor, l. c. 298. *Paspalum sanguinale* (L.) Lam. var. *ciliare* (Retz.) Hook. f., F.B.I. 7 : 15. 1896. *D. sanguinale* (L.) Lam. var. *rottleri* Hook. f., l. c.

Annual grass. Culms decumbent, rooting at nodes. Leaf-blades linear. Racemes digitate or whorled. Spikelets appressed lanceolate; lower glumes obscure, margins of nerves of lower smooth. Caryopsis ellipsoid, grey or light brown.

Melli, Rumtek ; up to 1500 m.

Fl. & Fr.: June - Nov.

D. compacta (Roth ex R. & S.) Weldk. in Blumea 21 : 71. 1973. *Paspalum compactum* Roth ex R. & S., Syst. Veg. 2 : 316. 1817 ; Hook. f., F.B.I. 7 : 12. 1896 ; Bor, Grass. Ind. 336. 1960.

Grass with decumbent leafy culms. Leaf-blades elliptic, lanceolate, sparsely hairy on both sides. Sheaths densely hairy ; hairs bulbous at base. Spikelets hemispheric, without a lower glume.

Moplong.

Fl. & Fr. : June - Dec.

D. cruciata (Nees ex Steud.) A. Camus in Lecomte, Fl. Gen. de l' Indo-Chine 7 : 399. 1922 ; Bor, Grass. Ind. 300. 1960. *Panicum cruciatum* Nees ex Steud., Syn. Pl. Glum. 1 : 30. 1854. *Paspalum sanguinale* Lam. var. *cruciatum* (Nees ex Steud.) Hook. f., F.B.I. 7 : 14. 1896.

Annual grass with decumbent culms creeping at base. Leaves linear-lanceolate, membranous. Ligule truncate. Racemes subdigitate. Caryopsis apiculate, protruding above lower lemmas.

Deorali ; 100 - 1500 m.

Fl. & Fr. : June - Dec.

D. radicosa (C. B. Presl.) Miq., Fl. Ind. Bat. 3 : 437. 1857. *Panicum radicosum* C. B. Presl., Rel. Hanek. 1 : 297. 1830. *D. tinorensis* (Kunth) Bal. in Morot., J. de. Bot. 4 : 138. 1890 ; Bor, l. c. 306. 1960. *Paspalum sanguinale* var. *debile* Hook. f., F.B.I. 7 : 16. 1896.

Annuals. Culms decumbent, rooting at nodes. Leaves linear, scabrous. Ligule truncate. Rachis winged. Caryopsis narrow, lanceolate, acute, yellowish to pale brown.

Rangpo ; up to 1500 m.

Fl. & Fr. : Sept. - Nov.

D. sanguinalis (L.) Socp., Fl. Carn. ed. 2, 1 : 52. 1772 ; Bor, l. c. 304. 1960. *Panicum sanguinale* L., Spl. Pl. 57. 1753. *Paspalum sanguinale* (L.) Lam., Ill. 1 : 176. 1791 ; Hook. f., F.B.I. 7 : 13. 1896.

Annuals with ascending culms. Leaves linear ; sheath basally hirsute. Racemes subdigitate. Spikelets narrowly elliptic to ovate ; lower lemma with distinct minute spines. Caryopsis lanceolate, grey or light brown.

Phodonchen ; up to 1800 m.

Fl. & Fr. : July. - Sept.

D. setigera Roth. apud R. & S., Syst. Veg. 2 : 474. 1817 ; Bor, l. c. 305. 1960. *Paspalum sanguinale* Lam. var. *extensum* Hook. f., F.B.I.

7 : 15. 1896. *P. sanguinale* var. *calliblepharata* (Henr.) Veldk. in Blumea 21 : 40. 1973.

Annual grass with creeping culms. Leaf-blades linear-lanceolate. Racemes subdigitate large. Caryopsis lanceolate, not apiculate, grey or yellowish brown.

Sikkim, s. l. ; up to 600 m.

Fl. & Fl. : July - Oct.

D. strocta Roth ex R. & S., Syst. Veg. 2 : 474. 1817 ; Bor, Grass. Ind. 305. 1960. *Paspalum coyleanum* Nees ex Hook. f., F.B.I. 7 : 18. 1896. *D. stricta* var. *glabrescens* Bor in Webbia 11 : 336. 1955 et l.c. *D. stricta* var. *denudata* (Link.) Hern., Digitaria 175. 1950 ; Bor, l.c. 306.

Perennial grass with erect, leafy, culms. Leaf-blades linear, acuminate. Racemes subdigitate on a slender peduncle ; rachis narrowly winged ; pedicels dilated at tip. Caryopsis elliptic, apiculate.

Sikkim, s. l. ; up to 2100 m.

Fl. & Fr. : Aug. - Oct.

DIMERIA R. Br.

Dimeria ornithopoda Trin., Fund. Agrost. 167.t. 14. 1820 ; Hook. f., F.B.I. 7 : 104. 1896 ; Bor, Grass. Ind. 142. 1960.

Annual grass with weak culms, bearded at nodes, Leaf-blades linear, setaceous. Ligule ciliate. Spikes with solitary, shortly pedicelled, spikelets. Caryopsis cylindrical.

Sikkim, s. l..

Fl. & Fr. : July - Dec..

ECHINOCHLOA P. Beauv.

Echinochloa colonum (L.) Link., Hort. Berol. 2 : 209. 1833 ; Bor, l.c. 308. 1960. *Panicum colonum* L., Syst. Nat. ed. 10, 2 : 870. 1759. Hook. f., F.B.I. 7 : 32. 1892.

Annuals with erect or ascending culms. Leaves marked with purple streaks, eligulate. Racemes 4 - rowed, with uncrowded spikelets. Caryopsis broadly elliptic, deciduous.

Sikkim, s. l. ; up to 1800 m.

Fl. & Fr. : May - June.

Notes : It is one of the finest fodder grass.

E. crus-galli (L.) P. Beauv., Ess. Agrost. 53. 161. 1812 ; Bor, l. c. 310. 1960. *Panicum crus-galli* L., Sp. Pl. 1 : 56. 1753 ; Hook. f., F.B.I. 7 : 30. 1896. *E. glabrescens* Munro ex Eggel., Ann. List. Grass Uganda 16. 1974 ; Bor, l.c. 311.

Annuals with erect or ascending culms. Leaves linear, eligulate. Racemes in 2 to several irregular rows. Spikelets ovate-elliptic, hispid. Caryopsis broadly elliptic

Lower Sichey Basti, Gangtok.

Fl. & Fr. : June - Oct.

Notes : Found in marshy places like rice fields etc. It is a good fodder grass.

E. grumentacea Link., Hort. Berol. 1 : 204. 1827 ; *Panicum curs - galli* var. *frumentaceum* Hook. f., F.B.I. 7 : 31. 1896.

Robust herbs with erect culms. Leaves broadly linear. Spikelets broadly elliptic to orbicular, pubescent to hispid ; lower lemma acute.

Sikkim, s.l.

Fl. & Fr. : Aug. - Oct.

Notes : It is a source of potent beer.

EREMOPOGON (Hack.) Stapf

Eremopogon foveolatus (Del.) Stapf in Prain, Fl. Trop. Afr. 9: 183. 1917; Bor, l. c. 148. 1960. *Andropogon foveolatus* Del., Fl. d' Egypte 16. t. 8. f. 2. 1812 ; Hook. f., F.B.I. 7 :168. 1896.

Perennials with densely tufted culms. Leaves linear, scaberulous beneath, basal sheath villous ; ligule hyaline, ciliate. Racemes solitary at the end of branches ; joints of raceme and pedicels silky hairy. Sessile spikelets glabrous, pitted, purplish. Caryopsis oblong.

Sikkim, s.l.

Fl. & Fr. : July - Dec.

EULALIA Kunth

Eulalia fastigiata (Nees ex Steud.) Haines, Bot. Bihar & Orissa 1014. 1924 et 3 : 1061 (rep. ed.) Bor, l.c. 155. 1960. *Saccharum fastigiatum* Nees ex Steud., Syn. Pl. Glum. 1: 409. 1855. *Erianthus fastigiatus* Nees ex Steud., l.c. ; Hook. f., F.B.I. 7 : 125. 1896.

Robust grass with erect glabrous culms. Leaves linear, acuminate; sheath fimbriate, throated. Racemes veriticellate at nodes of panicle, densely silky hairy, mauve. Spikelets linear-lanceolate ; upper lemma bidentate.

Sikkim, s. l. ; 500 - 1400 m.

Fl. & Fr. : Aug. - Jan.

E. hirtifolia (Hack.) A. Camus in Ann. Soc. Linn. Lyon n. s. 68 : 202. 1922 ; Bor, l.c. 155. 1960. *Pollinia hirtifolia* Hack. in DC., Mongr. Phan. 6 : 165. 1889 ; Hook. f., F.B.I. 7: 111. 1896.

Annuals with erect leafy culms. Leaves linear, pubescent. Rachis with golden violet stiff hairs. Sessile spikelets lanceolate ; lower glume shiny, ciliate, chartaceous.

Reshi ; ca 2400 m.

Fl. & Fr. : Oct. - Dec.

E. leschenaultiana (Decne.) Ohwi in Bull. Tokyo Muse. No. 18 : 2. 1974 ; Bor, Grass. Ind. 155. 1960. *Andropogon leschenaultianus* Decne., Herb. Timor Descr. 29. 1835. *Pollinia cumingii* Nees in Hook. Kew J. Bot. 2: 98. 1850 ; Hook. f., F. B.I. 7 : 114. 1896. *E. cumingii* (Nees) A. Camus in Lecomte, Fl. Gen. de l' Indo-Chine 7 : 250. 1922.

Perennials. Culms slender, ascending, pubescent. Leaves linear, acuminate, margins scabrid. Racemes with densely golden hairs on sides and back of joints and pedicels. Lower floret absent.

Sikkim ; s.l.

Fl. & Fr. : Oct. - Jan.

E. mollis (Griseb.) O. Ktze., Rev. Bot. Gen. Pl. 2 : 775. 1891 ; Bor, Grass. Ind. 156. 1960. *Erianthus mollis* Griseb. in Nachr. Ges. Wiss. Gottingen 92. 1868. *Pollinia mollis* (Griseb.) Hack. in DC., Mongr. Phan. 6 : 161. 1889 ; Hook. f., F.B.I. 7 : 111. 1896.

Perennials. Culms erect, slender, glabrous. Leaves narrowly lanceolate, acute, glabrous ; ligule small, bearded. Spikes crowded. Spikelets concealed by long hairs.

Sikkim, s.l. ; up to 2400 m.

Fl. & Fr. : Sept. - Oct.

E. quadrinervis (Hack.) O. Ktze., Rev. Bot. Gen. Pl. 2 : 775. 1891 ; Bor, Grass. Ind. 157. 1960. *Pollinia quadrinervis* Hack. in DC., Mongr. Phan. 6 : 158. 1889 ; Hook. f., F.B.I. 7 : 110. 1896.

Perennials. Culms slender, leafy. Leaves linear, acuminate, hairy; margins scabrid. Racemes spreading, white-hairy. Spikelets pale brown, with lower glume distinctly 4-nerved.

Richingpong to Gyalzing ; up to 1500 m.

Fl. & Fr. : Sept. - Dec.

EULALIOPSIS Honda

Eulaliopsis binata (Retz.) C.E. Hubb. in Hook., Ic. Pl. Sub. t. 3262. 1935 ; Bor, l. c. 158. 1960. *Andropogon binatus* Retz., Obs. Bot. 5 : 21. 1789. *Ischaemum angustifolium* (Trin.) Hack. in DC., Mongr. Phan. 6 : 241. 1889; Hook. f., F.B.I. 7 : 129. 1896.

Perennials. Culms erect, densely tufted with woolly base. Leaves narrow, hirsute below, hairs tubercled at base. Upper lemma bidentate, awned. Caryopsis black.

Sikkim, s.l. ; up to 2100 m.

Fl. & Fr. : June - Sept.

HEMARTHRIA R. Br.

Hemarthria compressa (L. f.) R. Br., Prodr. 207. 1810; Bor, l.c. 161. 1960. *Rottboellia compressa* L. f., Suppl. Pl. 114. 1781 ; Hook. f., F.B.I. 7 : 153. 1896.

Perennials. Culms erect, branched, compressed, leafy. Leaves linear, acuminate at apex, rounded at base. Racemes fascicled with almost articulate rachis. Spikelets all similar, lower glume bluntly acuminate.

Ronglikhola, Burtuk Basti, Gangtok ; up to 1800 m.

Fl. & Fr. : Sept. - Dec.

ICHNANTHUS P. Beauv.

Ichnanthus cicinus (F. M. Bail.) Merr. in Enum. Philipp. Fl. Pl. 1 : 70. 1924 ; Bor, l.c. 314. 1960. *Panicum vicinum* F. M. Bail., Syn. Queens. Fl. Suppl 3: 82. 1890. *Ichnanthus pallens* sensu Hook. f., F.B.I. 7 : 60. 1896, non Munro 1861.

Perennials. Culms weak. Leaves oblong-lanceolate, attenuate, amplexicaul or obliquely cordate at base. Panicles emerging from the sheath of flag leaf, wth erect angular rhachis and hispid or woolly branches. Fertile florets with lateral basal appendages or scars.

Sikkim, s.l.

Fl. & Fr. : Sept. - Dec.

IMPERATA Cyr.

Imperata cylindrica (L.) Reauschel., Nom. Bot. ed. 3 : 10. 1797 ; Bor, l.c. 169. 1960. *Lagureos cylindricus* L., Syst. Nat. ed. 10 : 878. 1759. *Imperata arundinacea* Cyr., Pl. Rar. Neap. 2 : 26. t. 11. 1792 ; Hook. f., F.B.I. 7 : 106. 1896. *I. cylindrica* var. *major* (Nees) C. E. Hubb. ex E. E. Hubb. et Vaughan, Grass Manr. 96. 1940 ; Bor, l.c. 170.

Chungthang, Gyalshing, Dentam, Baluakhani.

Fl. & Fr. : May - Nov.

ISCHAEMUM L.

Ischaemum barbatum Retz., Obs. Bot. App. 6 : 35. 1791 ; Bor in Kew Bull. 15 : 411. 1961 *et* Grass. Ind. (rep. ed.) App. I. 1973. *Ischaemum aristatum sensu* Hack. ssp. *imberbe* Hack. var. *imbricatum* Hack. in DC., Mongr. Phan. 6: 203. 1889 ; Hook. f., F.B.I. 7: 127. 1896. *I. goebelli* Hack. in Ost. Bot. Z. 51 : 149. 1901 ; Bor, Grass. Ind. 179. 1960.

Tall grass with smooth terete culms. Leaves lanceolate with cartilaginous margins and very long ligule. Inflorescence of 2 false noded racemes, without a sheath. Sessile spikelets with 1-nerved upper glume ; lower glume with 2 - 3 nodules in lower region only.

Sikkim, s. l. ; up to 1500 m.

Fl. & Fr. : Aug. - Dec.

I. indicum (Houtt.) Merr. in J. Arnold Arbor. 19 : 320. 1938 ; Bor, l.c. 180. 1960. *Paleum indicum* Houtt., Nat. Hist. 2. 13 : 198. t. 90. f. 2. 1782. *I. ciliare* Retz., Obs. Bot. 6 : 36. 1791 ; Hook. f., F.B.I. 7 : 133. 1896. *I. aristatum* auct. non L. : Hook. f., l.c. 126.

Grass with prostrate decumbent culms, rooting at nodes. Leaves linear - lanceolate, hirsute ; ligule small, membranous. Racemes with glabrous peduncle, joints hairy along margins. Sessile spikelets greenish, upper glume many nerved, shortly aristate; lower glume with nodules.

Sikkim, s.l.

Fl. & Fr. : Aug. - Dec.

I. rugosum Salisb., Ic. Stri. Rar. 1. t. 1. 1791 ; Hook. f., F.B.I. 7 : 127. 1896 ; Bor, l.c. 182. 1960.

Tall grass with erect or ascending culms. Leaves linear-lanceolate, almost glabrous. Racemes slightly appressed to each other. Sessile spikelets yellow with distinct nodules on lower glume.

Sikkim, s.l.

MICROSTEGIUM Nees

Microstegium ciliatum (Trin.) A. Camus in Ann. Soc. Linn. Lyon. s.n. 68 : 201. 1921 ; Bor, Grass. Ind. 193. 1960. *Pollinia ciliata* Trin. in Mem. Acad. Sci. Petersb. 6 : 306. 1832 ; Hook. f., F.B.I. 7 : 117. 1896. *Pollinia monantha* Nees ex Steud., Hook. f., l.c.

Perennials. Culms highly branched in lower region. Racemes subdigitate, joints linear, compressed, ciliate in two rows. Sessile spikelets with V-shaped median groove on lower glume, bearded with stiffly ciliate white or purplish hairs. Stamens 3.

Mangan ; up to 1800 m.

Fl. & Fr. : Oct. - Dec.

M. nudum (Trin.) A. Camus in Ann. Soc. Linn. Lyon. n.s. 1832. 68 : 201. 1921 ; Bor, l.c. 194. 1960. *Pollinia nuda* Trin. in Mem. Acad. Sci. Petersb. 6, 2 : 307. 1832 ; Hook. f., F.B.I. 7 : 117. 1896.

Annuals, culms decumbent, multinoded, glabrous. Leaves lanceolate, acuminate. Spikelets with shortly bearded callus ; lower glume with a broad shallow median groove ; upper lemma with a long awn. Caryopsis subovoid, abruptly narrowed at tip. Stamens 2.

Sikkim, s.l. ; up to 1800 m.

Fl. & Fr. : Sept. - Oct.

M. petiolare (Trin.) Bor in Ind. For. Rec. (Bot.) 1 : 87. 1938 et Grass. Ind. 1194. 1960. *Spodiopogon petiolaris* Trin. in Mem. Acad. Sci. Petersb. 6, 2 : 301. 1832. *Ischaemum petiolaris* (Trin.) Hack. in DC., Mongr. Phan. 6 : 238. 1889 ; Hook. f., F.B.I. 7 : 138. 1896.

Grass with tall, glabrous stout culms. Leaves lanceolate, acuminate, hairy ; margins scabrid. Racemes with joints and pedicels subtrigonous. Sessile spikelets with deeply bifid upper lemma. Stamens 2.

Dikchu.

Fl. & Fr. : Sept. - Nov.

Notes : A new record for Sikkim.

M. vagans (Nees ex Steud.) A Camus in Ann. Soc. Linn. Lyon n.s. 68 : 200. 1921 ; Bor, Grass. Ind. 195. 1960. *Pollinia vagans* Nees ex Steud., Pl. Glum. 1 : 410. 1855 ; Hook. f., F.B.I. 7 : 115. 1896. *Pollinia grata* Hack. in DC., Mongr. Phan. 6 : 238. 1889 ; Hook. f., l.c.

Tall grass with branched glabrous culms, pubescent at nodes. Leaves linear-lanceolate, acuminate. Racemes with hairy joints ; joints dilated at top. Sessile spikelets with obtuse lower and upper glume. Stamens 3.

Sikkim ; s.l.

Fl. & Fr. : Oct. - Dec.

Notes : It is used as a fodder grass.

M. vimineum (Trin.) A. Camus in Ann. Soc. Linn. Lyon n.s. 68 : 201. 1921 ; Bor, l.c. 195. 1960. *Andropogon vimineous* Trin. in Mem. Acad. Sci. Petersb. 6, 2 : 268. 1832. *Pollinia imberbis* Nees ex Steud. Syn. Pl. Glum. 1 : 410. 1855 ; Hook.f., F.B.I. 7 : 117. 1896. *M. vimineum* var. *willdenowianum* (Nees) Sur in J. Econ. Tax. Bot. 6 : 175. 1985.

Annuals with glabrous, furrowed culms. Leaves linear-lanceolate, leaf-blades hairy on upper surface. Inflorescence of false spikes ; joints of rachis channelled. Stamens 3, anthers less than 1 mm long. Caryopsis rounded.

1500 - 2000 m.

Fl. & Fr. : Oct. - Dec.

MISCANTHUS Anders.

Miscanthus nepalensis (Trin.) Hack. in DC., Mongr. Phan. 6 : 104. 1889 ; Hook., f. F.B.I. 7 : 107. 1896 ; Bor, Grass. Ind. 196. 1960. *Eulalia nepalensis* Trin. in Mem. Acad. Sci. Petersb. 6, 2 : 333. 1832.

Tufted perennials with small rhizomes. Leaves linear, glabrous. Racemes bearded, with golden brown hairs, 3 - 4 times longer than spikelets. Spikelets all pedicelled with awned lemmas. Caryopsis oblong-ellipsoid.

Baluwakhani, Gangtok, Rate - Chu, Phadomchan, Soreng ; up to 1500 - 2400 m.

Fl. & Fr. : April - June.

M. nudipes (Griseb.) Hack. in DC., Mongr. Phan. 6 : 109. 1889 ; Hook. f., F.B.I. 7 : 107. 1896 ; Bor, Grass. Ind. 196. 1960. *Erianthus nudipes* Griseb. in Nachr. Koen. Akad. Wiss. Goet. 92. 1868.

Perennials with subrobust culms. Leaves linear, pubescent on both surfaces. Inflorescence of fasciculate racemes. Spikelets brownish hairy.

Lachung to Donbyon, Lachen to Tallam, Samdong ; 2100 - 2700 m.

MNESITHEA Kunth

Mnesithea khasiana (Hack.) Koning & Sosef in Blumea 31 : 291. 1986. *Rottboellia striata* Nees ex Steud. ssp. *khasiana* Hack. in DC., Mongr. Phan. 6 : 302. 1889 ; Hook. f., F.B.I. 7 : 157. 1896. *Coelorachis khasiana* (Hack.) Stapf ex Bor in Ind. For. Rec. (Bot.) 1 (3) : 101. 1938 et Grass. Ind. 121. 1960.

Culms solid, erect, branched. Leaves - linear-lanceolate. Racemes of 2 - 4 - jointed fascicles arising from spathe like sheaths. Sessile spikelets in pairs, at each node of axis.

Sikkim, s.l.

M. striata (Nees ex Steud.) Koning & Sosef. in Blumea 31 : 292. 1986. *Rottboellia striata* Nees ex Steud., Syn. Pl. Glum 1 : 361. 1854 ; Hook. f., F.B.I. 7 : 157. 1896. *Coelorachis striata* (Nees ex Steud.) A. Camus in Ann. Soc. Linn. Lyon. n.s. 68 : 1197. 1921. Bor, Grass. Ind. 121. 1960.

Culms glabrous. Leaves linear-lanceolate. Racemes exerted from sheath, nodes without a spathe ; rachis striate. Sessile spikelets not spherical.

Sikkim, s.l.

M. granularis (L.) Koning & Sosef in Blumea 31 : 295. 1986. *Cenchrus granularis* L., Mant. Fl. 2. App. 575. 1771. *Manisuris granularis* (L.) L.f., Nov. Gram. Gen. 40. 1779 ; Hook. f., F.B.I. 7 : 159. 1896. *Hackelochloa granularis* (L.) Ktze., Rev. Gen. Pl. 2 : 776. 1891. Bor, Grass. Ind. 159. 1960. *Manisuris porifera* Hack. in Oest. Bot. Z. 41 : 48. 1891 ; Hook. f., l.c. 160. *Hackelochloa porifera* (Hack.) Rhind, Grass Burma 77. 1954 ; Bor, l.c.

Culms compressed, softly hairy. Leaflets linear-lanceolate, hairy on both surfaces with ciliate margins. Racemes solitary or from axillary branches. Sessile spikelets almost spherical in shape.

Rangeet valley ; 1000 - 1500 m.

OPHIUROS Gaertn. f.

Ophiuros exaltatus (L.) O.Ktze., Rev. Gen. Pl. 2: 780. 1891 ; Bor, Grass. Ind. 199. 1960. *Aegilops exalata* L., Mant. 2 : 575. 1771. *Ophiuros corymbosus* Gaertn. f., Fruct. 3 : 4. t. 181. 1805. Hook. f., F.B.I. 7 : 160. 1896.

Perennials with erect, solid culms, leafy throughout. Leaves linear-lanceolate, with spinulose margins. Spikes corymbosely fasciculate in dense purple tinged fascicles with short peducles. Spikelet solitary, sessile.

Sikkim, s.l.

OPLISMENUS P. Beauv.

Oplismenus burmanii (Retz.) P. Beauv., Ess. Agrost. 54. 168. 1812; Hook. f., F.B.I. 7 : 68. 1896. Bor, l.c. 317. 1960. *Panicum burmanii* Retz., Obs. Bot. 3 : 10. 1783.

Annuals with prostrate or trailing culms. Leaves lanceolate-ovate, sparingly white hairy. Racemes 3 - 8, with hirsute rachis ; spikelets with crossband of hairs and scaberulous awn.

Kabi, Mangan, Singtam, Sangtok ; up to 1800 m.

Fl. & Fr. : July - Sept.

O. compositus (L.) P. Beauv., Ess. Agrost. 54. 168. 1812; Hook. f., F.B.I. 7 : 66. 1896 ; Bor, l.c. 317. 1960. *Panicum compositum* L., Sp. Pl. 57. 1753.

Perennial grass with ascending decumbent culms. Leaves lanceolate to narrowly ovate, minutely hairy below. Spikelets distant, up to 12 mm in the lowest racemes. Awns viscid, smooth.

Gangtok, Mangan ; up to 2400 m.

Fl. & Fr. : June - Dec.

O. undulatifolius (Ard.) R. & S., Syst. Veg. 2 : 482. 1817, non P. Beauv. *nom. nud.* ; Hook. f., F.B.I. 7 : 66. 1896 ; Bor, l.c. 318. 1960. *Panicum undulatifolium* Ard., Animd. Spec. Ait. 14 : t. 4. 1764.

Perennials with trailing culms, rooting at nodes. Leaves undulating. Spikelets contiguous, less than 4 mm apart in lowest racemes.

Sikkim s.l. ; up to 2700 m.

PANICUM L.

Panicum auritum Presl. ex Nees, Agrost. Bras. 176. 1829 ; Hook. f., F.B.I. 7 : 40. 1896 ; Bor, Grass. Ind. 324. 1960.

Perennials with erect leafy culms ; internodes long. Leaves narrowly linear-lanceolate, hairy. Panicles contracted with subsecund fascicles of racemes ; rachis channelled. Spikelets oblong-lanceolate with strongly 3 - 5-nerved lower glumes.

Sikkim, s.l.

P. breviflorium L., Sp. Pl. 59. 1753 ; Bor, Grass. Ind. 324. 1960.
Panicum ovalifolium Poir. in Lam., Encycl. 4 : 279. 1816. Hook. f., F.B.I. 7 : 44. 1896.

Annual grass with ascending, striate culms, rooting at basal nodes. Leaves ovate-lanceolate, acuminate at apex, cordate amplexicaul at base. Branches of panicles with numerous alternate or verticillate branchlets. Spikelets purplish pubescent, with faintly nerved lower glume.

Lower Sichey Busty, Gangtok ; up to 1800 m.

Fl. & Fr. : May - Oct.

P. humidorum Buch.-Ham. ex Hook. f., F.B.I. 7 : 53. 1896 ; Bor, l.c. 326. 1960.

Perennial grass with creeping, procumbent culms, rooting at nodes. Leaves linear with obtuse base. Spikelets smaller, purple, with epaleate lower florets.

Sikkim, s.l.

Fl. & Fr. : July - Dec.

P. khasianum Munro ex Hook. f., F.B.I. 7 : 54. 1896 ; Bor, l.c. 327. 1960.

Perennials. Culms prostrate and creeping, with glabrous nodes. Leaves lanceolate. Ligule narrow, ciliate. Spikelets ellipsoid ; lower glume deltoid and obscurely nerved.

Sikkim, s.l.

P. miliaceum L., Sp. Pl. 58. 1753 ; Hook. f., F.B.I. 7 : 45. 1896 ; Bor, l.c. 327. 1960.

Annuals. Culms stout, clothed with long spreading hairs ; nodes bearded. Leaves linear. Spikelets largest of all species occurring in state. Caryopsis orbicular, plano-convex, white.

Sikkim, s.l.

P. notatum Retz., Obs. Bot. 4 : 18. 1756 ; Bor, Grass. Ind. 701. 1960. *Panicum montanum* Roxb., Fl. Ind. 1 : 315. 1820 ; Hook. f., F.B.I. 7 : 53. 1896.

Perennials with woody base. Culms erect, branched. Leaves lanceolate, cordate at base. Ligule obscure. Spikelets ellipsoid-oblong.

Sikkim, s.l.

P. paludosum Roxb., Fl. Ind. 1 : 310. 1820 ; Bor, Grass. Ind. 329. 1960. *P. proliferum* sensu Hook. f., F.B.I. 7 : 50. 1896, non Lam.

Perennials. Culms erect or ascending, spongy, rooting at nodes. Leaves linear - ensiform. Spikelets ovate, acuminate, green ; lower glume orbicular or reniform, obscurely nerved.

Sikkim, s.l.

Fl. & Fr. : Aug. - Feb.

P. psilopodium Trin., Gram. Panic. 217. 1826 ; Hook. f., F.B.I. 7 : 46. 1896 ; Bor, Grass. Ind. 329. 1960 ; Hsu in Hara, Fl. East Him. 370. 1966 ; Ohwi in Ohashi, Fl. East. Him. 3 : 121. 1975. *P. psilopodium* var. *coloratum* Hook. f., l.c. 47.

Annuals. Culms tufted, with black nodes. Leaves linear - lanceolate ; ligule a ciliate ring. Panicles lax ; spikelets deciduous. Upper glume 11-nerved. Caryopsis elliptic.

Sikkim, s.l.

Fl. & Fr. : July - Nov.

P. repens L., Sp. Pl. ed. 2, 87. 1762 ; Hook. f., F.B.I. 7 : 49. 1896 ; Bor, l.c. 330. 1960.

Perennials with stoloniferous rhizomes. Culms solid. Leaves distichous, linear-lanceolate with involute margins. Panicles lax, with angular rachis. Caryopsis oblong.

Sikkim, s.l.

Fl. & Fr. : Aug. - Sept.

P. sumatrense Roth ex R. & S., Syst. Veg. 2 : 434. 1817 ; Bor, Grass. Ind. 701. 1960. *Panicum millare* Lam., Ill. 1 : 173. 1791 ; Hook. f., F.B.I. 7 : 46. 1896.

Annuals. Culms erect or ascending. Leaves linear ; ligule ciliate, truncate. Panicles with triquetrous rachis, contracted. Spikelets persistent, green or purple tinged. Caryopsis subglobose, dull white.

Sikkim, s.l. ; 600 m.

Fl. & Fr. : July - Oct.

P. walens Mes. in Bot. Jahrb. 34 : 146. 1904, (*watens*). *Panicum humile* Nees ex Steud. Syn. Pl. Glum. 1 : 84. 1854 non Thunb. ex Trin.

1826 ; Hook. f., F.B.I. 7 : 48. 1896. *P. austroasiaticum* Ohwi in Act. Phytotax. Geobot. 2 : 45. 1924 ; Bor, l.c. 324. 1960.

Dwarf annuals with erect, slender, glabrous culms. Leaves linea ; ligule a ciliate rim. Panicles open, ovate-oblong. Spikelets ellipsoid, shortly acuminate with paleate lower floret.

Sikkim, s.l.

PASPALIDIUM Stapf

Paspalidium flavidum (Retz.) A. Camus in Lecomte, Fl. Gen. de l' Indo - Chine 7 : 419. 1922 ; Bor, l.c. 333. 1960. *Panicum flavidum* Retz., Obs. Bot. 4 : 15. 1786 ; Hook. f., F.B.I. 7 : 28. 1896.

Tufted perennials. Culms slender, glabrous, somewhat compressed. Leaves linear-lanceolate. Inflorescence of pseudo spikes. Spikelets abaxial with developed lower glume and paleate male lower floret ; upper lemma granulose.

Sikkim, s.l.

Fl. & Fr. : Aug. - Sept.

P. punctatum (Burm. f.) A. Camus in Lecomte, Fl. Gen. de l' Indo - Chine 7 : 419. 1922 ; Bor, l.c. 333. 1960. *Panicum puctatum* Burm. f., Fl. Ind. 26. 1768 ; Hook.f., F.B.I. 7 : 29. 1896.

Perennials with spongy rhizomes. Culms decumbent or prostrate, rooting at nodes. Leaves linear. Inflorescence consisting of distantly placed pseudo spikes. Spikelets with neuter, lower floret without palea ; upper lemma rugulose.

Sikkim, s.l.

Fl. & Fr. : Sept. - Oct.

PASPALUM L.

Paspalum conjugatum Bewrg. in Act. Helvet. Phys. Manth. 7 : 129. t.8. 1773 ; Hook. f., F.B.I. 7 : 11. 1896.

Creeping, stoloniferous herbs. Leaves lanceolate, sparsely hairy above. Spikelets 2-seriate, plano-convex, with a ciliate fringe from margins.

Sikkim, s.l.

P. dilatatum Poir. in Lam., Encyclop. 5 : 35. 1804 ; Bor, Grass. Ind. 338. 1960.

Tufted perennials. Culms robust, leafy at base. Leaves linear-lanceolate, acuminate. Inflorescence of 3 - 5 (-10) racemes with 2 - 4 rows of paired spikelets. Spikelets ovate, pubescent, yellowish green ; upper glume concave, ciliate at margins, sparsely pilose above.

Gangtok - Penangla (on way to Tashi View Point, TNA Campus).

P. paspalodes (Mich.) Scribn. in Mem. Tor. Bot. Club. 5 : 129. 1894. (*paspaloides*). *Digitaria paspalodes* Michx., Fl. Bor. Amer. 1 : 46. 1803. *Paspalum distichum* auct. non L. 1759. *nom. confus* : Hook. f., F.B.I. 338. 1960.

Creeping stoloniferous perennials. Culms dark, hairy, noded. Leaves with rounded ciliate base. Spikelets solitary in two rows, plano-convex pale green ; lower glumes minute scaly. Upper glumes appressely puberulent, herbaceous.

Lindok, East Distt.

Fl. & Fr. : April - May ; Aug. - Sept.

P. scobiculatum L., Mant. 1: 29. 1767 ; Hook. f., F.B.I. 7 : 10. 1896, *p.p.*; Bor, *l.c.* 340. 1960. *P. orbiculare* Forst. *f.*, Fl. Insul. Austr. Prodr. 7. 1786 ; Bor, *l.c.* *P. commersonii* Lam., Ill. 1 : 175. t. 43. f. 1. 1791 ; Bor, *l.c.* 335. *Paspalum cartilagineum* J.S. ex C.B. Presl., Rel. Haenk. 1 : 216. 1830 ; Bor, *l.c.* 335.

Perennials with tufted spongy culms, rooting at nodes. Leaves lanceolate. Racemes digitate, borne on a long axis. Spikelets in two rows ; lower glume lacking ; upper glume papyraceous, lower floret sterile.

Lingdok, East Distt.

Fl. & Fr. : April - Dec.

PENNSETUM Rich. ex Pers.

Pennisetum americanum (L.) K. Schum. in Engl, Pflanz. Oest. Afr. B. 51. c. t. 4. f. A. & B. 1895. *Panicum americanum*. L., Sp. Pl. 56. 1753. *Pennisetum typhodeum* L.C. Rich. in Pers., Syn. Pl. 1: 72. 1805 ; Hook. *f.*, F.B.I. 7 : 86. 1896. *P. typhodes* (Burm. *f.*) Stapf & C.E. Hubb. in Kew Bull. 1933 ; 271. 1933 ; Bor, Grass. Ind. 350. 1960.

Tall annuals. Culms erect with bearded nodes. Leaves linear-lanceolate. Panicles compact, cylindrical, pubescent. Spikelets ovate-elliptic, subtended by cream coloured or brownish involucral bristles. Caryopsis ellipsoid-orbicular.

Cultivated in tropical zone.

Fl. & Fr. : Sept. - Oct.

P. clandestinum Hochst. ex Chiov. in Ann. Ist Bot. Roma 8 : 41. t. 5. f. 2. 1903, Bor, Grass. Ind. 344. 1960.

Perennials with many creeping rhizomes. Culms short, copiously branched, appressed. Leaves linear, obtuse. Inflorescence consisting of cluster of 2 - 4 spikelets. Caryopsis oblong-ellipsoid, brown.

Introduced, naturalized.

Notes : A fodder grass locally known as "Kikuya Grass".

P. flaccidum Griseb. in Nachr. Ges. Wiss. Goett. 86. 1868 ; Hook.f., F.B.I. 7 : 84. 1896 ; Bor, Grass. Ind. 344. 1960.

Tufted perennials with thick rhizomes. Culms erect, branched. Leaves flat or folded. Panicles linear, with scabrid rachis having shallow ribs below cupular scars.

Sikkim, s.l.

P. glaucum (L.) R. Br., Prodr. 1 : 195. 1810. *Panicum glaucum* L., Sp. Pl. 56. 1753. *Setaria glauca* (L.) P. Beauv., Ess. Agrost. 51 : 169. 178. 1812 ; Hook. f., F.B.I. 7 : 78. 1896 ; Bor, l.c. 344. 1960.

Annuals. Culms loosely tufted, geniculate. Leaves linear-lanceolate. Inflorescence consisting of pseudospikes, erect, dense, with minutely pubescent rachis. Involucre persistent.

Sikkim, s. l.

Fl. & Fr. : Sept. - Oct.

P. orientale L.C. Rich. in Pers, Syn. Pl. 1 : 72. 1805 ; Hook.f., F.B.I. 7 : 86. 1896 ; Bor, Grass. Ind., 34. 1960.

Perennials with creeping stout rhizome. Culms tall, erect, with glabrous nodes. Leaves linear. Spikes erect or nodding, pinkish, with flexuous villous or woolly rachis. Outer spikelets imperfect, anther-cells not bearded at apex. Caryopsis ellipsoid.

Sikkim, s.l.

Fl. & Fr. : Aug. - Dec.

P. pedicellatum Trin. in Mem. Acad. Petersb. ser. 6,3 : 184. 1834 ; Hook.f., F.B.I. 7 : 86. 1896 ; Bor, l.c. 346. 1960.

Annuals with tall, glabrous, noded culms. Leaves linear-lanceolate. Spikes reddish purple, dense, erect or nodding, with angular, winged or ribbed rachis. Spikelets with silky hairs on back. Caryopsis oblong-lanceolate.

Jorethang.

Fl. & Fr. : Aug. - Jan.

A. purpureum K. Schum., Beskr. Guin. Pl. 44. 1827 ; Bor, I.c. 348. 1960. ELEPHANT or NAPIER GRASS (Eng.).

Perennials with creeping rhizomes. Culms erect, branched ; nodes with a ring of stiff, long appressed hairs. Leaves linear, acuminate, purple tinged, glabrous beneath. Inflorescence consisting of dense cylindric yellow or purple tinged pseudospikes; lateral pedicelled spikelets surrounded by involucre of bristles of unequal length.

Sikkim, s.l.

PHACELURUS Griseb.

Phacelurus zea (C.B.Clarke) W.D. Clayton in Kew Bull. 33 : 177. 1978. *Rotboella zea* C.B. Clarke in J. Linn. Soc. 25 : 86. f. 35. 1889 ; Hook. f., F.B.I. 7 : 152. 1896. *Thysis zea* (C.B. Clarke) Stapf in Hook., Ic. Pl. sub. t. 3078. 1892 ; Bor, I.c. 255. 1960.

Culms tall, glabrous. Leaves with serrulate-ciliate margins. Particles of subverticillate spikes ; joints of rachis swollen, dorsally compressed ; margins of lower glume narrowly inflexed.

Sikkim, s.l.

Fl. & Fr. : Aug. - Dec.

POGONATHERUM P. Beauv.

Pogonatherum crinitum (Thumb.) Kunth, Enum. Pl. 1 : 478. 1833 ; Hook. f., F.B.I. 7 : 141. 1896 ; Bor, Grass. Ind. 200. 1960. *Andropogon crinitus* Thunb., Fl. Japan 40 : t. 1748.

Perennials. Culms slender, highly branched. Leaves linear-lanceolate. Racemes dense with long delicate yellowish brown awns. Spikelets up to 2 mm ; upper lemma deeply cleft ; lower floret empty.

Rhenock, on way to Jorethan.

Fl. & Fr. : May - June.

P. paniceum (Lam.) Hack. in Allg. Bot. Z. 178. 1906 ; Bor, Grass. Ind. 202. 1906. *Saccharum paniceum* Lam., Encycl. 1 : 595. 1785. *P. saccharoideum* P. Beauv., Ess. Agrost. 176. 1812 ; Hook. f., F.B.I. 7 : 141. 1896.

Tufted perennials. Culms erect, stiff. Leaves linear-lanceolate. Racemes with many delicate yellow-brown awns. Spikelets more than 2 mm long, lower floret male.

Singtam, Ranipool, Gangtok, Burtuk Busti.

POLYTOCA R. Br.

Polytoca digitata (L.f.) Druce in Rep. Bot. Exch. Cl. Brit. Is. 4 : 641. 1916 ; Bor, l.c. 267. 1960. *Apluda digitata* L.f., Suppl. Pl. 434. 1781. *Polytoca bracteata* R.Br., Pl. Jav. Rar. 20. t. 5. 1838 ; Hook. f., F.B.I. 7 : 101. 1896.

Perennials. Culms with one sided grooved sheath at base, tomentose at nodes. Leaves linear. Racemes male, terminal, in lateral ones female or male at top, or female always at base. Lower glume obtuse, mucronate without an arista.

Sikkim, s.l.

Fl. & Fr. : July - Dec.

P. wallichiana (Nees ex Steud.) Benth. in J. Linn. Bot. 19 : 52. 1881 ; Hook.f., F.B.I. 7 : 101. 1896 ; Bor, l.c. 267. 1960. *Cyathorhachis wallichiana* Nees ex Steud., Syn. Pl. Glum. 1 : 403. 1855.

Perennials with well developed stolons. Culms leafy glabrous. Leaves linear. Panicles compound, with terminal node and lateral female spikelets. Lower glume toothed on one side, produced into a caudate arista.

Sikkim, s.l.

PSEUDECHINOLAENA (Hook.f.) Stapf

Pseudechinolaena polystachya (H.B.K.) Stapf in Prain, Fl. Trop. Afr. 9 : 495. 1919 ; Bor, Grass. Ind. 352. 1960. *Echinolaena polystachya* H.B.K., Nov. Gen. Sp. Pl. 1 : 119. 1816. *Panicum uncinatum* Raddi, Agrost. Bras. 41. 1823 ; Hook. f., F.B.I. 7 : 58. 1896.

Perennials. Culms pubescent, noded, branched, rooting at base. Leaves ovate-lanceolate. Panicles with slender rachis, filiform branches. Spikelets laterally compressed, obliquely ovate, upper glume hispid.

Sikkim, s.l.

Notes : It forms thick masses in secondary forest. Its fruits stick to the legs of grazing animals by means of hooked spines of upper glume.

PSEUDOPOGONATHERUM A. Camus

Pseudopogonatherum contortum (Brongn.) A. Camus in Lecomte, Fl. Gen. de l' Indo - Chine 7 : 255. 1922 ; Bor, l.c. 204. 1960. *Pollinia articulata* Trin. in Hook. f., F.B.I. 7 : 109. 1896. *P. articulata* Trin. var. *pedicellata* Hack. ex Hook. f., l.c. 110.

Annuals. Culms slender, densely tufted. Leaves filiform to linear. Panicles silky white. Spikelets all pedicelled; upper glume with a long slender awn.

Sikkim, s.l.

Notes : A gregarious grass of open hill side; morphologically very variable.

P. brunonianus Griff., Not. 3, 29. 1851; Bor, l.c. 353. 1960. *Holcus natans* Roxb. ex Hook. f., F.B.I. 7 : 62. 1896. *Chamaeraphis spinescens* Poir. var. *brunonianus* Hook. f., l.c.

On way to Tinbon, Lachung; up to 2700 m.

Notes : Found in swampy places.

PSEUDOSORGHUM A. Camus

Pseudosorghum fasciculare (Roxb.) A. Camus in Bull. Mus. Hist. Nat. Paris 26 : 662. 1920; Bor, l.c. 205. 1960. *Andropogon fascicularis* Roxb., Fl. Ind. 1 : 269. 1820; Hook. f., F.B.I. 7 : 177. 1896. *A. nitidulus* Hook. f., l.c. 199.

Annuals. Culms glabrous, decumbent, slender. Leaves lanceolate. Racemes spiciform with joints of rachis and pedicels without a median furrow. All pairs of spikelets heterogamous; upper lemma of sessile spikelets bifid.

Sikkim, s.l.

Fl. & Fr. : Sept. - Dec.

ROTTBOELLIA L.f. nom. cons.

Rottboellia cochinchinensis (Lour.) Clayton in Kew Bull. 35 : 1817. 1981. *Stegosia cochinchinensis* Lour., Fl. Cochinch. 1 : 51. 1790. *R. exaltata* L.f., Suppl. Pl. 114. 1781, non (L.) L.f. 1779; Hook. f. in F.B.I. 7 : 156. 1896; Bor, Grass. Ind. 206. 1960. *R. exaltata* L.f. var. *robusta* Hook. f., l.c.

Perennials. Culms erect, leafy, solid, branched at base. Leaves linear-lanceolate. Racemes solitary, terete, with dorsally glabrous internodes. Sessile spikelets solitary at each node of axis. Pedicelled spikelets imperfect with narrowly winged lower glume.

Sikkim, s.l.

Fl. & Fr. : Oct. - Feb.

SACCHARUM L. Pl.

Saccharum longisetosum (Anders. ex Benth.) Narayanasw. ex Bor, Fl. Assam 5 : 461. 1940 et Grass. Ind. 211. 1960. *Erianthus longisetous* Anders. ex Benth. in J. Linn. Soc. Bot. 19 : 67. 1881 ; Hook. f., F.B.I. 7 : 124. 1896.

Perennials. Culms solid, arising from stout rhizomes. Leaves lanceolate with scabrid margins. Panicles densely packed with drooping racemes, clothed with reddish brown hairs. Lower glume chartaceous, glabrous, upper glume 1-nerved.

Gangtok.

var. hookeri (Hack.) Bor, Grass. Ind. 212. 1960. *Erianthus hookeri* Hack. in DC., Mongr. Phan. 6 : 143. 1889 ; Hook.f., F.B.I. 7 : 125. 1896.

Perennials with closely fitted sheath. Panicles of silky racemes; rachis ciliate at angles. Lower glume dorsally villous, upper glume 3-nerved.

Lachen, Rangpo, Rorathang.

S. narenga (Nees ex Steud.) Hack. in DC., Mongr. Phan. 6 : 119. 1889. *Erianthus narenga* Nees ex Steud., Syn. Pl. Glum. 1 : 411. 1855. *Narenga porphyrocoma* (Hance ex Trim.) Bor in Ind. For. 66 : 267. 1940 et Grass. Ind. 198. 1960.

Stout perennials with tall erect culms. Densely bearded at nodes, appressed silky below panicle. Panicles narrow, erect, woolly, purplish or whitish. Glumes coriaceous, obtuse.

Sikkim, s.l.

S. officinatum L., Sp. Pl. 54. 1753. Hook.f., F.B.I. 7 : 118. 1896 ; Bor, Grass. Ind. 212. 1960.

Grass with juicy, glabrous culms, waxy below nodes. Leaves linear-lanceolate. Panicles dense, ovate pyramidal. Spikelets surrounded by dense white silky hairs.

Cultivated in valleys.

S. procerum Roxb., Fl. Ind. 1 : 248. 1820 ; Bor, l.c 213. 1960.

Tall grass with solid, terete culms. Leaves linear - lanceolate. Panicles large, with silvery - grey hairy. Sessile spikelets with setaceous upper glume, curved at apex. Pedicelled spikelets with dorsally villous glumes.

Sikkim, s.l.

S. rufipilum Steud., *Syn. Pl. Glum.* 1 : 409. 1855 ; Bor, *I.c.* 213. 1960. *Erianthus versicolor* Nees ex Steud., *I.c.* ; Hook. f., *F.B.I.* 7 : 124. 1896. *E. rufipilus* (Steud.) Griesb. in *Nachr. Ges. Wiss. 'Gottingen'* 93. 1868 ; Bor, *I.c.* 152. *E. versicolor* var. *palens* (Hack.) Hook. f., *I.c.* *E. fulvus* Nees ex Steud. *Syn. Pl. Glum.* 1 : 409. 1855, *E. fulvus* Nees ex Hack. in DC., *Mongr. Phan.* 6 : 147. 1889 ; Hook. f., *I.c.* 123.

Perennials. Culms erect, robust. Leaves linear, flat or convolute. Panicles with appressed silky peduncles. Spikelets brownish, with reddish silky hairs and membranous lower glumes.

Gangtok, Bop near Chungthang, on way to Rongli.

Fl. & Fr. : July - Sept.

S. sikkimensis (Hook.f.) Narayana. ex Bor, *Fl. Assam* 5 : 462. 1940 et *Grass. Ind.* 214. 1960. *Erianthus sikkimense* Hook. f., *F.B.I.* 7 : 123. 1896.

Perennials. Culms tall, slender. Leaves narrow, flat. Panicles silky villous below ; rachis villous or appressed hairy. Spikelets with coriaceous lower glume and entire upper lemma with a short awn.

Lachen.

S. spontaneum L., *Mant. Pl.* 2 : 183. 1771 ; Hook. f., *F.B.I.* 7 : 118. 1896 ; Bor, *Grass. Ind.* 214. 1960.

Perennials, with thick rhizomes. Culms erect, solid, glabrous, with hairy nodes. Leaves petiolate towards base. Panicles lax, silvery white, pedicels ciliate. Spikelets with long silky hairs and acuminate lower glume.

On way to Rumtek, Rangeet, Rishop, Teesta, below Chhangu.

SACCIOLEPIS Nash.

Sacciolepis indica (L.) A. Chase in *Proc. Biol. Soc. Wash.* 218. 1908 ; Bor, *I.c.* 357. 1960. *Aira indica* L., *Sp. Pl.* 1753 in '*Errata*'. *Panicum indicum* L., *Mant. Pl.* 2 : 184. 1771, *non* Mill. 1786. *P. indicum* L. var. *angustum* (Trin.) Hook. f., *I.c.* 428. *P. indicum* L. var. *gracile* Hook. f., *I.c.*

Tufted annuals. Culms glabrous, striate. Leaves linear-lanceolate. Panicles cylindrical, spiciform, continuous. Spikelets obliquely lanceolate, acute, relatively smaller, greenish purple. Caryopsis oblong.

Rongli - khola, Ranipool, Deorali, Lindok.

S. interrupta (Willd.) Stapf in Prian, Fl. Trop. Afr. 9 : 757. 1920 ; Bor, l.c. 358. 1960. *Panicum interruptum* Willd., Sp. Pl. 1 : 341. 1798 ; Hook. f., F.B.I. 7 : 40. 1896.

Perennials. Culms ascending from a stout creeping rootstock. Rootstock spongy. Leaves linear, flat, rounded or subcordate at base. Panicles cylindrical, interrupted, spiciform. Spikelets densely crowded, acute, green. Caryopsis obovoid, apiculate.

Sikkim, s.l.

Notes : A very good fodder grass relished by cattle.

S. myosuroides (R. Br.) A. Camus in Lecomte, Fl. Gen. de l' Indo - Chine 7 : 460. 1922 ; Bor, l.c. 358. 1960. *Panicum myosuroides* R. Br., Prodr. 189. 1810 ; Hook. f., F.B.I. 7 : 42. 1896.

Annuals. Culms glabrous. Leaves narrow linear, flat, wooly. Panicles narrow, spiciform, cylindric, dense. Spikelets obtuse, pale green, purple tinged at tip.

Chungthang, Soreng, Rumtek.

SCHIZACHYRIUM Nees

Schizachyrium brevifolium (Swartz) Nees ex Buse in Miq, Pl. Jungh. 359. 1854 ; Bor, l.c. 215. *Andropogon brevifolium* Swartz, Prodr. Veg. Ind. Occ. 26. 1788 ; Hook. f., F.B.I. 7 : 165. 1896.

Annuals. Culms weak, slender. Leaves linear, sharply pointed at apex. Racemes slender, solitary, concealed in a narrow spathe, joints glabrous. Caryopsis compressed, glabrous.

Sikkim, s.l.

Fl. & Fr. : Aug. - Jan.

S. delaveyi (Hack.) Bor in Ind. For. Rec.(Bot.) 1 : 95. 1938 et Grass. Ind. 215. 1960. *Andropogon delaveyi* Hack. in DC., Mongr. Phan. 6 : 40. 1889. *A. bootanensis* Hook. f., F.B.I. 7 : 166. 1896.

Perennials. Culms many noded, erect. Leaves linear. Spathe reddish. Racemes compressed.

Sikkim, s.l. ; 2400 - 2600 m.

SETARIA P. Beauv. *nom. cons.*

Setaria barbata (Lam.) Kunth, Rev. Gram. 1 : 47. 1829 ; Bor, Grass. Ind. 360. 1960. *Panicum barbatum* Lam., Ill. 1 : 171. 1791. *P. flavescens* sensu Hook. f., F.B.I. 7 : 56. 1896, non Swartz 1788. *P. rhachitrichum* Hochst. in Flora 27 : 254. 1844 ; Hook. f., l.c. 56.

Annuals. Culms slender. Leaves linear-lanceolate, plicate, broad, softly hairy on both surfaces. Panicles narrow, loose. Spikelets subsessile, green, subtended by persistent bristles. Upper lemma transversely rugose. Caryopsis apiculate.

Jorethang ; 1500 - 1800 m.

Fl. & Fr. : July - Oct.

S. forbesiana (Nees ex Steud.) Hook. *f.*, F.B.I. 7 : 81. 1896 ; Bor, *I.c.* 360. 1960. *Panicum forbesianum* Nees ex Steud., *Syn. Pl. Glum.* 1 : 98. 1854.

Perennials. Culms tufted. Leaves linear-lanceolate, acuminate. Panicles narrowly pyramidal, erect. Spikelets apiculate. Caryopsis *ca* 2.5 mm long, elliptic.

Sikkim, *s.l.* ; 1000 - 1800 m.

Fl. & Fr. : Aug. - Dec.

S. intermedia R. & S., *Syst. Veg.* 2 : 489. 1817 ; Hook. *f.*, F.B.I. 7 : 79. 1896. *Panicum tomentosum* Roxb., *Fl. Ind.* 1 : 303. 1820. *Setaria tomentosa* (Roxb.) Kunth, *Rev. Gram.* 1 : 47. 1829 ; Bor, *Grass. Ind.* 365. 1960.

Annuals. Culms weak, ascending. Leaves broadly linear, acuminate, with bristles along midrib. Panicles contracted. Caryopsis *ca* 1.5 mm long, obtuse.

Sikkim, *s.l.*

Fl. & Fr. : Sept. - Oct.

S. italicica (L.) P. Beauv., *Ess. Agrost.* 51. 170. 178. 1812 ; Hook. *f.*, F.B.I. 7 : 78. 1896 ; Bor, *I.c.* 362. 1960. *Panicum italicum* L., *Sp. Pl.* 56. 1753.

Annuals. Culm erect or ascending. Leaves linear. Panicles spiciform, cylindric. Spikelets deciduous, oblong-elliptic. Caryopsis broadly elliptic, *ca* 2 mm long.

Cultivated in tropical zone.

S. palmifolia (Koen.) Stapf in J. Linn. Soc. Bot. 42 : 186. 1914. *Panicum palmifolium* Koen. in *Naturf.* 23. 208. 1788. *P. plicatum* Willd., *Enum. Pl.* 1023. 1809, *non* Lam. 1791.

Perennials. Rootstocks branched, woody. Culms erect or ascending. Leaves linear-lanceolate, acuminate, plicate between nerves. Spikelets ovoid, lower lemma apiculate.

Sikkim, s.l.

Fl. & Fr. : July - Nov.

S. plicata (Lam.) T. Cooke, Fl. Pres. Bombay 2 : 919. 1908 ; Bor, l.c. 364. 1960. *Panicum plicatum* Lam., Encycl. 4 : 736. 1797 ; Hook. f., F.B.I. 7 : 55. 1896.

Perennials. Culms tufted, leafy. Leaves elliptic-oblong to linear-lanceolate, plicate. Panicles erect or drooping with purplish green branches. Bristles stiff at the end of branchlets. Spikelets ovoid ; lower lemma dorsally flattened. Caryopsis globose.

On way to Pakyong, Baluwakhani, Gangtok.

Fl. & Fr. : Aug. - Jan.

S. pumila (Poir.) R. & S., Syst. Veg. 2 : 891. 1817. *Panicum pumillum* Poir. in Lam., Encycl. 4 : 273. 1816. *Setaria pallida-fusca* (K. Schum.) Stapf & C. E. Hubb. in Kew. Bull. 1930 : 250. 1930 ; Bor, Grass. Ind. 363. 1960. *S. glauca* auct. non (L.) P. Beauv.

Annuals. Culms loosely tufted. Leaves linear. Panicles spiciform, continuous. Bristles antrorsely banded, 7 - 8 in each involucre. Spikelets ovate. Caryopsis elliptic, plano-convex.

Burtuk, Gangtok, Lindok, Reshi, Rumtek, Gyalshing, Rate - Pani, Melli.

Fl. & Fr. : June - Oct.

S. verticillata (L.) P. Beauv., Ess. Agrost. 51. 178. 1812 ; Hook. f., F.B.I. 7 : 80. 1896 ; Bor, l.c. 365. 1960. *Panicum verticillatum* L., Sp. Pl. ed. 2 : 82. 1762.

Annuals. Culms tall erect or ascending. Leaves linear-lanceolate, sparsely pilose or hispid. Panicles interrupted spiciform. Bristles retrorsely barbed. Spikelets pale green. Caryopsis ca 2 mm long, elliptic.

Sikkim, s.l. ; 1500 - 1800 m.

Fl. & Fr. : July - Oct.

SORGHUM Moench *nom. cons.*

Sorghum miliiforme (Hack.) Snowden in Kew Bull. 1935 : 237. 1935 ; Bor, l.c. 239. 1960. *Andropogon sorghum* ssp. *sativus* var. *miliiformis* Hack. in DC., Mongr. Phan. 6 : 518. 1889 ; Hook. f., F.B.I. 7 : 184. 1896.

var. **miliiforme**

Annuals. Culms up to 2 m high. Leaves linear, very large. Panicles 5 - 10 cm broad. Spikelets paired, straw coloured ; lower glume reddish. Caryopsis *ca* 3 mm, orbicular, dark brown to violet brown.

Sikkim, *s.l.* ; 1500 m. Cultivated.

var. sikkimensis Snowden in Kew Bull. 1935 ; 237. 1935 ; Bor, *I.c.* 239. 1960.

Annuals. Panicle erect, contracted, 15 - 20 cm broad. Caryopsis orbicular, *ca* 4 mm.

Sikkim, *s.l.* ; *ca* 1500 m. Cultivated.

THEMEDA Forssk.

Themeda arundinacea (Roxb.) Ridley in Trans. Linn. Soc. ser. 2, 3 : 401. 1893 ; Bor, *I. c.* 250. 1960. *Anthistria arundinacea* Roxb., Fl. Ind. 1 : 256. 1820. *A. gigantea* (Roxb.) Hack. in DC., Mongr. Phan. 6 : 774. 1889 ; Hook. *f.*, F.B.I. 7 : 216. 1896.

Tufted perennials with solid, erect, smooth, polished culms up to 5 m high. Leaves linear with subulate tip. Panicles nodding at apex ; involucral spikelets with golden hairs, sessile spikelets bearded. Awns up to 5 cm long.

Rhenock ; *ca* 1200 m.

Fl. & Fr. : Aug. - Jan.

T. hookeri (Griseb.) A. Camus in Bull. Mus. Hist. Nat. Paris 26 : 425. 1920 ; Bor, *I.c.* 251. 1960. *Anthistria hookeri* Griseb. in Goett. Nachr. 91. 1868 ; Hook. *f.*, F.B.I. 7 : 216. 1896.

Perennials. Culm up to 60 cm tall, slender. Leaves narrow, linear, glabrous. Awns up to 2.5 cm long.

Chungthang, Lachung ; up to 2500 m.

T. quadrivalvis (L.) O. Ktze., Gen. Pl. 2 : 994. 1891 ; Bor, *I.c.* 252. 1960. *Andropogon quadrivalvis* L. in Mur. Syst. Veg. ed. 13 : 758. 1774. *Anthistiria ciliata* L. *f.*, Suppl. 113. 1781 ; Hook. *f.*, F.B.I. 7 : 213. 1896.

Annuals. Culms tall, erect or ascending, up to 2 cm tall ; nodes glabrous. Involucral spikelets in pairs, inserted at different leaves, less than 6 mm long. Awns up to 3.5 mm long.

Mangan ; up to 2000 m.

Fl. & Fr. : Aug. - Jan.

T. triandra Forssk., Fl. Aeg. - Arab. 1238. 178. 1775 ; Bor, l.c. 254. 1960. *Anthistiria imberbis* Retz., Obs. Bot. 3 : 11. 1783 ; Hook. f., F.B.I. 7 : 211. 1896. *Themeda imberbis* (Retz.) T. Cooke, Fl. Pres. Bombay 2 : 993. 1908.

Perennials. Culms erect, up to 1 m tall. Panicles enclosed in spatheoles. Awns up to 6 mm long. Caryopsis up to 3.5 mm long, truncate at apex.

Chalisay on way to Rongli.

Fl. & Fr. : Oct. - Nov.

VETIVERIA Bory. de St. Vincent.

Vetiveria zizanoides (L.) Nash in Small, Fl. South-east U.S. 67. 1903; Bor, l.c. 258. 1960. *Phalaris zizanioides* L., Mant. Pl. 2 : 183. 1771. *Andropogon squarrosus* L.f., Suppl. 433. 1781. Hook. f., F.B.I. 7 : 186. 1896.

Perennials. Roots stout, spongy, aromatic. Culms up to 2 m tall, tufted. Panicles contracted, with many pairs of spikelets arranged in whorled racemes.

Sikkim, s.l. ; 1200 m.

Fl. & Fr. : Sept.

ZEA L.

Zea mays L., Sp. Pl. 971. 1753 ; Hook. f., F.B.I. 7 : 102. 1896.

Annual robust grass with tall erect culms. Leaves broadly lanceolate, hairy on upper surface. Male inflorescence a panicle. Female inflorescence axillary, sessile, enclosed in a leafy sheath surrounded by bracts. Spikelets conical, arranged in longitudinal rows. Caryopsis subglobose, shiny, hard, pale yellow or white-orange to dark purple.

Cultivated in tropical and subtropical regions.

Fl. & Fr. : July - Sept.

Notes : It is one of the main crops of Sikkim.

Sub fam. POOIDEAE

(Paramjit Singh)

Agrostis filipes Hook. f., F.B.I. 7 : 256. 1896 ; Bor, Grass. Ind. 387. 1960.

Small tufted perennials. Panicles narrow, interrupted ; glumes ca 2.5 mm long, equal.

Changu, Chamnago.

Fl. & Fr. : July - Sept.

A. gigantea Roth, Tent. Fl. Germ. 1 : 31. 1788 ; Bor, Grass. Ind. 387. 1960. *A. stolonifera* L. subsp. *gigantea* (Roth) Naie and Weiller, Fl. Afr. Nord. 2 : 120. 1893. *A. alba* auct. non Hook. f., F.B.I. 7 : 254. 1896.

Rhizomatous perennials. Panicles very effuse, glumes equal.

Sikkim, s.l. ; 2900 - 4000 m (*fide* Hook.f., *op.cit.*).

A. himalayana Bor in Kew Bull. 1953 ; 269. 1953 ; Bor, Grass. Ind. 387. 1960.

Culms up to 30 cm high. Spikelets 1 - 1.2 mm long ; glumes equal ; palea minute.

Mainam, Dentam R.F.

Fl. & Fr. : July - Aug.

A. hookeriana Clarke ex Hook. f., F.B.I. 7 : 256. 1896 ; Bor, Grass. Ind. 387. 1960. *A. poluninii* Bor in Kew Bull. 1953 : 269. 1953.

Culms 30 - 60 cm high, slender. Spikelets 3 - 4 mm long ; glumes unequal ; lemma awned.

Changu, Dzongri ; 3000 - 3400 m.

Fl. & Fr. : July - Oct.

A. inaequiglumis Griseb. in Goett. Nachr. 80. 1868 ; Hook. f., F.B.I. 7 : 257. 1896 ; Bor, Grass. Ind. 387. 1960.

Culms 5 - 15 cm high. Panicles spiciform. Spikelets ca 2.5 mm long ; glumes unequal ; palea absent.

Chakung Chu, Karponang, Momay - Samdong ; 3000 - 4500 m.

Fl. & Fr. : July - Oct.

A. micrantha Steud., Syn. Pl. Glum. 1 : 170. 1854 ; Hook. f., F.B.I. 7 : 256. 1896 ; Bor, Grass. Ind. 388. 1960.

Culms up to 1 m high. Spikelets ca 2 mm long ; pedicels clavate ; glumes subequal ; lemma ovate-truncate ; palea rectangular.

Lachen.

Fl. & Fr. : July - Nov.

A. myriantha Hook. f., F.B.I. 7 : 257. 1896 ; Bor, Grass Ind. 388. 1960. *A. myriantha* Hook. f. var. *sikkimensis* Hook. f., F.B.I. 7 : 257. 1896.

Culms up to 70 cm high. Panicles 15 - 30 cm long; glumes equal ; lemma 1.25 mm long ; palea minute.

Dikchu, Phadomchen.

Fl. & Fr. : July - Sept.

A. nervosa Nees ex Trin. in Mem. Acad. Sci. Petersb. Ser. 6, 6 : 328. 1814 ; Bor, Grass Ind. 388. 1960. *A. clarkei* Hook. f., F.B.I. 7 : 257. 1896.

Culms 15 - 30 cm long. Panicles pyramidal, 14 - 15 cm high. Spikelets 2 - 2.5 mm long ; glumes unequal, lemma 1.5 mm long ; palea absent or minute.

Changu, Dzongri ; 3300 - 4000 m.

Fl. & Fr. : Aug. - Sept.

A. pilosula Trin. in Mem. Acad. Sci. Petersb. Ser. 6, 6 : 372. 1814 ; Bor, Grass. Ind. 388. 1960. *Calamagrostis pilosula* (Trin.) Hook. f., F.B.I. 7 : 263. 1896.

Culms 30 - 90 cm high. Panicles 10 - 20 cm long. Spikelets 2.5 - 3.5 mm long ; glumes acute ; lemma 2 - 3 mm long, hairy, awned ; palea up to one - third of the lemma.

Changu ; 1400 - 3700 m.

Fl. & Fr. : Aug. - Sept.

A. sikkimensis Bor in Kew Bull. 1954 : 502. 1954 ; Bor, Grass. Ind. 390. 1960. *A. divaricata* Griseb. in Goett. Nachr. 81. 1868, non Holzman 1800 ; Hook. f., F.B.I. 7 : 258. 1896.

Culms 10 - 15 cm high. Panicles 3 - 5 cm long and broad. Spikelets 1.5 - 1.75 mm long ; glumes unequal ; lemma three-fourth of upper glumes ; palea minute, orbicular.

Ningbil, Karponang ; 2800 - 4000 m.

Fl. & Fr. : July - Aug.

A. stolonifera L., Sp. Pl. 1 : 62. 1753 ; Bor, Grass. Ind. 390. 1960.
A. alba auct. non L. : Hook.f., F.B.I. 7 : 254. 1896.

Culms 15 - 70 cm high. Panicles 3 - 10 cm long. Spikelets 2 - 3 mm long ; glumes acute ; lemma 1.5 - 2.5 mm long ; awnless or with a short awn ; anther 1 - 1.5 mm long.

Sikkim, s.l. ; 1500 - 4000 m (*fide* Hook. f., *op.cit.*)

A. triaristata (Hook.f.) Bor, Grass. Ind. 391. 1960. *Deyeuxia triaristata* Hook.f., F.B.I. 7 : 266. 1896. *Calamagrostis triplifera* Hook. f., F.B.I. 7 : 262. 1896.

Culms 20 - 45 cm high. Panicles of few very long, wide spreading branches. Spikelets *ca* 4 mm long ; glumes glabrous ; lemma with mucros in between two lateral and median awn.

Changu, Sherabthang.

Fl. & Fr. : Aug. - Sep.

ALOPECURUS L.

Alopecurus aequalis Sobol, Fl. Petrop. 16. 1799 ; Bor, Grass. Ind. 392. 1960. *A. aristulatus* Michx., Fl. Bor. Am. 1 : 43. 1803 ; Hook. f., F.B.I. 7 : 238. 1896.

Annulas or perennials. Spike 2.5 - 7.5 cm long. Spikelets *ca* 2.5 mm long ; awn *ca* 2 mm long, median - dorsal.

Gangtok.

Fl. & Fr. : Apr. - Aug.

A. geniculatus L., Sp. Pl. 60. 1753 ; Hook.f., F.B.I. 7 : 239. 1896 ; Bor, Grass. Ind. 393. 1960.

Erect or geniculately ascending perennials. Spikelets strongly compressed, *ca* 2.75 mm long, awn up to 4.5 mm long from the base of lemma.

Sikkim (*fide* Hook. f., *op. cit.*).

ANISELYTRON Merr.

Aniselytron treutleri (O. Ktze.) Sojak in Cas. Nat. Muz. 148. 202. 1980 ; Bennet & Raizada in Ind. Forester 107 : 434. 1981. *Milium treutleri* O. Ktze., Rev. Gen. Pl. 2 : 780. 1891, *p.p.* *Deyeuxia treutleri* (O.Ktze.) Stapf in Hook., Ic. Pl. t. 2396. 1895 ; Hook.f., F.B.I. 7 : 269.

1896. *Aulacolepis treutleri* (O. Ktze.) Hack. in Fedde Rep. Sp. Nov. 3 : 242. 1907 ; Bor, Grass. Ind. 394. 1960.

Culms up to 50 cm high, glabrous. Leaf-blades up to 25 cm long, 1 cm wide. Spikelets 1-flowered, lemma slightly curved.

Lachen ; 2100 - 3500 m.

Fl. & Fr. : Aug. - Sept.

ANTHOXANTHUM L.

Anthoxanthum flexuosa (Hook. f.) Veldk. in Blumea 30 : 347.

1895. *Hierochloe flexuosa* Hook. f., F.B.I. 7 : 222. 1896 ; Bor, Grass. Ind. 441. 1960.

Rhizoma tous perennials. Culms 30 - 75 cm high. Leaves up to 10 cm long, 4 mm broad. Panicles 5 - 8 cm long and 4 mm broad. Spikelets long, pedicelled.

Changu, Bakhim.

Fl. & Fr. : July - Sept.

A. hookeri (Griseb.) Rendle in J. Linn. Soc. 36 : 380. 1904 ; Bor, Grass. Ind. 431. 1960. *Ataxia hookeri* Griseb. in Goett. Nachr. 77. 1868. *Hierochloe hookeri* (Griseb.) Clarke ex Hook. f., F.B.I. 7 : 223. 1896.

Culms 15 - 16 cm high. Spikelets 6 - 8.5 mm long ; short pedicelled.

Thangu, Zemu ; 2600 - 3400 m.

Fl. & Fr. : June - July.

A. sikkimense (Maxim.) Ohwi in Bull. Tokyo Sci. Mus. 18 : 8. 1947 ; Bor, Grass. Ind. 431. 1960. *Hierochloe sikkimensis* Maxim. in Bull. Acad. Sci. St. Petersb. 32 : 262. 1888. *H. gracillima* Hook. f., F.B.I. 7 : 223. 1896.

Slender grass. Spikelets up to 5 mm long, short pedicelled.

Sikkim (*fide* Hook. f., *op. cit.*).

ARUNDINELLA Reddl.

Arundinella bengalensis (Spreng.) Druce in Rep. Bot. Exch. Cl. Brit. Isles 605. 1917 ; Bor, Grass. Ind. 421. 1960. *Panicum bengalense* Spreng., Syst. Veg. 1 : 311. 1852. *Arundinella wallichii* Nees ex Steud., Syn. Gram. 114. 1854 ; Hook. f., F.B.I. 7 : 75. 1896.

Stout perennial. Panicle up to 30 cm long ; branches 2.5 - 4 cm long. Spikelets *ca* 3 mm long, lower floret male, upper bisexual, lemma awned.

Rangeet, Laltung.

Fl. & Fr. : Aug. - Nov.

A. decempedalis (O. Ktze.) Jano. in Fedde Rep. 17 : 84. 1921 ; Bor, Grass. Ind. 422. 1960. *Panicum decempedalis* O. Ktze, Rev. Gen. Pl. 2 : 783. 1891. *Arundinella clarkei* Hook. f., F.B.I. 7 : 75. 1896.

Perennial up to 2 m tall. Panicle up to 45 cm, branches long, filiform, lemma awnless.

Saramsa ; 700 - 1000 m.

Fl. & Fr. : Aug. - Oct.

A. hookeri Munro ex King in Nat. Contr. Univ. Sci. Rep. Ser. B. 2 (3) : 50. 1936 ; Bor, Grass. Ind. 422. 1960. *A. villosa* Steud. var. *himalaica* Hook. f., F.B.I. 7 : 73. 1896.

Densely tufted perennials, up to 45 cm high. Panicles spiciform. Spikelets with bristly hairs on glumes.

Bakhim, Lachung ; 2500 - 3000 m.

Fl. & Fr. : July - Oct.

A. nepalensis Trin., Gram. Panic. 62. 1826 ; Bor, Grass. Ind. 423. 1960. *A. brasiliensis* auct. non Raddl. 1832 ; Hook. f., F.B.I. 7 : 73. 1896.

Perennials with scaly rhizomes. Panicles up to 30 cm long. Spikelets glabrous. Fertile lemma awned.

Singtam.

Fl. & Fr. : Oct. - Nov.

ARUNDO L.

Arundo donax L., Sp. Pl. 81. 1753 ; Hook. f., F.B.I. 7 : 302. 1896 ; Bor, Grass. Ind. 413. 1960.

Perennial with creeping rhizomes and 2 - 4 m high culms. Panicles 30 - 60 cm long.

Gangtok ; *ca* 1700 m.

Fl. & Fr. : Aug. - Dec.

AVENA L.

Avena fatua L. ssp. **fatua** var. **glabrata** Peterm., Fl. Bieuitz. 13 : 1841 ; Bor, Grass. Ind. 434. 1960. *Avena sativa* L. var. *sericea* Hook. f., F.B.I. 7 : 275. 1896.

Annual. Culms few or solitary, 30 - 80 cm high, stout. Panicle open or contracted. Spikelets 2.2 - 2.8 cm long, with 2 -florets. Lemma bearded at base only.

Yamthang ; 3200 m (*fide* Hook. f., *op. cit.*)

A. sativa L., Sp. Pl. 79. 1753 ; Hook. f., F.B.I. 7 : 275. 1896 ; Bor, Grass. Ind. 434. 1960.

Culms simple. Panicle open. Spikelets 2.2 - 3 cm long. Lemmas glabrous.

Chungthang. Weed in the cultivated fields.

Fl. & Fr. : Mar. - May.

BRACHYPODIUM P. Beauv.

Brachypodium sylvaticum (Huds.) P. Beauv., Ess. Agrost. 101. 155. 1812 ; Hook. f., F.B.I. 7 : 362. 1896 ; Bor, Grass. Ind. 450. 1960. *Festuca sylvatica* Huds., Fl. Angl. 1 : 38. 1762. *Brachypodium wattii* Clarke in J. Linn. Soc. 25 : 90. t. 40. 1887.

Perennials, culms up to 1.5 m high. Raceme up to 15 cm long, nodding. Spikelets subsessile, cylindrical at first, becoming compressed later.

Gangtok.

Fl. & Fr. : Apr - Sept.

BRIZA L.

Briza media L., Sp. Pl. 70. 1753 ; Hook. f., F.B.I. 7 : 336. 1896 ; Bor, Grass. Ind. 528. 1960.

Perennial. Culms 30 - 100 cm high, very slender. Inflorescence an erect spreading pyramidal panicle.

Gnaton (*fide* Hook. f., *op. cit.*)

BROMUS L.

Bromus himalaicus Stapf in Hook. f., F.B.I. 7 : 358. 1896 ; Bor, Grass. Ind. 453. 1960. *Zerna himalaica* (Stapf) Henr. in Blume 4(3) : 499. 1941.

Culms up to 70 cm high. Leaves linear, acuminate, pubescent. Spikelets linear-oblong, up to 2.5 cm long ; lemma glabrous or hairy along the margins, awn slender.

Sikkim (*fide* Hook. f., *op. cit.* ; Bor, *op. cit.*).

3000 -3700 m.

Fl. & Fr. : June - Oct.

B. porphyranthus T.A. Cope in Nasir & Ali, Fl. Pak. 143 : 574. 1892. *B. himalaicus* Stapf var. *grandis* Stapf in Hook. f., F.B.I. 359. 1896. Bor, Grass. Ind. 454. 1960. *B. grandis* (Stapf) Meld. in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 125. 1978, *non* Hitchc. 1912.

Loosely tufted perennial. Panicles branched, simple, ending in a solitary spikelet. Spikelets 20 - 25 mm long, lemmas densely hairy, lowest usually 13 - 15 mm long, awns usually robust.

Tsangu (Thangu) ; 3300 - 3600 m.

B. ramosus Huds., Fl. Angl. 40.1762. Bor, Grass. Ind. 456. 1960 ; Hook. f., F.B.I. 7 : 358. 1896.

Culms up to 1 m high. Panicle wide and loose, the lowest branch with a ciliate scale at base. Spikelets up to 4 cm long ; lemmas with straight awn.

Lachen.

Fl. & Fr. : July - Aug.

B. staintonii Melderis in Hara *et al.* Enum. Fl. Pl. Nepal 1 : 125. 1978. *Bromus ramosus sensu* Bor, Grass. Ind. 456. 1960, *non* Huds. 1853.

Sikkim, s.l.

CALAMAGROSTIS Adans.

Calamagrostis emodensis Griseb. in Goett. Nachr. 80. 1868 ; Hook. f., F.B.I. 7 : 261. 1896 ; Bor, Grass. Ind. 395. 1960.

Tufted perennial, 80 - 130 cm high. Panicles nodding, 13 - 35 cm long. Spikelets ca 6 mm long, lemma with excluded awn ; palea almost as long as lemma.

Gangtok, Zemu Valley ; 2200 - 4000 m.

Fl. & Fr. : July - Oct.

C. pseudophragmites (Hall. f.) Koeler, Descr. Gram. 106. 1802 ; Bor, Grass. Ind. 396. 1960. *Arundo pseudophragmites* Hall. f. in Roem., Arch. für. Bot. 1 : 11. 1797. *Calamagrostis littorea* (Schrad.) P. Beauv., Ess. Agrost. 15. 157. 1812 ; Hook. f., F.B.I. 7 : 261. 1896.

Perennials, with creeping rhizomes ; culms 40 - 150 cm high. Panicles lax, 10 - 14 cm long. Spikelets 5 - 9 mm long, lemma with included or shortly excluded awn, palea half as long as lemma.

Lachen, Chungthang ; 1500 - 2700 m.

Fl. & Fr. : July - Sept.

CATABROSA P. Beauv.

Catabrosa sikkimensis Stapf in Hook. f., F.B.I. 7 : 311. 1896.

Stoloniferous perennial, culms up to 15 cm long, slender. Panicles ovate to oblong, 4 - 7 cm long. Spikelets 2 - 2.5 mm long, purple.

Lhonk valley.

Fl. & Fr. : June - Aug.

CENTOTHECA Desr.

Centotheca lappacea (L.) Desv. in Nouv. Bull. Sci. Soc. Philom. Paris. 2 : 189. 1818 ; Hook. f., F.B.I. 7 : 32. 1896 ; Bor, Grass. Ind. 457. 1960. *Cenchrus lappaceus* L., Sp. Pl. ed. 2. 1488. 1763.

var. lappacea

Culms 15 -90 cm high. Leaves ovate-lanceolate. Panicles loose. Spikelets ovate-oblong, glumes keeled, lemmas with bristles near the margins.

Teesta (West Bengal).

COELACHNE R. Br.

Coelachne simpliciuscula (Steud.) Benth. in J. Linn. Soc. 19 : 93. 1881 ; Bor, Grass. Ind. 576. 1860. *Panicum simpliciusculum* Steud., Syn. Pl. Glum. 1 : 96. 1854. *C. pulchella* sensu Hook. f., F.B.I. 7 : 270. 1896, *non* R. Br. 1810.

Annuals. Culms 15 - 40 cm high, slender. Panicles, simple, spike - like. Spikelets globose.

In marshy places near Gangtok.

Fl. & Fr. : Throughout the year.

COLPODIUM Trin.

Colpodium wallichii (Hook. f.,) Bor in Kew Bull. 1953 : 270. 1953 ; Bor, Grass. Ind. 530. 1960. *Catabrosa wallichii* Hook. f., F.B.I. 7 : 312. 1896.

Culms 15 - 40 cm high. Leaf-blades 3 - 7 mm long. Panicle narrow. Spikelets pendulous, glumes oblong ; lemma with shortly hairy outer nerves.

Kankola pass (*fide* Hook. f., *op. cit.*)

CYATHOPUS Stapf

Cyathopus sikkimensis Stapf in Hook., Ic. Pl. t. 2395. 1895 ; Hook. f., F.B.I. 7 : 240. 1896 ; Bor, Grass. Ind. 565. 1960.

Culms about 1 m high. Panicles effuse ; spikelets scaberulous; glumes beaked.

Lachung valley ; ca 3300 m.

Fl. & Fr. : Oct. - Dec.

CYNODON L.

Cynodon dactylon (L.) Pers., Syn. Pl. 1 : 85. 1805 ; Hook. f., F.B.I. 7 : 288. 1896 ; Bor, Grass. Ind. 469. 1960. *Panicum dactylon* L., Sp. Pl. 58. 1753.

Creeping perennial. Inflorescence of 3 - 6 ascending to spreading digitate slender spikes.

Gangtok ; up to 2000 m.

Fl. & Fr. : Throughout the year.

DACTYLOCTENIUM Willd.

Dactyloctenium aegyptium (L.) Beauv., Ess. Agrost. Expl. Pl. 15. 1812 ; Bor, Grass. Ind. 489. t. 54. 1960. *Cynosurus aegyptius* L., Sp. Pl. 72. 1753. *Eleusine aegyptia* (L.) Desf., Fl. Atlant. 1 : 85. 1789 ; Hook. f., F.B.I. 7 : 295. 1896.

Annuals. Culms up to 70 cm high. Inflorescence of 2 - 6 mm linear to oblong spikes, naked at the tip for 2 - 3 mm. Spikelets with upper glume, mucronate or awned.

Sikkim. Only one sheet without locality (LBG) represents its distribution is Sikkim.

DANTHONIA DC.

Danthonia cumminsii Hook. f., F.B.I. 7 : 282. 1896.
D.cachemeryiana sensu Hook. f., F.B.I. 7 : 281. 1896, non Jaub. & Spach. 1851. *P.jacquemotii* Bor in Kew Bull. 7 : 80. 1952 ; Bor, Grass. Ind. 479. 1960.

Perennials with woody rootstock ; culms up to 60 cm high. Panicles 10 - 30 cm long. Spikelets 15 - 25 mm long.

Jongu, Tangu, Gnatong, Legyep.

Fl. & Fr. : July - Sept.

DESCHAMPSIA L.

Deschampsia caespitosa (L.) P. Beauv., Ess. Agrost. 91, t. 18. f. 2. 1812 ; Hook. f., F.B.I. 7 : 273. 1896 ; Bor, Grass. Ind. 435. 1960. *Aira caespitosa* L., Sp. Pl. 64. 1753.

Densely tufted perennial. Panicles oblong. Spikelets narrowly oblong, 2-flowered, lemma awned from near the base.

Zemu valley, Lhonak, Yumthang, Yakla ; 3600 - 4800 m.

Fl. & Fr. : June - Nov.

DEYEUSIA P. Beauv.

Deyeusia abnormis Hook. f., F.B.I. 7 : 268. 1896 ; Korthof & Veld. in Gard. Bull. Sing. 37 : 218. 1985.

Densely tufted perennial. Panicles contracted or effuse ; branches and branchlets capillary. Callus hairs half the length of lemma, awn absent.

Lanchung valley (*fide* Hook. f., *op. cit.*).

D. debilis (Hook. f.) Veldk. in Gard. Bull. Sing. 37 : 220. 1985. *Calamagrostis debelis* Hook. f., F.B.I. 7 : 262. 1896. *Agrostis debilis* (Hook. f.) Bor, Grass. Ind. 387. 1960, non Poir. 1810. *A. neodebilis* Bennet & Raizada in Ind. For. 107 : 433. 1981.

Tufted perennial. Panicles narrow. Spikelets purplish brown, glumes unequal, lemma shorter than glumes, anthers very small.

Chola, 3700 m (*fide* Hook. f., *op. cit.*).

D. nivicola Hook. f., F.B.I. 7 : 267. 1896 ; Bor, Grass. Ind. 399. 1960.

Densely tufted perennial. Panicles 2.5 - 4 cm long, contracted, spiciform. Spikelets 6 mm long, glumes unequal, lanceolate-acuminate, lemma shorter than glumes, awn basal, included.

Momay - Samdong (*fide* Hook. *f.*, *op. cit.*).

D. pulchella (Griseb.) Hook. *f.*, F.B.I. 7 : 268. 1896 ; Bor, Grass. Ind. 399. 1960. *Calamagrostis pulchella* Griseb. in Goett. Nachr. 78. 1868. *Calamagrostis lahulensis* G. Singh in Taxon 33 : 94. 1984.

Tufted perennial. Panicles oblong up to 5 cm long, branches scaberulous. Spikelets 4 - 5 mm long, callus shortly bearded, glumes unequal, lemma with submedian awn.

Changu, Momay - Samdong ; 3500 - 4800 m.

Fl. & Fr. : July - Oct.

D. scabrescens (Griseb.) Duthie in Atkins., Gaz. N.W. India 628. 1882. Hook. *f.*, F.B.I. 7 : 267. 1896 ; Bor, Grass. Ind. 399. 1960. *Calamagrostis scabrescens* Griseb. in Goett. Nachr. 79. 1868. *Calamagrostis filiformis* Griseb. *l.c.* 79. 1868. *Deyeusia filiformis* (Griseb.) Hook. *f.*, F.B.I. 7 : 268. 1896. *D. scabrescens* var. *humilis* Hook. *f.*, *l.c.* 1896.

Perennials up to 1.5 m high. Panicle up to 20 cm long, contracted. Spikelets 6 - 6.5 mm long ; glumes unequal, lemma minutely scabrid, awn inserted about one third below tip, scabrid.

Changu, Zemu valley ; 2700 - 3900 m.

Fl. & Fr. : Aug. - Oct.

D. tibetic Bor in Kew Bull. 1949 ; 66. 1949 ; Bor, Grass. Ind. 400. 1960. *Calamagrostis tibetica* (Bor) G. Singh in Taxon 32 : 95. 1984.

Perennial up to 20 cm high, pilose and angular below panicle. Panicle compact, spiciform. Spikelets purple, glumes equal, pilose.

Chakalunga ; ca 5000 m.

Fl. & Fr. : June.

ELEUSINE Gaertn.

Eleusine coracana (L.) Gaertn., Fruct. 1 : 8. t.1.f. 11. 1789 ; Hook. *f.*, F.B.I. 7 : 294. 1896 ; Bor, Grass. Ind. 492. 1960. *Cynosurus coracanus* L., Syst. Nat. ed. 10, 2 : 875. 1759. KODO (Nep.).

Tufted annual ; culms 15 - 85 cm high. Leaf-blades usually folded. Inflorescence digitate, of 1 - 10 slender ascending spikes. Spikelets 3 - 9 - flowered, elliptic.

Cultivated.

E. indica (L.) Gaertn., Fruct. 1 : 8. 1789 ; Hook. f., F.B.I. 7 : 293. 1896 ; Bor, Grass. Ind. 493. 1960. *Cynosurus indicus* L., Sp. Pl. 72. 1753.

Tufted annual ; culms 15 - 85 cm high. Leaf-blades usually folded. Inflorescence digitate, of 1 - 10 slender ascending spikes. Spikelets 3 - 9-flowered, elliptic.

Throughout subtropical and temperate Sikkim as a weed.

ELYMUS L.

Elymus nutans Griseb. in Goett. Nach. 72. 1868 ; Bor, Grass. Ind. 670. 1960. *E. sibiricus* auct. non L. : Hook. f., F.B.I. 7 : 373. 1896, p.p.

Tufted perennial. Spike often flexuous, usually nodding, 5 - 15 cm long. Spikelets paired at the nodes of rachis.

Naku - la ; 4200 - 4700 m.

Fl. & Fr. : Aug. - Nov.

E. semicostatus (Nees ex Steud.) Meld. in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 132. 1978. *Triticum semicostatum* Nees ex Steud., Syn. Glum. 1 : 346. 1854. *Agropyron semicostatus* (Nees ex Steud.) Boiss., Fl. Or. 5 : 662. 1884 ; Hook. f., F.B.I. 7 : 369. 1896 ; Bor, Grass. Ind. 665. 1960.

Tufted perennial, erect or geniculately ascending. Spike 8 - 30 cm long. Spikelets 6 - 8-flowered, glumes unequal, palea shorter than lemma.

Sikkim (*fide* Hook. f., *op. cit.* & Bor, *op. cit.*).

E. sikkimensis (Melderis) Melderis in Hara *et al.*, Enum. Fl. Pl. Nepal 1 : 132. 1978. *Agropyron sikkimense* Melderis in Bor, Grass. Ind. 666. 694. 1960.

Perennial, 70 - 80 cm high ; culms scabrous. Spike lax. Spikelets 3 - 6-flowered, palea as long or longer than lemma.

Lachen, Momay - Samdong ; ca 4500 m.

Fl. & Fr. : May-Aug.

ERAGROSTIS Wolf.

Eragrostis atrovirens (Desf.) Trin. ex Steud., Nom. Bot. ed. 2. 1 : 562. 1840 ; Bor, Grass. Ind. 503. 1960. *Poa atrovirens* Desf., Fl. Atlant. 1 : 73, t. 14. 1798.

Tufted perennial. Panicles ovate or oblong ; spikelets 6 - 59 flowered, narrowly oblong, 3 - 20 mm long, breaking up from the base.

Jora ; 500 - 1000 m.

Fl. & Fr. : Almost throughout the year.

E. coarctata Stapf in Hook. f., F.B.I. 7 : 313. 1896 ; Bor. Grass. Ind. 507. 1960.

Perennial ; culms tufted. Inflorescence an elongate, subspiciform panicle. Spikelets on short pedicels, glumes equal, lemmas ciliate on the margins, palea strongly ciliate on the keels.

Rongli, Rongpo ; 300 - 700 m.

Fl. & Fr. : May.

E. curvula (Schrad.) Nees, Fl. Afr. Austr. 397. 1841 ; Bor, Grass. Ind. 507. 1960. *Poa curvula* Schrad. in Goett. Anz. Ges. Wiss. 3 : 2073. 1821.

Densely tufted perennial ; leaves usually rolled or filiform. Panicle spreading. Spikelets breaking up from the base.

Gangtok. Introduced.

E. ferruginea (Thunb.) P. Beauv., Ess. Agrost. 71. 1812 ; Stapf in Hook. f., F.B.I. 7 : 324. 1896 ; Bor, Grass. Ind. 508. 1960. *Poa ferruginea* Thunb., Fl. Jap. 50. 1784.

Perennial up to 45 cm high. Panicles 10 - 50 cm long, oblong. Spikelets 6 mm long, pedicels with glandular band ; lemma rounded on the back, palea shorter than lemma.

Lindok.

Fl. & Fr. : May - June.

E. gangetic (Roxb.) Steud., Syn. Pl. Glum. 1 : 266. 1854 ; Bor, Grass. Ind. 508. 1960. *Poa gangetica* Roxb., Fl. Ind. 1 : 321. 1820. *Eragrostis stenophylla* Hochst. ex Miq. in Verh. Konink -Ned. Inst. 3, 4 : 39. 1851. Stapf in Hook. f., F.B.I. 7 : 318. 1896.

Annual, tufted grass. Panicles oblong, nodding. Spikelets linear-oblong, 10 - 30-flowered, lemmas broadly ovate, subacute, palea deciduous.

Gyalshing, Rishi, Rhenock ; 400 - 1300 m.

Fl. & Fr. : Almost throughout the year.

E. japonica (Thunb.) Trin. in Mem. Acad. Sci. Petersb. 6, 1 : 405. 1831. Bor, Grass. Ind. 509. 1960. *Poa japonica* Thunb., Fl. Jap. 51. 1784. *Eragrostis diarrhena* (Schult.) Steud., Syn. Pl. Glum. 1 : 266. 1854 ; Bor, Grass. Ind. 507. 1960. *E. diplachnoides* Steud., l.c. 268 ; Bor, Grass. Ind. 508. 1960. *E. interrupta* var. *tenuissima* Stapf in Hook. f., F.B.I. 7 : 316. 1896.

Annual or short-lived perennial. Panicles linear-lanceolate to narrowly ovate ; 4 - 5 cm long. Spikelets 4 - 14-flowered, ovate-oblong, 1.3 - 3 mm long, breaking up from the apex.

Rangeet valley.

Fl. & Fr. : Aug. - Oct.

E. minor Host, Gram. Austr. 4 : 15. 1809 ; Stapf in Hook. f., F.B.I. 7 : 321. 1896. *E. poaeoides* P. Beauv., Ess. Agrost. 162. 1812 ; Bor, Grass. Ind. 512. 1960. *E. pappiana* Chiov. in Ann. Ist. Bot. Roma 8 : 371. 1908 ; Bor, Grass. Ind. 511. 1960 ; Jain in Bull. Bot. Surv. Ind. 10 : 161. 1969.

Loosely tufted annual, culms 6 - 60 cm high, ascending. Panicles ovate. Spikelets narrowly oblong or almost linear, breaking up from base, glumes subequal, often glandular on the keel.

Gangtok.

Fl. & Fr. : June - Sept.

E. nigra Nees ex Steud., Syn. Pl. Glum. 1 : 267. 1854 ; Stapf in Hook. f., F.B.I. 7 : 324. 1896 ; Bor, Grass. Ind. 511. 1960.

Perennial, leafy at base. Panicles large spreading, lax. Spikelets linear to oblong, glumes subequal, palea persistent.

Chungthang, Lachen ; 900 - 3000 m.

Fl. & Fr. : June - Aug.

E. nutans (Retz.) Nees ex Steud., Nom. Bot. ed. 2, 1 : 563. 1840 ; Bor, Grass. Ind. 511. 1960. *Poa nutans* Retz., Obs. Bot. 4 : 19. 1786. *Eragrostis stenophylla* auct. non Miq : Hook. f., F.B.I. 7 : 318. 1896. p.p.

Perennial ; culms densely tufted, 30 - 40 cm high. Panicles ovate or oblong or sublinear, branches distant, pedicels of spikelets filiform. Spikelets 4 - 6 mm long, linear-oblong, palea caducous, keels scabrid.

Deorali, Gangtok.

Fl. & Fr. : Sept. - Nov.

E. papposa (Roem. & Schult.) Steud., Nom. Bot. ed. 2, 1 : 564. 1840 ; Hook *f.*, F.B.I. 7 : 322. 1896 ; Bor, Grass. Ind. 511. 1960. *Megastachye papposa* Roem. & Schult., Syst. Veg. 2 : 585. 1817.

Tufted perennial ; culms 10 - 40 cm high. Panicles ovate. Spikelets grey-green to purplish green, glumes unequal, paleas minutely scaberulous on the keels, persistent.

Gyalshing, Soreng.

Fl. & Fr. : Apr. - Oct.

E. pilosa (L.) P. Beauv., Ess. Agrost. 71. 162. 175. 1812 ; Stapf in Hook. *f.*, F.B.I. 7 : 323. 1896 ; Bor, Grass. Ind. 512. 1960. *Poa pilosa* L., Sp. Pl. 68. 1753.

Loosely tufted annual. Panicles elliptic to ovate, 4 - 25 cm long, open, usually with a few long hairs in the axis of lower baranches.

Sikkim, *s.l.*

Fl. & Fr. : July - Oct.

E. tenella (L.) P. Beauv. ex R. & S., Syst. Veg. 2 : 576. 1817 ; Stapf in Hook. *f.*, F.B.I. 7 : 315. 1896 ; Bor, Grass. Ind. 513. 1960 (incl. vars.). *Poa tenella* L., Sp. Pl. 69. 1753. *E. tenella* var. *plumosa* (Retz.) Stapf in Hook. *f.*, F.B.I. 7 : 315. 1896.

Tufted annual. Panicles elliptic or pyramidal. Spikelets 4 - 8-flowered, 1.5 - 2.5 mm long, breaking from the apex, palea ciliate on keels.

Rishikhola, Chakung ; 200 - 1000 m.

Fl. & Fr. : Mar. - Sept.

E. tremula (Lam.) Steud., Syn. Pl. Glum. 1 : 269. 1854 ; Stapf in Hook. *f.*, F.B.I. 7 : 320. 1896 ; Bor, Grass. Ind. 514. 1960. *Poa tremula* Lam., Tab. Encycl. 1 : 185. 1791.

Loosely tufted annual ; culms 30 - 100 cm high. Panicle ovate, 7-30 cm long, loose and open. Spikelets linear, 5 - 25 mm long, 1.5 - 2 mm wide, glumes unequal, palea scabrid on keels, persistent.

Sikkim, s.l.

Fl. & Fr. : Mar. - Nov.

E. unioloides (Retz.) Steud., *Syn. Pl. Glum.* 1 : 264. 1854 ; Bor, *Grass. Ind.* 515. 1960. *Poa unioloides* Retz., *Obs. Bot.* 5 : 19. 1789. *Eragrostis amabilis* Stapf in *Hook. f.*, *F.B.I.* 7 : 317. 1896.

Tufted annual or short lived perennial. Spikelets 11 - 80-flowered, ovoid to oblong, 3.5 - 16 mm long, 2.5 - 4 mm wide, florets slightly overlapping, breaking from base, glumes unequal, narrowly lanceolate, 1-nerved, palea deciduous.

Gangtok, Rangpo ; 400 - 1700 m.

E. viscosa (Retz.) Trin. in *Mem. Acad. Sci. Petersb. Ser. 6*, 1 : 397. 1830 ; Bor, *Grass. Ind.* 515. 1960. *Poa viscosa* Retz., *Obs. Bot.* 4 : 20. 1986. *Eragrostis tenella* var. *viscosa* (Retz.) Stapf in *Hook. f.*, *F.B.I.* 7 : 315. 1896.

Tufted annual ; culms 10 - 30 cm high, glandular. Panicle dense, cylindrical or oblong, 7 - 18 cm long, up to 3.5 cm broad. Spikelets 5 - 20-flowered, often purplish, about 3 mm long, glumes almost equal, obtuse, lemmas broadly elliptic, obtuse or rounded.

Teesta ; ca 300 m.

Fl. & Fr. : April.

FESTUCA L.

Festuca coelestis (St.Yves) Krecz. & Bobrov in Kum. *Fl. USSR* 2 : 514. 770. 1934 ; T.A. Cope in Nasir & Ali, *Fl. Pak.* 143. 359. 1982. *Festuca ovina* L. ssp. *coelestris* St. Yves in Candollea 3 : 376. 1928 ; Bor, *Grass. Ind.* 539. 1960. *F. valesiaca* var. *tibetica* Stapf in *Hook. f.*, *F.B.I.* 7 : 349. 1896 ; Bor, *Grass. Ind.* 542. 1960.

A dwarf tufted perennial. Culms slender. Leaves folded, setaceous. Spikelets 3 - 4-flowered, glumes 1-nerved, lemmas scabrid towards tip with 1 - 2 mm long awn, ovary glabrous.

Donkia La ; 5100 m (*fide Hook. f.*, *op. cit* ; Bor, *op. cit*).

F. cumminisii Stapf in *Hook. f.*, *F.B.I.* 7 : 349. 1896 ; Bor, *Grass. Ind.* 538. 1960.

Tufted perennials. Culms slender. Panicles nodding or flexuous, 5 - 10 cm long. Spikelets pale green or purple, glumes unequal, lemma awned, palea scabrid.

Changu, Sherabthang, Chamnago.

Fl. & Fr. : July - Aug.

F. leptobogon Stapf in Hook. *f.*, F.B.I. 7 : 354. 1896 ; Bor, Grass. Ind. 538. 1960.

Tufted perennial. Panicles 15 - 30 cm long ; open or contracted. Spikelets 2 - 3 - flowered, glumes unequal, lemmas 7 - 8 mm long, 5-nerved, 2-lobed at apex, awn straight, scabrid, 7 mm long. Ovary minutely hispid on the top.

Lachen.

Fl. & Fr. : July.

F. parvigluma Steud., Syn. Pl. Glum. 1 : 305. 1854 ; Bor, Grass. Ind. 540. 1960.

Glumes short, broadly oblong, obtuse, lemmas entire at tip.

Yatung (Bor, *op. cit.*).

F. polycolea Stapf in Hook. *f.*, F.B.I. 7 : 349. 1896 ; Bor, Grass. Ind. 540. 1960. *F. ovina* ssp. *polycolea* (Stapf) St. Yves in Candollea 3 : 373. 1928 ; Bor, Grass. Ind. 540. 1960.

var. **polycolea**

Densely tufted perennial. Leaves rigid, setaceous. Panicles ovate or narrow, 5 - 8 cm long, erect, rigid. Spikelets 8 - 12 mm long, glumes scabrid, lemma involute, awned, ovary glabrous.

Dzongri, Yakla, Zemu ; 3000 - 5000 m.

Fl. & Fr. : July - Oct.

var. **brevis** Stapf in Hook. *f.*, F.B.I. 7 : 350. 1896 ; Bor, Grass. Ind. 540. 1960.

Culms 10 - 15 cm long. Panicles 1.5 - 3.5 cm long, contracted.

Lachen (*fide* Hook. *f.*, *op.cit.*).

F. rubra L., Sp. Pl. 74. 1753 ; Stapf in Hook. *f.*, F.B.I. 7 : 352. 1896 ; Bor, Grass. Ind. 540. 1960.

Culms 15 - 90 cm high. Leaves folded, setaceous. Panicles loose. Spikelets 5 - 14 mm long, lemma 4 - 7.5 mm long, scaberulous towards tip, ovary glabrous or sparserly hairy.

Tosa ; 3500 - 4200 m.

Fl. & Fr. : July - Aug.

F. undata Stapf in Hook. *f.*, F.B.I. 7 : 350. 1896 ; Bor, Grass. Ind. 542. 1960.

var. undata

Tufted perennial, 30 - 60 cm high. Leaves subsetaceous. Culms scabrid below the panicle, awn of lemma 1.5 mm long.

Gnathang ; 3500 m.

Fl. & Fr. : Sept.

var. aristata Stapf in Hook. *f.*, F.B.I. 7 : 387. 1896 ; Bor, Grass. Ind. 542. 1960.

Awn slender, longer than lemma ; anthers 1.3 - 1.5 mm long. Ovary glabrous.

Chamnago ; 3200 - 4000 m.

Fl. & Fr. : July - Aug.

F. valesiaca Schleich ex Gaud., Agrost. Helvet. 1 : 242. 1811 ; Bor, Grass. Ind. 542. 1960 ; Hook. *f.*, F.B.I. 7 : 348. 1896. *F. ovina* L. ssp. *sulcata* Hack. var. *valesiaca* (Schleich ex Gaud.) Koch., Syn. ed. 1 : 812. 1837 ; Bor, Grass. Ind. 540. 1960.

Tufted perennial ; culms 10 - 40 cm high, erect, slender. Leaves folded, setaceous. Panicles narrowly oblong, contracted. Spikelets 5 - 7-flowered, lemma scabrid in the upper part, awn 1 - 3 mm long.

Sikkim (*fide* Hook. *f.*, *op.cit.*).

GARNOTIA Brongn.

Garnotia acutigluma (Steud.) Ohwi in Bot. Mag. Tokyo 55 : 393. 1941 ; Gould in Kew Bull. 27 : 540. 1972. V. Prakash & Jain in Fasc. Fl. Ind. 3 : 5. 1979. *Urachne acutigluma* Steud., Syn. Pl. Glum. 1 : 121. 1854. *Garnotia stricta* auct. non Brongn. 1832 : Hook. *f.*, F.B.I. 7 : 243. 1896, *p.p.* ; Bor, Grass. Ind. 569. 1960, *p.p.*

Perennial ; culms 25 - 70 cm long. Spikelets with bearded base, glumes awned, lemma with straight awn.

Burtuk, Rongli, Yoksum.

Fl. & Fr. : Aug. - Nov.

var. logiseta (Hack.) Ved Prakash & Jain in Fasc. Fl. Ind. 3 : 6. 1979.

G. polypogonoides Munro ex Oliver in Hook. f., Ic. Pl. 15 : 64. t. 1481. 1885 ; Hook. f., F.B.I. 7 : 243. 1896 ; V. Prakash & Jain, Fasc. Fl. Ind. 3 : 10. 1978.

Perennial ; culms erect from decumbent base. Panicles 8 - 18 cm long. Spikelets conspicuously bearded at base, lemma with geniculate, 13 - 25 mm long awn.

Neebay ; 2000 - 3000 m.

Fl. & Fr. : Sept. - Dec.

G. tenella (Arn. ex Miq.) Jan. in Fedde Repert. 17 : 86. 1921 ; Gould in Kew Bull. 27 ; 528. 1972. *Berghanisa tenella* Arn. ex. Miq. in Vash. Konink. Nederl. Inst. 3, 4 : 34. 1851. *Garnotia stricta* sensu Hook. f., F.B.I. 7 : 243. 1896 ; Bor, Grass. Ind. 569. 1960, *non* Brongn. 1831.

Annuals. Leaf-sheaths glabrous. Panicles, 3 - 29 cm long, contracted. Spikelets scabrid ; lemma with geniculate, 5 - 12 mm long awn.

Richingpon to Gayzing.

Fl. & Fr. : Aug. - Dec.

GLYCERIA R. Br.

Glyceria tonglensis C.B. Clarke in J. Linn. Soc. 15 : 19. 1876 ; Hook. f., F.B.I. 7 : 346. 1896 ; Bor, Grass. Ind. 571. 1960. *G. kasmiriensis* L. Kelso in Rhodora 37 : 262. 1935.

Tufted or creeping perennial ; culms 20 - 50 cm high. Panicles 10 - 156 cm long. Spikelets ovate, 4 - 6-flowered, glumes acute, lemma obtuse.

Chakung Chu.

Fl. & Fr. : July - Sept.

HELIOTRICHON Roem. & Schult.

Helictotrichon virescens (Nees ex Steud.) Henr. in Blumea 3 : 452. 1940 ; Bor, Grass. Ind. 439. 1960 ; Sevenster & Veldkemp in Blumea 28 : 338. 1983. *Trisetum virescens* Nees ex Steud., Syn. Pl. Glum. 1 : 226. 1854. *Avena aspera* Munro ex Thw., Enum. Pl. Zeyl. 372. 1864 ; Hook. f., F.B.I. 7 : 276. 1896 (incl. vars.). *Helictotrichon sperum* (Munro ex Thw.) Bor in Ind. For. Rec. n.s. Bot. 68. 1938 ; Bor, Grass. Ind. 438. 1960. *H. parviflorum* (Hook. f.) Bor in Kew Bull. 1951 ; 445. 1952 ; Bor, Grass. Ind. 439. 1960.

Tufted perennials ; culms 40 - 120 cm high. Leaves flat. Panicles loose. Spikelets 3-flowered.

Gnatong, Tallum - Samdong, Zemu Valley ; 2100 - 4400 m.

Fl. & Fr. : July - Aug.

HOLCUS L.

Holcus lanatus L., Sp. Pl. 1048. 1753 ; Hook. f., F.B.I. 7 : 274. 1896 ; Bor, Grass. Ind. 443. 1960.

Annual or perennial, up to 1 m high. Leaves flat. Panicles laterally compressed.

Sikkim, s. l. Naturalized in Darjeeling.

HORDEUM L.

Hordeum aegiceras Nees ex Rolye, Ill. Bot. Himal. 418. t. 17.f. 2. 1836 ; Bor, Grass. Ind. 675. 1960. *H. vulgare* ssp. *aegiceras* (Nees ex Royle) A. Love in Fedde Repert. 95 : 435. 1984. *H. vulgare* var. *aegiceras* Aitch. In Hook. f., F.B.I. 7 : 371. 1896.

Erect annual. Leaves flat. Spikelets one flowered, lemma produced into a dilated 3-fid appendage.

Sikkim. Cultivated.

H. vulgare L., Sp. Pl. 84. 1753 ; Hook. f., F.B.I. 7 : 371. 1896 ; Bor, Grass. Ind. 677. 1960.

ssp. *vulgare*

Erect annual grass. Leaves flat. Spikelets two ranked.

Sikkim. Cultivated.

ssp. *agriocriton* (Aberg) A. Love in Fedde Repert. 95 : 435. 1984. *Hordeum agriocriton* Aberg in Ann. Agri. Coll. Sweden 6 : 159. 1938. Bor, Grass. Ind. 676. 1960.

Notes : It is believed to be the ancestral form of the cultivated Barley. May be found in Sikkim (*cf.* Bor, *op. cit.*).

ISACHNE R. Br.

Isachne albens Trin., Sp. Gram. 1. Ic. t. 85. 1826 ; Hook. f., F.B.I. 7 : 22. 1896 ; Bor, Grass. Ind. 579. 1960.

Tall perennials ; culms erect or spreading, 30 - 120 cm high. Spikelets 1.5 - 16 mm long, florets equal in size and texture, glumes glabrous.

Throughout subtropical Sikkim.

Fl. & Fr. : Almost throughout the year.

I. clarkei Hook. f., F.B.I. 7 : 24. 1896 ; Bor, Grass. Ind. 579. 1960.

Annuals ; culms slender, often rooting at nodes. Spikelets up to 1 mm long, pedicels longer than spikelets, lemmas involute, scabrid.

Lachung valley ; 1500 - 3300 m.

Fl. & Fr. : Aug. - Dec.

I. himalaica Hook. f., F.B.I. 7 : 23. 1896 ; Bor, Grass. Ind. 581. 1960.

Perennials ; culms 30 - 60 cm high, erect or spreading. Panicles pyramidal. Spikelets 1.8 - 2.1 mm long, ellipsoid to subglobose on long capillary, glandular pedicels, lemmas glabrous.

Sikkim, s.l.

Fl. & Fr. : Aug. - Dec.

I. miliacea Roth ex R. & S., Syst. Veg. 2 : 476. 1817 ; Hook. f., F.B.I. 7 : 25. 1896 ; Bor, Grass. Ind. 582. 1960.

Annuals ; culms up to 20 cm tall. Leaves small, smooth. Spikelets ca 2 mm long, florets unequal in size and of different textures.

Gangtok ; 700 - 1700 m.

Fl. & Fr. : Aug. - Nov.

I. sikkimensis Bor in Kew Bull. 1949 ; 115. 1949 & Bor, Grass. Ind. 582. 1960.

Annual ; culms up to 40 cm tall. Panicles 5 - 10 cm long, effuse, branches flexuous, eglandular. Spikelets ca 2 cm long, ellipsoid, smooth, florets of similar texture.

Karponang, Lachung.

Fl. & Fr. : July - Sept.

LOLIUM L.

Lolium perenne L., Sp. Pl. 83. 1753 ; Hook. f., F.B.I. 7 : 365. 1896 ; Bor, Grass. Ind. 545. 1960.

Stoloniferous perennials. Leaves linear, complicate. Spikelets oblong, 3 - 11-fid.

Sikkim, s.l. (*fide Hook. f., op. cit.* ; Bor, *op. cit.*).

MUHLENBERGIA Schreb.

Muhlenbergia huegelii Trin. in Mem. Acad. Sci. Petersb. Ser. 6, 4 : 41. 198. 1841 ; Bor, Grass. Ind. 401. 1960. *M. viridissima* Nees ex Steud., Syn. Pl. Glum. 1 : 178. 1854 ; Hook. f., F.B.I. 7 : 259. 1896 ; Hsu in Hara, Fl. E. Himal. 368. 1966.

Annual or short perennials. Panicles elongate, branches and pedicels slender. Spikelets awned, glumes 1-nerved, lemmas twice as long as the glumes.

Sikkim (*fide Hook. f., op. cit* ; Bor, *op. cit.*).

NEYRAUDIA Hook.f.

Neyraudia reynaudiana (Kunth) Keng *ex* Hitchc., Amer. Jour. Bot. 21 : 131. 1934 ; Bor, Grass. Ind. 518. 1960. *Arundo reynaudiana* Kunth, Rev. Gram. 1 : 275. t. 49. 1830. *Neyraudia madagascariensis* (Kunth) Hook. f. var. *zollingeri* Hook. f., F.B.I. 7 : 305. 1896. GUNGRING (Nep.).

A perennial grass up to 4 m tall ; culms terete, shining. Panicles large, nodding. Spikelets 4 - 8-flowered, lowest lemma empty.

Jorethang, Chatesay, Deorali ; 1000 - 1800 m.

Fl. & Fr. : July - Sept.

PHAENOSPERMUM Benth. & Hook. f.

Phaenospermum globosum Munro *ex* Oliver in Hook., Ic. Pl. Sub. t. 1991. 1891 ; Bor, Grass. Ind. 614. 1960.

A robust 2 - 3 m tall perennial. Leaves broad, acuminate, tessellately - nerved.

Sikkim (*fide* Bor, *op. cit.*).

PHALARIS L.

Phalaris arundinacea L., Sp. Pl. 55. 1753 ; Hook. f., F.B.I. 7 : 221. 1896 ; Bor, Grass. Ind. 615. 1960.

Perennial, rhizomatous grass. Culms tall, up to 1.5 m high. Panicles 7 - 40 cm long, dense. Glumes acute, wingless or with inconspicuous wing on the keel, sterile florets 2, below the fertile floret.

Sikkim (*fide* Bor, *op. cit.*).

P. minor Retz., Obs. Bot. 3 : 8. 1783 ; Hook. f., F.B.I. 7 : 221. 1896 ; Bor, Grass. Ind. 616. 1960.

Annuals ; culms 20 - 100 cm high. Panicles 1 - 6 cm long, ovate - oblong. Glumes equal, broadly winged, sterile floret one.

Betwo.

Fl. & Fr. : April - June.

PHLEUM L.

Phleum alpinum L., Sp Pl. 1 : 59. 1753 ; Hook. f., F.B.I. 7 : 236. 1896 ; Bor, Grass. Ind. 402. 1960.

Loosely tufted, shortly rhizomatous perennials ; culms erect or ascending. Panicles 1 - 5 cm long, broadly cylindrical to ovoid. Glumes truncate, stiffly ciliate on keel, awn equal in length to glumes, lemma shorter than glumes.

Zemu, Thangu ; 3300 - 3900 m.

Fl. & Fr. : July - Sept.

PHRAGMITES Adans.

Phragmites australis (Cav.) Trin. ex Steud., Nom. Bot. ed. 2. 324. 1854 ; Clayton in Taxon 17 : 168. 1968. *Arundo australis* Cav. in Anales Hist. Nat. Madrid. 1 : 100. 1799. *P. communis* Trin, Fund. Agrost. 134. 1820 ; Hook. f., F.B.I. 7 : 303. 1896 ; Bor, Grass. Ind. 416. 1960.

Perennial weed with creeping rhizomes. Panicles 20 - 30 cm long, 6 - 10 cm wide ; lowest branches with a silky beard at the base.

Chungthang ; ca 1600 m.

Fl. & Fr. : July - Oct.

PIPTATHERUM P. Beauv.

Piptatherum aeguiglume (Duthie ex Hook.f.) Duthie ex Rozhev. in Bot. Mater. Gart. Bot. Inst. Komarova. Acad. Nauk. USSR. 14 : 113. 1951. *Oryzopsis aeguiglumis* Duthie ex Hook. f., F.B.I. 7 : 234. 1896.

Loosely tufted perennial. Culms stout. Leaf-blades flat. Panicles lax. Spikelets lanceolate, lemma densely hairy, awn long, exserted, deciduous or subpersistent.

Giagong ; 2700 - 4500 m.

Fl. & Fr. : Oct. - Nov.

POA L.

Poa annua L., Sp. Pl. 68. 1753 ; Stapf in Hook. f., F.B.I. 7 : 345. 1896 ; Bor, Grass. Ind. 555. 1960.

Annual or subperennial ; culms erect, up to 30 cm tall, smooth. Panicle ovate, lax. Spikelets oblong ovate, 3 - 7-flowered, lemma silky ciliate along keel and outer-side nerves.

Sikkim, s. l. ; 1800 - 2500 m.

Fl. & Fr. : March - Sept.

P. eleanorae Bor in Kew Bull. 1948 ; 142. 1948 ; Bor, Grass. Ind. 556. 1960.

Perennial ; culms up to 40 cm tall, geniculate at base. Panicles up to 24 cm long. Spikelets cuneate, 2 - 3-flowered, glumes unequal, lemma 5 - 6 mm long, keels ciliate below, palea scabrous on keels.

N.E. Sikkim.

Fl. & Fr. : Aug.

P. gammieana Hook. f., F.B.I. 7 : 345. 1896 ; Bor, Grass. Ind. 557. 1960.

Perennial ; culms 30 - 60 cm high, leafy throughout. Panicles up to 10 cm long. Spikelets pale green, 3 - 2-flowered, lemma hairy on nerves and keel, palea ciliate on keels.

Changu, Naku Chu, Tankra Mt.

Fl. & Fr. : June - July.

P. himalayana Nees ex Steud., Syn. Pl. Glum. 1 : 256. 1854 ; Stapf in Hook. f., F.B.I. 7 : 344. 1896 ; Bor, Grass. Ind. 557. 1960.

Perennials. Culms ascending from geniculate base. Panicles lax, up to 15 cm long, branches capillary, flexuous. Spikelets 3 - 4-flowered, pale green, lemma subobtuse, 5-nerved, silky hairy on keel and side nerves, palea ciliate on keels.

Chola range, Lachung valley ; 3000 - 4000 m.

Fl. & Fr. : June - Aug.

P. hirtiglumis Stapf in Hook. f., F.B.I. 7 : 343. 1896. Nelderis in Hara *et al.*, Enum. Fl. Pl. Nepal 140. 1978.

Rhizomatous perennials. Culms 20 - 30 cm high, smooth. Panicles up to 5 cm long, pyramidal. Spikelets obovate, green or violet-tinged, glumes unequal, lemmas silky hairy at base, palea ciliate on the keels.

Lachung valley ; 4000 - 4900 m.

Fl. & Fr. : June - Aug.

P. khasiana Stapf in Hook. *f.*, F.B.I. 7 : 343. 1896 ; Bor, Grass. Ind. 557. 1960.

Annual ; culms 30 - 60 cm high. Panicles pyramidal. Spikelets ovate-oblong, pale green, glumes unequal, rough, lemmas oblong, keel and outer nerves sparsely silky or subglabrous, palea scabrid on keels.

Chamnango, Singalela range ; 3300 - 4200 m.

Fl. & Fr. : May - June.

P. ludens R.R. Stewart in Brittonia 5 : 420. 1945 ; Bor, Grass. Ind. 558. 1960. *P. pseudo-pratensis* Hook. *f.*, F.B.I. 7 : 340. 1896, non Beyer 1891 *nec* Serbn. et Ruyd. 1896.

Perennials ; culms smooth. Panicles up to 20 cm long, branches up to 7 cm long. Spikelets ovate-oblong, 6 - 8 mm long, 3 - 2-flowered, bluish, glumes unequal, lemma, oblong, silky hairy on keel beyond the middle, palea ciliate below.

Changu, Jelep - La, Zemu valley ; 3500 - 4000 m.

Fl. & Fr. : June - July.

P. nemoralis L., Sp. Pl. 69. 1753 ; Stapf in Hook. *f.*, F.B.I. 7 : 341. 1896 ; Bor, Grass. Ind. 558. 1960. *P. nemoralis* L. var. *ligulata* Stapf in Hook. *f.*, F.B.I. 7 : 341. 1896.

Perennial ; culms loosely tufted, 30 - 60 cm high. Panicles lax, branches erect or spreading. Spikelets 2 - 4-flowered, glumes unequal, rough, lemma oblong, subobtuse, with keel silky hairy at the base ; palea scaberulous on the keels.

Chamnago ; 3300 - 3600 m.

Fl. & Fr. : June - July.

P. nitide-spiculata Bor in Kew Bull. 1948 ; 139. 1948 ; Bor, Grass. Ind. 558. 1960.

Perennial ; culms up to 30 cm tall, glabrous. Panicles lax, up to 16 cm long. Spikelets elliptic-oblong, 2 - 3-flowered, long pedicellate, glumes unequal, lemma oblong, obtuse, ciliate on the keels at base, scabrous above, palea pilose between the keels.

Thangu, Jelep - la.

Fl. & Fr. : June - July.

P. pagophila Bor in Kew Bull. 1949 ; 239. 1949 ; Bor, Grass. Ind. 558. 1960. *P. flexuosa* Stapf in Hook. f., F.B.I. 7 : 342. 1896, non Sm. 1800.

Perennial, stoloniferous ; culms 20 - 40 cm high. Panicles pyramidal, lax. Spikelets green, usually suffused with purple, glumes equal, lemmas usually smooth, rarely scaberulous on the keels, palea scabrid on the keels.

Giagong, Yakla ; 3800 - 4600 m.

Fl. & Fr. : July - Sept.

P. poophagorum Bor in Kew Bull. 1948 : 143. 1948 ; Bor, Grass. Ind. 559. 1960.

Perennial ; culms 15 cm long, glabrous. Panicles up to 5 cm long. Spikelets on small scabrous branches, lemma glabrous, palea scabrous on keel.

Temu - la.

Fl. & Fr. : July.

P. polyneuron Bor in Kew Bull. 1952 ; 223. 1952 ; Bor, Grass. Ind. 559. 1960.

Perennials ; culms up to 30 cm tall. Panicles loose. Spikelets 3 - 4-flowered, compressed, elliptic ; glumes unequal ; lemma ciliate on nerves and keels, dorsally hairy in lower half, palea pilose in lower half.

Nathu - La.

Fl. & Fr. : June - July.

P. sikkimensis (Stapf) Bor in Kew Bull. 1952 ; 130. 1952 ; Bor, Grass. Ind. 560. 1960. *Poa annua* L. var. *sikkimensis* Stapf in F.B.I. 7 : 346. 1896.

Perennial, stoloniferous ; culms up to 30 cm high. Panicles up to 15 cm long, glumes firm, lemma obtuse, glabrous.

Patang La, Sebu valley ; 3500 - 4700 m.

Fl. & Fr. : July - Sept.

P. tibeticola Bor in Kew Bull. 1948 ; 139. 1948 ; Bor, Grass. Ind. 561. 1960.

Annual to perennial, culms up to 25 cm tall, glabrous. Panicles up to 10 cm long. Spikelets elliptic-oblong, 2 - 3-flowered, glumes unequal, lemma oblong-elliptic, obtuse, scabrous on keel and lateral nerves, palea scabrous on keel.

Chungyu ; ca 5000 m.

Fl. & Fr. : Sept.

POLYPOGON Desf.

Polypogon fugax Nees ex Steud., *Syn. Pl. Glum.* 1 : 184. 1854 ; Bor, *Grass. Ind.* 403. 1960. *P. higagaweri* Steud., *I.c.* 1 : 422. 1854. *P. littoralis* var. *higagaweri* (Steud.) Hook. *f.*, *F.B.I.* 7 : 246. 1896.

Annual. Panicles narrowly ovate, oblong or cylindrical, usually lobed. Spikelets pale green or yellowish, glumes slightly notched at the apex.

Sikkim, *s.l.* (*fide* Hook. *f.*, *op. cit.*).

There is only one collection without precise locality at CAL.

SPOROBOLUS R.Br.

Sporobolus diander (Retz.) P. Beauv., *Ess. Agrost.* 26. 147. 178. 1812 ; Hook. *f.*, *F.B.I.* 7 : 247. 1896 ; Bor, *Grass. Ind.* 629. 1960. *Agrostis diandra* Retz., *Obs. Bot.* 5 : 19. 1789.

Tufted perennials ; culms erect or geniculately ascending. Panicles effuse, stamens 2.

Sikkim, *s.l.*

S. fertilis (Steud.) W.D. Clayton in *Kew Bull.* 19 : 294. 1965. *Agrostis fertilis* Steud., *Syn. Pl. Glum.* 1 : 170. 1854. *Sporobolus indicus* auct. non R. Br., : Hook. *f.*, *F.B.I.* 7 : 247. 1896 ; Bor, *Grass. Ind.* 630. f. 76. 1960.

Perennials ; culms slender, erect. Panicles contracted. Spikelets pale green, upper glume shorter than lemma, stamens 3.

Phassing, Tassiding.

Fl. & Fr. : Apr. - Nov.

S. piliferus (Trin.) Kunth, *Enum. Pl.* 1 : 211. 1833 ; Hook. *f.*, *F.B.I.* 7 : 251. 1896 ; Bor, *Grass. Ind.* 632. 1960. *Vilfa pilifera* Trin., *Diss. Bot.* 157. 1824.

Densely tufted annuals. Panicles interrupted, linear, strict. Spikelets brown, upper glume as long as or longer than lemmas.

Sikkim ; wayside weed in tropical and subtropical parts.

Fl. : May - Sept.

STIPA L.

Stipa concinna Hook. *f.*, F.B.I. 7 : 230. 1896.

Tufted perennials, 8 - 22 cm high. Leaf-blades involute, setaceous. Panicles narrow, 2.5 - 4 cm long. Spikelets purplish ; lemma pilose at base only ; awn plumose throughout.

Sikkim (*fide* Hook. *f.* *op. cit.* ; Bor, *op. cit.*).

S. consanguinea Trin. & Rupr., Sp. Gram. Stirp. 78. 1842 & in Mem. Acad. Sci. Petersb. 6, 7 : 78. 1843 ; Bor, Grass. Ind. 664. 1960. *Stipa koelzii* R.R. Steward in Brittonia 5 : 441. 1945.

Tufted perennial, 17 - 25 cm high. Panicles contracted ; absent 2.5 cm long, lemma thinly pubescent to glabrous.

Sikkim, *s.l.*, G.H. Cave 276 (CAL).

S. mongholica Turcz. ex. Trin. in Bull. Sci. Acad. Petersb 1 : 67. 1836 ; Hook. *f.*, F.B.I. 7 : 229. 1896 ; Bor, Grass. Ind. 645. 1960.

Tufted perennials. Leaf-blades involute, setaceous. Panicles loose. Spikelets 3 - 5 mm long ; pedicels capillary ; glumes subequal, lemma pilose, awn flexuous, plumose throughout.

Chola, Lhonak ; 4000 - 4500 m.

Fl. & *Fr.* : July - Aug.

S. purpurea Griseb. in Nachr. Ges. Wiss., Gottingen 1868 : 82. 1868 ; Hook. *f.*, F.B.I. 7 : 229. 1896 ; Bor, Grass. Ind. 645. 1960.

Tufted perennials, 15 - 25 cm high. Panicles loose, 6 - 12 cm long ; glumes shortly acuminate, lemma pubescent, awn 6 - 8 cm long, bigeniculate, plumose throughout.

Naku - la ; 4800 - 5800 m.

Fl. & *Fr.* : Oct. - Nov.

***S. roylei* (Nees)** Mez. in Fedde Repert. 17 : 207. 1921 ; Bor, Grass. Ind. 646. 1960. *Orthoraphium roylei* Nees in Proc. Linn. Soc. 1 : 94. 1841. *S. orthoraphium* Steud., Syn. Pl. Glum. 131. 1854 ; Hook. *f.*, F.B.I. 7 : 233. 1896.

Densely tufted perennials. Culms slender. Panicles 15 - 25 cm long. Spikelets on short, erect pedicels ; glumes unequal, lemma subulate, awn up to 2.5 cm long, not twisted, glabrous.

Chakung - chu, Gnatong, Jelep la ; 3000 - 3700 m.

Fl. & Fr. : July - Oct.

THYSANOLAENA Nees

Thysanolaena latifolia (Roxb. ex Hornem.) Honda in J. Fac. Sc. Tokyo Sect. Ill. Bot. 3 : 312. 1930 ; Mebberley in Taxon 33 (3) : 437. 1984. *Agrostis maxima* Roxb., Fl. Ind. 1: 319. 1820. *Thysanolaena maxima* (Roxb.) O. Ktze., Rev. Gen. Pl. 2 : 794. 1891. *Thysanolaena agrostis* Nees in Edinb. New. Phil. J. 18 : 180. 1835 ; Hook. f., F.B.I. 7 : 61. 1896. *T. sikkimensis* Gdger. in Bull. Soc. Bot. Fr. 66 : 303. 1919.

Perennials. Culms 1.5 - 3.5 high. Leaves lanceolate. Panicles 30 - 60 cm long. Spikelets *ca* 1.5 mm long, lower florets sterile, upper bisexual.

Rangeet, Rebong, Dikchu, Gangtok ; 500 - 1700 m.

Fl. & Fr. : Feb. - May.

TRIKERAIA Bor

Trikeria hookeri (Stapf) Bor in Kew Bull. 1954 ; 555. 1955 ; Bor, Grass. Ind. 647. 1960. *Stipa hookeri* Stapf in J. Linn. Soc. Bot. 30 : 120. 1894 ; Hook. f., F.B.I 7 : 232. 1896.

Culms up to 70 cm high. Panicles 10 - 15 cm long. Spikelets 1-flowered, glumes membranous, lemma oblong, elliptic, hyaline, hairy, deeply bifid at the apex, lateral nerves produced to two stout scabrid bristles, awn from back of the lemma.

Kankola Pass (*fide* Hook. f., *op. cit.*).

TRIPOGON Roem. & Schult.

Tripogon filiformis Nees ex Steud., Syn. Pl. Glum. 1 : 301. 1854 ; Hook. f., F.B.I. 7 : 288. 1896 ; Bor, Grass. Ind. 521. 1960.

Tufted perennials ; culms simple. Spikelets 5 - 8-flowered ; lemma with a median awn, shortly awned, outer lobes short or long, obtuse or acute or awned inner lobes.

Lachung ; Reunig Chu.

Fl. & Fr. : Sept. - Oct.

T. trifidus Munro ex Stapf in Kew Bull. 1892 ; 85. 1892 ; Hook. f., F.B.I. 7 : 286. 1896 ; Bor, Grass. Ind. 524. 1960.

Tufted perennials, up to 45 cm high, lower glume lobed on one side, lemma bearded at the base.

Sikkim (*fide* Hook. f. *op. cit.* ; Bor, *op. cit.*).

TRISETUM Pers.

Trisetum flavescens (L.) P. Beauv., Ess. Agrost. 88. 153. 180. 1812 ; Bor, Grass. Ind. 448. 1960. *Avena flavesceps* L., Sp. Pl. 80. 1753. *A. sikkimensis* Hook. f., F.B.I. 7 : 280. 1896.

Perennials. Panicles very effuse. Spikelets yellowish green, glumes unequal, hyaline.

Lachen & Lachung valley (*fide* Hook. f., *op. cit.*).

T. scitulum Bor in Kew Bull. 1956 ; 212. 1956 ; Bor, Grass. Ind. 448. 1960. *Avena flavescens* auct. non L. : Hook. f., F.B.I. 7 : 279. 1896.

Perennials. Panicles effuse. Spikelets brownish or bronze, glumes unequal, lemma shining.

Yumthang, Zemu valley, Sherabthang.

Fl. & Fr. : Aug. - Sept.

T. spicatum (L.) Richt., Pl. Eur. 1 : 59. 1890 ; Bor, Grass. Ind. 448. 1960. *Aira spicata* L., Sp. Pl. 64. 1753. *Avena subspicata* L., Syst. Nat. ed. 10 : 873. 1759. Hook. f., F.B.I. 7 : 278. 1896. *Trisetum subspicatum* (L.) P. Beauv., Ess. Agrost. 88. 149. 1812.

Perennials. Panicles spiciform or ovoid, erect, densely spiculate.

Llonak, Zemu valley, Dongri ; 3600 - 4800 m.

Fl. & Fr. : June - Aug.

TRITICUM L.

Triticum aestivum L., Sp. Pl. 85. 1753 ; Bor, Grass. Ind. 679. 1960. *T. vulgare* Vill., Hist. Pl. Dauph. 2 : 153. 1787 ; Hook. f., F.B.I. 7 : 367. 1896.

Cultivated.

INDEX TO BOTANICAL NAMES

(R.C. Srivastava)

- Acampe** Lindl. 24
 - longifolia* (Lindl.) Lindl. 25
 - ochracea* (Lindl.) Hochr. 24
 - papillosa* (Lindl.) Lindl. 24
 - rigida* (Buch. - Ham. ex J.E. Smith) P.F. Hunt 25
- Acanthophippium** Bl. 25
 - striatum* Lindl. 25
 - sylhetense* Lindl. 25
- Aceras** *angustifolia* Lindl. 80
- Acianthus** *petiolatus* D. Don 85
- Aclisia** *subumbellata* (Clarke) Brueck. 171
- ACORACEAE** 181
- Acorus** L. 185
 - calamus* L. 185
 - gramineus* 185
- Acriopsis** Reinw. ex. Bl. 25
 - harai* Tuyama 25
 - javanica* Reinw. ex Bl. 25
- Achrochaene** Lindl. 26
 - punctata* Lindl. 26
- Aegilops** *exaltata* L. 260
- Aerides** Lour. 26
 - ampullaceum* Roxb. 30
 - appendiculatum* Lindl. 46
 - calceolare* J.E. Smith
 - cylindricum* auct.
 - dasytopogon* J.E. Smith 74
 - decumbens* Griff. 82
 - diformis* Wall. ex Lindl. 96
 - longicornu* Hook. f. 98
 - multiflora* Roxb. 26
 - odorata* Lour. 26
 - rigida* Buch. - Ham. ex J.E. Smith 25
 - suaveolens* Roxb. 108
 - taeniale* Lindl. 82
 - testaceum* Lindl. 118
 - uniflora* (Lindl.) Summerh. 98
 - vandarm* Reichb. f. 99
- Aetheria** *fusca* Lindl. 76
- AGAVACEAE** 139
- Agave** L. 139
 - americana* sensu Hook. f.
 - cantula* Roxb.
 - foetida* L. 140
 - sisalana* Perrine 139
 - vera-cruz* Mill. 140
- Agropyron** *semicostatus* (Nees ex Steud.) Boiss. 288
 - sikkimensis* Melderis 288
- Agrostis** *alba* auct. 277, 279
 - clarkei* Hook. f. 278
 - debilis* (Hook. f.) Bor 286
 - diandra* Retz. 303
 - divaricata* Griseb. 278
 - fertilis* Steud. 303
 - filipes* Hook. f. 276
 - gigantea* Roth 277
 - himalayana* Bor 277
 - hookeriana* Clarke ex Hook. f. 277
 - inaequiglumis* Griseb. 277
 - maxima* Roxb. 305
 - micrantha* Steud. 277
 - myriantha* Steud. 278
 - var. *sikkimentis* Hook. f. 278
 - neodebilis* Bennet & Raiz. 286
 - nervosa* Nees ex Trin. 278
 - pilosula* Trin. 278
 - poluninii* Bor 277
 - sikkimensis* Bor 278
 - stolonifera* L. 279
 - ssp. *gigantea* (Roth) Naie & Weiller 277
 - triaristata* (Hook. f.) Bor 279
- Agrostophyllum** Bl. 26
 - brevipes* King & Pantl. 26
 - callosum* Reichb. f. 26
 - khasianum* Griff. 27
 - myrianthum* King & Pantl. 27
- Aira** *caespitosa* L. 286
 - indica* L. 271
 - Aletris* L. 142
 - gracilis* Rendle 142
 - nepalensis* Hook. f. 143
 - pauciflora* (Klotzsch.) Hand. - Mazz. 142
 - sikkimensis* Hook. f. 143
 - spicata* L. 306
- Allium** L. 143
 - cepa* L. 143
 - ellipticum* Wall.
 - fasciculatum* Rendle. 143
 - gageanum* W.W. Smith apud W.W. Smith & Cave 143
 - hookeri* Thw. 143
 - jacquemontii* Regel 144
 - macranthum* Baker 144
 - pratii* var. *ellipticum* Wang & Tang. 144
 - pratii* C.H. Wright apud Forbes & Heinsley 144
 - prezwalskium* Regel 144

- sativum** L. 144
sikkimense Baker 144
victorialis auct. 144
 var. *angustifolium* Hook. f. 144
wallachii Kunth. 145
- Allotropopsis** C.B. Presl. 243
semialata (R. Br.) Hitch. 243
- Alocasia** (Schott.) D. Donn 185
fallax Schott 185
macrorrhizus L. 186
navicularis (Kotch. & Bouche) Koch. &
 Bouche 186
odora Roxb. 186
- Aloe** L. 145
vera (Lindl.) Burm. f. 145
- Alopecurus aequalis** Sobol. 279
aristatus Michx. 279
geniculatus L. 279
- Alpinia** Roxb. 120
allughas (Retz.) Rosc. 121
bracteata Roxb. 120
linguiforme Roxb.
malaccensis (Burm. f.) Rosc. 121
nigra (Gaertn.) Burtn 121
- AMARYLLIDACEAE** 137
Amia latifolia Lindl. 114
- Amischophacelus** Rolla Rao &
 Kammathy 167
axillaris (L.) Rolla Rao & Kammathy
 167
- Amischotolype** Hassk. 167
hookeri (Hassk.) Hara 167
mollissima (Bl.) Hassk. 168
- Amitostigma** Schltr. 27
puberulum (King & Pantl.) Schltr. 27
- Amomum** Roxb. 121
aromaticum Roxb. 121
corynostachyum Wall. 122
dealbatum Roxb. 122
 var. **dealbatum** 122
 var. **sericeum** (Roxb.) Baker 122
- kingii** Baker 122
linguiforme (Roxb.) Baker 131
sericeum Roxb. 122
subulatum Roxb. 122
zedoaria Christm. 126
zerumbent. L. 134
- Amorphophallus** Bl. ex Decne 186
bulbifer (Schott.) Bl. 186
napalensis Bogner & Mayo 186
- Androcorys** Schltr.
gracilis (King & Pantl.) Schltr.
- Andropogon** L.
aciculatus Retz 247
annulatum Forssk 251
assimilis Steud. 247
binatus Retz. 256
bladhii Retz. 245
bootanensis Hook. f. 272
brevifolium Swartz 272
caricosus L. 251
castratus Griff. 244
crinitus Thunb. 267
delavayi Hook 272
dulce Burm. f. 218
fascicularis Roxb. 269
flexuosus Nees ex Steud. 248
foveolatus Del. 254
gryllus L. 247
intermedius R. Br. 245
ischaemum L. 246
lancearius Hook. 248
lancifolius Trin. 245
leschenaultiamus Decne 255
martinii Roxb. 249
micranthus Kunth 247
microphyllus Trin. 245
microtheca Hook. f. 249
nardus L. 248
 var. **flexuosus** Hack 248
 var. **grandis** Hook. 249
 var. **khasianus** Hook. 249
 var. **sikkimensis** Bor 248
nitidulus Hook. f. 269
odoratus Donna Lisboa 245
pendulus Nees ex Steud. 249
quadrivalvis L. 275
schoenanthus L. 249
 var. **martinii** (Roxb.) Hook. f. 249
sorghum 274
ssp. *sativus* 274
 var. **miliiformis** Hook. 274
 var. **sikkimensis** Snowden. 275
squarrosum L. f. 276
vimineus Trin. 259
- Aneilema compressa** Dalz. 171
herbaceum 170
 var. **divergens** Clarke 170
lineolatum Kunth 170
malabaricum (L.) Brueck. 171
nudiflorum 171
 var. **compressa** (Dalz) Clarke 171
scaberrimum (Bl.) Kunth. 172
scapiflorum (Roxb.) Kostel. 170
vaginatum (L.) R. Br. 171
indica (Lindl.) R. Br.

- Anguillaria indica* (Lindl.) R. Br. 150
Aniselytron Merr. 279
treutleri (O. Ktze.) Soják. 279
Anisopetalon careyanum Hook. 32
Anoectochilus Bl. 27
crispus Lindl. 27
elwesii (Clarke ex Hook. f.) King & Pantl. 27
grandiflorus Lindl. 28
lanceolatus Lindl. 28
luteus Lindl.
roxburghii (Wall.) Lindl. 28
sikkimensis King & Pantl. 28
Anthistiria arundinacea Roxb. 275
ciliata L. f. 275
gigantea 275
 ssp. *arundinacea* (Roxb.) Hook. 275
hookeri Griseb. 275
imberbis Retz. 276
Anthogonium Wall. ex Lindl. 29
gracile Wall. ex Lindl. 29
Anthoxanthum L. 280
flexuosa (Hook. f.) Veldk. 280
hookeri (Griseb.) Rendle 280
sikkimense (Maxim.) Ohwi 280
Aorchis Vermeulen
spathulata (Lindl.) Vermeulen
Apaturia lindleyana Wt. 97
montana Thw. 29
senile Lindl. 97
Aphyllorchis Bl. 29
alpina King & Pantl. 29
montana (Thw.) Reichb. f. 29
pantlingii W.W. Smith 29
parviflora King & Pantl. 29
prainii Hook. f. 29
Apluda L. 243
digitata L. f. 268
mutica L. 243
varia Hack 243
Apocopsis Nees 244
palaeacea Trin 244
royleanus Nees 244
Apostasia Bl. 29
odorata 29
wallichii R. Br. 29
Appendicula Bl. 30
bifaria Lindl. 30
cornuta Bl. 30
teres Griff. 44
ARACEAE 185
Arachnanthe 30
cathcartii (Lindl.) Benth. 30
clarkei (Reichb. f.) Rolfe 70
Arachnis Bl. 30
cathcartii (Lindl.) J.J. Smith 30
clarkei Rolf
lobrosa (Lindl. ex Paxt.) Reichb. f.
Areca L. 180
catechu L. 180
gracilis Roxb. 183
ARECACEAE 180
Ariopsis Niimmo 187
peltata Niimmo 187
Arisaema Mart. 187
biflagellatum Hara 189
concinnum Schott. 187
consanguineum Schott. 187
costatum (Wall.) Mart. ex Schott 187
 var. *sikkimensis* (Stapf.) Hara 190
echinatum (Wall.) Schott 188
erubescens (Wall.) Schott 188
exile Schott 189
flavum (Forssk.) Schott. 188
galeatum N.E. Br. 188
griffithii Schott. 188, 191
 var. *griffithii* 188
 var. *verrucosum* (Schott) Hara 188
hookorianum schott 188
intermedium Bl. 189
 var. *intermedium* 189
 var. *biflagellatum* (Hara) Hara 189
jacquemontii Bl. 189
mirabile Schott 190
nepenthoides (Wall.) Mart. 189
ochraceum Schott 190
ostiolatum Hara 190
propinquum Schott 190
sikkimensis Stapf. ex Chatterjee 190
speciosum Wall. 190
tortuosum (Wall.) Schott. 191
utile Hook. f. ex Schott. 191
verrucosum Schott 189
wallichianum 190
 var. *sikkimensis* (Stapf. ex Chatterjee)
 Hara 190
Annodorum labrosum (Lindl. ex Paxt.)
 Schltr.
Arrhynchium labrosum Lindl. ex Paxt.
Arthraxon P. Beauv. 244
breviaristatus Hack. 244
castratus (Griff.) Narayanswami ex Bor.
 244
ciliaris B. Beauv. 244
hispidus (Thunb.) Makino 244
hookeri (Hack.) Henr.

- inermis* Hook. f. 244
lancifolius (Trin.) Hochst. 245
microphyllus sensu Hook. f. 245
microphylus (Trin.) Hochst.
nudus Steud. 244
quartinianus (A. Rich.) Nash. 244
rudis (Steud.) Hochst. 244
sikkimensis Bor 245
submuticus (Steud.) Hochst. 244
Arum L. 186
bulbiferum Roxb. *apud* Sims. 186
Colocasia L. 191
castatum Wall. 187
echinatum Wall. 188
erubescens Wall. 188
esculentum L. 191
flavum Forssk. 188
macrorhizon L. 186
nephenthoides Wall. 189
odorum Roxb. 186
tortuosum Wall. 191
trilobatum L. 195
viviparum Roxb.
Arundina Bl. 30
bambusifolia Lindl. 30
chinensis Bl. 30
graminifolia (D. Don) Hochr. 30
Arundinaria
aristata Gamble 242
falconeri Hook. f. 240
gracilis Blanch. 239
hookeriana Munro 237
intermedia Munro 239
maling Gamble 242
microphylla (Griff.) Munro 242
pantlingii Gamble 242
polystachya Kurz *ex* Gamble 240
racemosa Munro 243
suberecta Munro 240
Arundinella Raddi 280
bengalensis (Spreng.) Druce 280
brasiliensis auct. 281
clarkei Hook. f. 281
decempedalis (O. Ktze.) Jan. 281
hookeri Munro *ex* Keng. 281
nepalensis Trin. 281
villosa Steud. 281
 var. *himalaica* Hook. f. 281
wallichii Nees *ex* Steud. 280
Arundo L. 281
australis Cav. 299
donax L. 281
pseudo-phragmites Hall. f. 283
 reynaudiana Kunth 298
Ascocentrum Schltr. 30
 ampullaceum (Roxb.) Schltr. 30
 micranthum (Lindl.) Holttum 111
Asparagus L. 145
 filicinus Buch. - Ham. *apud*
 D. Don 145
 racemosus Willd. 145
 var. *racemosus* 145
 var. *subacerosus* Baker 145
Asphodelus L. 146
 tenuifolius Cav. 146
Ataxia *hookeri* Griseb. 280
Aulacolepis *treatleri* (O. Ktze.) Hack. 280
Avena L. 282
 aspera Munro *ex* Thw. 295
 fatua L. 282
 ssp. *fatua* 282
 var. *glabrata* Peterm. 282
 flavescens L. 306
 flavescens auct. 306
 sativa L. 282
 var. *sericea* Hook. f. 282
 sativa L. 282
 sikkimensis Hook. f. 306
 subspicata L. 306
Axonopus *semialatus* (R. Br.) Hook. f. 243
Bambos arundinacea Retz. 237
Bambusa Schreb. 237
 arundinacea (Retz.) Willd. 237
 longispiculata Gamble 237
 microphylla Griff. 242
 nutans Wall. *ex* Munro 237
 pallida Munro 237
 teres Buch. - Ham. *ex* Munro 237
 tulda Roxb. 237
BAMBUOIDEAE 237
Bonksia speciosa Koen. 124
Belamcanda Adans 136
 chinensis (Lindl.) DC. 136
Berghansia tenella Arn. *ex* Miq. 295
Biermannia King & Pantl. 31
 bimaculata (King & Pantl.) King &
 Pantl. 31
Bletia *bicallosa* D. Don 70
 graminifolia D. Don 30
 mascua D. Don 43
 woodfordii Hook. 102
Blysmus Panz. *ex* J. A. Schultes 198
 compressus (L.) Panz. *ex* Link. 198
 var. *sikkimensis* (Clarke)
 Karth. 198
Bothriochloa O. Ktze. 245

- bladhii** (Retz.) S.T. Blake 245
glabra (Roxb.) A. Camus 245
intermedia (R. Br.) A. Camus 245
ischaemum (L.) Keng 246
odorata (Donna Lisboa) A. Camus 245
Brachiaria Griseb. 246
 ramosa (L.) Stapf. 246
 reptans (L.) Gard. et Hubb. 246
 villosa (Lam.) A. Camus 246
Brachycorythis Lindl. 31
 obcordata (Buch. - Ham. ex D. Don)
 Summerh. 31
Brachypodium P. Beauv. 282
 sylvaticum (Huds.) P. Beauv. 282
 wattii Clarke 282
Briza L. 282
 media L. 282
Bromus L. 282
 grandis (Stapf.) Melderis 283
 himalaicus Stapf. 282
 var. **grandis** Stapf. 283
 porphyranthos T.A. Cope 283
Bulbodium serotinum L.
Bulbophyllum Du. Petit Thouars. 31
 affine Lindl. 31
 alopécurum Reichb. 40
 andersonii (Hook. f.) J.J. Smith 31
 bicolor (Lindl.) Hook. f. 112
 bisetum Lindl. 32
 bootanense (Griff.) Par. & Reichb. f. 32
 brevipes Ridl. 40
 careyanum Lindl. 32
 caudatum Lindl. 32
 cauliflorum Hook. f. 33
 clarkeanum King & Pantl. 33
 confertum Hook. f. 39
 cornu-cervi King & Pantl. 33
 cylindraceum Lindl. 33
 ebulbum King & Pantl. 33
 elatum (Hook. f.) J.J. Smith 34
 eublepharum Reichb. f. 33
 flavidum 36
 gamblei (Hook. f.) Hook. f. 34
 gracilipes King & Pantl. 41
 griffithii (Lindl.) Reichb. f. 34
 guttulatum Wall. ex Hook. f. 34
 gymnopus Hook. f. 34
 helenae (O. Ktze.) J.J. Smith 35
 hirtum (J.E. Smith) Lindl. 35
 hymenanthum Hook. f. 35
 leopardianum (Wall.) Lindl. 35
 leptanthum Hook. f. 34, 35
 var. **gamblei** Hook. f. 34
 listeri King & Pantl. 36
 micranthum Hook. f. 40
 odoratissimum (J.E. Smith) Lindl. 36
 ornatissimum (Reichb. f.) J.J. Smith 31
 paleaceum (Lindl.) Hook. f. 112
 pantlingii Lucksom 36
 parvulum Hook. f. 36
 piluliferum King & Pantl. 37
 polyrhizum Lindl. 37
 protractum Hook. f. 37
 reptans (Lindl.) Lindl. 37
 var. **reptans** 37
 var. **sub-racemosa** Hook. f. 37
 retusiusculum Reichb. f. 38
 rigidum King & Pantl. 38
 rolfei (O. Ktze.) Seid. 38
 sarcophyllum (King & Pantl.)
 J.J. Smith 38
 scabratum Reichb. f. 38
 secundum Hook. f. 39
 sikkimensis (King & Pantl.)
 J.J. Smith 39
 striatum (Griff.) Reichb. f. 39
 thomsonii Hook. f. 39
 triste Reichb. f. 40
 umbellatum auct 34
 umbellatum Lindl.
 viridiflorum (Hook. f.) Schltr. 40
 wallichii (Reichb. f.) 40
 xylophyllum Par. & Reichb. 41
 yoksunense J.E. Smith 40
Bulbostylis Kunth 198
 barbata (Rottb.) Clarke 198
 catrillaris Kunth. 198
 var. **trifida** (Nees) Clarke 198
 densa (Wall.) Hand. - Mazz. 198
 trifida (Nees) Nelmes 198
BURMANNIACEAE 23
Burmannia L. 23
 coelestis D. Don 23
 disticha L. 23
 nepalensis (Miers) Hook. f. 24
BUTOMACEAE
Butomopsis Kunth
 lanceolatus (Roxb.) Kunth
 latifolia (D. Don) Kunth
Butomus *lanceolatus* Roxb.
 latifolius D. Don
 Caladium pumilum D. Don
Calamagrostis Adans 283
 debilis Hook. f. 286
 modensis Griseb. 283
 filiformis Girseb. 287

- lahulensis* G. Singh 287
littorea (Schrad.) P. Beauv. 284
pilosula (Trin) Hook. f. 278
pseudophragmites (Hall. f.) Koeler 284
pulchella Griseb. 287
scabrescens Griseb. 287
tibetica (Bor.) G. Singh 287
tripilifera Hook. f. 279
Calamus L. 181
acanthospathus Griff. 181
erectus Roxb. 181
flagellum Griff. 181
jenkineanus Griff. 182
latifolius Roxb. 181
leptospadix Griff. 181
Calanthe R. Br. 41
alismaeifolia Lindl. 41
alpina Hook. f. ex Lindl. 41
angusta Lindl. 41
biloba Lindl. 41
brevicornu Lindl. 42
chloroleuca Lindl. 42
densiflora Lindl. 42
fulgens Lindl. 43
gracilis Lindl. 42, 44
herbacea Lindl. 42
keshabii Lucksom
mannii Hook. f.
mascua (D. Don) Lindl. 43
 var. *mascua* 43
 var. *fulgens* (Lindl.) Hook. f. 43
puberula Lindl. 43
tricarinata Lindl. 43
treutleri Hook. f. 43
trulliformis King & Pantl. 43
uncata Lindl. 43
whiteana King & Pantl. 44
obtusa Lindl. 91
Compelia mollissima Bl. 168
Campylandra Baker 146
 aurantica Baker 146
CANNACEAE 134
Canna L. 134
 edulis Ker - Gawler 134
 indica L. 135
 var. *orientalis* (Rosc.) Baker 135
 orientalis Rosc. 135
Capillipedium Stapf. 247
 assimile (Steud.) A. Camus 247
 parviflorum (R. Br.) Stapf. 247
Cardiocrinum Lindl. 146
 giganteum (Wall.) Makino 146
Carex L. 199
 alpecuroides D. Don ex Tiloch & Taylor 199
 alpina Sw. 199
 var. *alpina* 199
 var. *gracilenta* (Bott.) Clarke 203
 var. *Ichmannii* (Drej.) R.C. Srivastava
aristata D. Don ex Tiloch & Taylor 211
atrata L. 199
 var. *atrata* 199
 var. *glacialis* Boott 199
 var. *pullata* Boott 200
 ssp. *pullata* (Boott) Kükenth.
atrofusca Schkuhr. 200
 ssp. *atrofusca* 200
 ssp. *angustifructus* 200
 ssp. *minor* 200
baccans Nees 200
bigelowii Torr. ex Schweinitz. 200
breviculmis sensu Clarke 211
capillacea Boott 201
condensata Nees 201
continua Clarke 201
cruciata Wahdenb. 201
cruenta Nees 201
curvata Boott. 223
daltonii Boott 202
decora Boott 202
duthiei Clarke 200
 var. *glacialis* (Boott) Clarke 199
emineus Nees 206
esenbeckii Kunth 223
ferruginea sensu Clarke 201
filicina 202
 var. *meogyna* (Nees) Strachey 202
 var. *minor* Boott 202
finitima Boott 202
 var. *attenuata* Clarke 202
foliosa D. Don ex Tiloch & Taylor 201, 213
fragilis Boott 203
fucata Boott ex Clarke 203
 var. *sikkimensis* (Clarke) Kükenth. 211
fusiformis Nees 203
gracilenta Boott apud Strachey 203
haematostoma Nees 203
inaequalis Boott ex Clarke 204
inanis Kunth 204
inclinis Boott ex Clarke 204
indica L. 204
insignis Boott 205
japonica Thunb. 199
 var. *alopecuroides* (D. Don ex Tiloch & Taylor) Clarke 199

- kingiana* Clarke 211
laeta Boott 205
lehmanni Drejer 205
lenticularis D. Don
linearis Boott
var. *elachista* Clarke
longipes D. Don ex Tilloch & Taylor 205
var. *nepalenisis* (Spr.) Boott 205
var. *racemosa* Kükenth.
jurida Clarke 208
microglochin Wahlenb. 205
monopleura Krecz 206
montis - everestii Kükenth
moorcroftii Falc. ex Boott 206
munda Boott. 206
muricata L.
var. *muricata* 213
var. *foliosa* (D. Don ex Tilloch & Taylor) Clarke 213
myosurus Nees 206
ssp. *spiculata* (Boott) Kükenth. 212
var. *nobilis* Kükenth. 212
var. *eminens* (Nees) Clarke 206
var. *ratongensis* Clarke 207
nakaoana T. Koyama 203
nepalensis Spr. 205
nilagirica Steud. 202
nivalis Boott 207
nobilis Boott 212
notha Kunth 207
nubigena D. Don ex Tilloch & Taylor 207
var. *fallax* Clarke
obscura Nees 207
var. *brachycarpa* Clarke 208
obscuriaps Kükenth. Kullesh 208
oedorhampha Nelmes 208
olivacea Boott 208
orbicularis Boott 208
pamirensis Clarke ex Fedtsch. 209
parva Nees 209
phacota Spr. 209
polyccephala Boott 209
praeclara Helma 209
praelonga Clarke 209
pseudofoetida Kükenth. 210
psychrophila Nees 210
pulchra Boott 210
radicalis Boott 210
rara Boott 210
var. *rara*
var. *capillaris* (Boott) R.C. Srivastava
remota 206
ssp. *rochebrunii* (Franch. & Sav.)
Kükenth 206
rigida Gooden 200
rigida *sensu* Clarke 208
rochebrunii Franch. & Sav. 206
rostrata *sensu* Clarke 209
royleana Nees 211
Carex sahnii Ghildyal & Bhattacharyya 211
schlagintweitiana Boeck. 204
setigera D. Don 211
var. *schlagintweitiana* (Boeck.)
Kükenth. 204
setosa Boott. 211
sikkimensis Clarke 211
spiculata Boott 212
var. *nobilis* Clarke 212
stramentita Boott ex Boeck. 212
thomsonii Boott 212
trinervis Nees
tumida Boott 208
uncinoides Boott 226
ustulata *sensu* Clarke 200
var. *minor* Boott 200
vesiculos Boott. 212
var. *vesiculos* 212
var. *congesta* 213
forma *pallida* Kükenth. 213
var. *pallida* (Kükenth.) R.C. Srivast. 213
vidua Boott 227
vulgaris *sensu* Clarke 208
wallichiana Spr. 213
Caryota L. 182
urens L. 182
Catabrosa P. Beauv. 284
sikkimensis Stapf. 284
wallichii Hook. f. 285
Caulokaempferia K. Larsen 123
secunda (Wall.) K. Larsen 123
sikkimensis (King ex Baker) Larsen 123
Cautleya (Benth.) Royle ex Hook. f. 123
cathcartii Baker 123
gracilis (J.E. Smith) Dandy 123
lutea (Royle) Hook. f. 123
robusta Baker 124
spicata (J.E. Smith) Baker 124
Cenchrus granularis L. 260
lappaceus L. 284
Centotheca Desr. 284
lappacea (L.) Desv. 284
Cephalanthera L.C. Rich. 44
ensifolia L.C. Rich. 44

- longifolia** Fritsch. 44
Cephalantheropsis Guill. 44
gracilis (Lindl.) Hu 44
Cephalostachyum capitatum Munro 241
Iotifolium Munro 241
Ceratostylis Bl. 44
himalaica Hook. f. 109
subulata Bl. 44
teres (Griff.) Reichb. f. 44
Chamaeraphis spinescens Poir 269
 var. *brunonianana* (Griff.) Hook. f. 269
Chamaerops martiana Wall. 184
Cheirostylis Bl. 45
griffithii Lindl. 45
Chilocista Lindl. 45
lunifera (Reichb. f.) J.J. Smith ex
 Holttum 45
Chimonobambusa Makino 237
hookeriana (Munro) Nakai 237
intermedia (Munro) Nakai 239
jaini Das & Pal 239
polystachya (Kurz ex Gamble) Nakai 240
Chlorophytum Ker. - Gawler 146
arundinaceum Baker 146
breviscapum auct. 146, 147
comosum Thunb. 147
khasianum Hook. f. 147
nepalense (Lindl.) Baker 147
undulatum Wall ex Hook. f. 147
Chrysobaphus roxburghii Wall. 28
Chrysoglossum Bl. 45
erraticum Hook. 45
macrophyllum King & Pantl. 63
Chrysopogon Trin. 247
aciculatus (Retz.) Trin 247
gryllus (L.) Trin 247
lancearius (Hook. f.) Haines 248
Chusua whomship Maxim. 108
Cirrhopteridium andersonii Hook. f. 31
bootanense Griff. 32
brevipes Hook. f. 40
caespitosum Wall. ex Lindl. 38
cornutum Lindl. 35
elatum Hook. f. 34
guttulatum Hook. f. 34
maculosum Lindl. 40
ornatissimum Reichb. f. 36
parvulum Hook. f. 38
refractum sensu Hook. f. 40
sarcophyllum King & Pantl. 38
sikkimensis King & Pantl. 39
viridiflorum 40
wallichii Lindl. 38
- Cleiscocentron** Brühl 45
trichromum (Reichb. f.) Brühl 45
Cleisostoma Bl. 46
appendiculatum (Lindl.) Benth. & Hook.
 f. ex Jackson 46
armigera King & Pantl. 107
aspersum (Reichb. f.) Garay 46
bicuspidatum Hook. f.
brevipes Hook. f. 47
filiforme (Lindl.) Garay 46
gemmata (Lindl.) King & Pantl. 111
micrantha (Lindl.) King & Pantl. 111
racemiferum (Lindl.) Garay 46
spathulatum Bl. 109
spicatum Lindl. 109
subulatum Bl. 47
striatum (Reichb. f.) 47
wendlandorum Reichb. f. 107
Clintonia Raffin 147
alpina Kunth ex Baker 147
undensis Trautv. & Meyer 147
 var. *alpina* (Kunth ex Baker) Hara
 147
Cnemidia angulosa Lindl. 113
Coccus L. 182
nucifera L. 182
Codogyne coronaria Lindl. 67
Coelachne R. Br. 284
simpliciuscula (Steud.) Benth. 284
pulchella sersx 284
Coelogyne Lindl. 47
coronaria Lindl.
corymbosa Lindl. 47
cristata Lindl. 47
elata Lindl. 50
fimbriata Lindl. 47
flaccida Lindl. 48
flavida Wall. ex Lindl. 49
fuscescens Lindl. 48
hookeriana Lindl. 105
humilis (J.E. Smith) Lindl. 106
longipes auct. 49
longipes Lindl. 48
maculata Lindl. 106
nitida (Wall. ex D. Don) Lindl. 48
occultata Lindl. 48
ovalis Lindl. 49
parviflora Lindl. 97
praecox (J.E. Smith) Lindl. 106
prolifera Lindl. 49
punctulata Lindl. 49
raizadae Jain & Das 49
schultesii Jain & Das 49

- stricta** Schltr. 50
treutleri Hook. f. 50
uniflora Lindl. 50
Coelorachis stricta (Nees ex Steud.) A. Camus. 260
Coix L. 248
lacryma-jobi 248
 var. *lacryma-jobi* 248
 var. *ma-yuen* (Romanet) Stapf. ex Hook. f. 248
Colocasia Schott 191
affinis Schott. 191
antiquorum Schott 191
esculenta L. 191
fallax Schott 192
navicularis Koch. & Bouche 186
Colpodium Trin 285
 wallichii (Hook. f.) Bor 285
COMMELINACEAE 167
Commelina L. 168
 appendiculata Clarke 168
 axillaris L. 167
 benghalensis L. 168
 communis Dalz. 168
 diffusa Burm. f. 168
 edulis Stokes 170
 elata Vahl 170
 hasskarlii Clarke 168
 japonica Thunb. 170
 maculata Edgew.
 nudiflora L. 168, 171
 nudiflora sensu Hook. f.
 obliqua Buch. ex D. Don 169
 var. *viscida* 169
 paludosa Bl. 169
 scaberrima Bl. 172
 scapiflora Roxb. 170
 sikkimensis Clarke 169
 suffruticosa Bl. 169
 vaginata L. 171
Compsoea maculata D. Don 162
Conchidium pusillum Griff. 69
Convallaria cirrhifolia Wall. 158
 oppositifolia Wall. 159
 punctata Wall.
 verticilata L. 160
Corallorrhiza foliosa Londl. 96
Cordyline Comin. ex A.L. Jun. 148
 terminalis Kunth 148
Corybas Salisb. 50
 himalaicus (King & Pantl.) Schltr. 50
Corymborkis Du Petit Thouars 50
 veratrifolia (Reinw.) Bl. 50
Corynbia veratrifale Reinw. 50
Corysanthes himalaiia King & Pantl. 50
Costus L. 124
 lacerus Gagnep. 124
 speciosus (Koen) J.E. Smith 124
Cottonia championii Lindl. 64
Courtoisia cyperoides Roxb. 222
Cremastera Lindl. 51
 appendiculata (D. Don) Makino 51
 wallichiana Lindl. 51
Crinum L. 137
 amoenum Roxb. ex Ker - Gawler 137
 powellii Baker 138
Crocasmea Planth. 136
Crocosmiflora (Lemonine) N.E. Br.
Cryptochilus Wall. 51
 lutea Lindl. 51
 meirax Par. & Reichb. f. 108
 sanguineus Wall. 51
Curculigo Gaertn. 139
 brevifolia Dryand. 139
 capitulata (Lour.) O. Kuntze 139
 crassifolia (Baker) Hook. f. 139
 gracilis (Kurz) Wall. ex Hook. f. 139
 orchioides Gaertn. 139
 recurvata 139
Curcuma Roxb. 125
 angustifolia Roxb. 125
 aromatica Salisb. 125
 caesia Roxb. 125
 domestica Valet
 longa auct. 125
 zedoaria (Christni.) Rosc. 126
Curcumorpha Rao & Verma 126
 longiflora (Wall.) Rao & Verma 126
Cyanotis D. Don 169
 axillaris (L.) J.A. & J.H. Schult. 167
 barbata D. Don 169
 cristata (L.) D. Don 169
 nepalensis Miers ex Wall.
 vaga (Lour.) J. A. & J.H. Schult. 169
Cyathopus Stapf. 285
 sikkimensis Stapf. 285
Cyathorhachis wallichiana Nees ex Steud. 268
Cymbidium Sw. 51
 aloifolium (L.) Swartz. 51
 appendiculatum D. Don 51
 bicolor 52
 ssp. *obtusum* Du Puy & Cribb 52
 bituberculatum Hook.
 cochleare Lindl. 52
 cyperifolium Lindl. 52

- dayanum** Reichb. f. 52
devonianum Paxt. 52
eburneum Lindl. 53
 var. *dayana* (Reichb. f.) Hook. f. 52
elegans Lind. 53
ensifolium (Lindl.) Swatz. 53
gammieanum King & Pantl. 53
giganteum Lindl. 54
 var. *lowiana* Reichb. f.
grandiflorum Griff. 53
hookerianum Reichb. f. 53
imbricatum Roxb. 103
iridioides D. Don
ixioides D. Don 111
lancifolium Hook. 54
longifoliu D. Don 54
lowianum Reichb. f.
mastersii Griff. ex Lindl. 54
munronianum King & Pantl. 53
nitidum Wall. ex D. Don 48
pendulum (Roxb.) Sw. 52
pendulum sensu King & Pantl.
sikkimense Hook. f. 54
simonsianum King & Pantl. 52
tridioides D. Don 54
whiteae King & Pantl. 55
- Cymbopogon** Spr. 248
flexuosus (Nees ex Steud.) Wats. 248
 var. *flexuosus* 248
khasianus (Hack.) Stapf. ex Bor 249
martinii (Roxb.) Wats 249
microtheca (Hook.) A. Camus 249
motia B.K. Gupta
pendulus (Nees ex Steud.) Wats 249
- Cyanodon** L.C. Rich. 285
dactylon (L.) Pers. 285
- Cyanosurus aegyptius** L. 285
 coracana L. 287
 indicus L. 288
- CYPERACEAE** 198
- Cyperorchis cochleare** (Lindl.) Benth. 52
 elegans (L.) Bl. 53
 longifolia (D. Don) Schltr. 54
 mastersii (Griff. ex Lindl.) Benth. 54
- Cyperus** L. 213
 alulatus Kern 213
 aristatus Rottb. 230
 brevifolius (Rottb.) Hassk. 227
 capillaris Koen. ex Roxb. 231
 castaneus Willd. 214
 compactus Retz. 228
 compressus L. 214
 var. *pectiniformis* (R.R.S.) Clarke 214
- corymbosus* Rottb. 216
 var. *pangorei* (Rottb.) Clarke 216
 cuspidatus Kunth 214
 cypdroides 230
 var. *evolutior* (Clarke) Kükenth. 231
- cyperinus* (Retz.) Valck
 var. *cyperinus*
 var. *benghalensis* (Clarke) Kükenth
 cyperoides (L.) O. Ktze
 diaphanus Schrad. ex R. & S. 231
 var. *latespicatus* (Boeck.) Kern. 231
- difformis* L. 214
 diffusus Vahl 216
 dilutus Valhl 228
 distans L. f. 214
 dubius Rottb.
 exaltatus Retz. 215
 flavidus Retz. 217
 flavidus sensu Clarke 231
 var. *erectus* (Clarke) Korla 231
 globosus All. 231
 var. *erectus* (Clarke) Kükenth. 231
- haspan* L. 215
 hookeri Boeck.
 hookerianus Arn.
 involucratus Rottb. 215
 jria L. 215
 var. *rectangularis* Kükenth. 213
 var. *parviflorus* (Nees) Miq. 215
 kamphoeveneri Boeck. 215
 latespicatus Boeck. 231
 laxus Lam. 215
 melanocephalus Miq. 232
 niveus Retz. 216
 nutans Vahl 216
 obliquus Nees 216
 pangorei Rottb. 216
 paniceus (Rottb.) Boeck. 230
 paniciformis Franch. & Sav. 215
 parviflorus Nees 215
 pectiniformis R. & S. 214
 pilosus Vahl 216
 var. *obliquus* Nees 216
 var. *polyantha* Clarke 216
- pumilus* L. 232
 rectangularis (Kükenth) Bennet 213
 rotundus L. 217
 sanguinolentus Vahl 232
 ssp. *sanguinolentus* & f. *melanocephalus* (Miq.) Kükenth. 232
 ssp. *melanocephalus* (Miq.) Karthik. et al. 232
- squarrosum* L. 230

- sikkimensis* Kükenth
silletensis Nees 217
stramineus Nees 233
substramineus Kükenth. 233
sulcinux Clarke 233
tenuiculmis Boeck. 217
tenuispica Steud. 217
tegetum Roxb. 216
umbellatus Clarke
 var. *laxatus* Clarke
uncinatus sensu Clarke 214
uniloides R. Br. 233
zollingeri sensu Clarke 217
 var. *condensatus* Kükenth. 217
- Cypripedium** L. 55
elegans Reichb. 55
himalaicum Rolfe *apud* Hemsl. 55
macranthon Sw. 55
tibeticum King *ex* Hemsl. 55
venustum Sims 98
- Cyrtococcum** Stapf. 250
accrescens (Trin.) Stapf. 250
longipes (Wt. & Arn. *ex* Hook. f.) A.
- Camus** 250
 oxyphyllum (Steud.) Stapf. 250
 patens (L.) A. Camus 250
- Cyrtopera bicarinata** Lindl. 70
candida Lindl. 71
mannii Rechb. 71
sanguinea Lindl. 72
altissima Bl.
- Dactyloctenium** Willd. 285
aegypticum (L.) Willd. 285
- Dactylorhiza** Necker *ex* Nevski
hatagirea (D. Don) Soo
- Daemonorops** Bl. 182
jenkinsiana Griff. 182
- Danthonia** DC. 286
cumminsii Hook. f. 286
jacquemontii Bor 286
- Dendrobium** O. Swartz. 55
aduncum Wall. 55
aggregatum
 var. *jenkinsii* King & Pantl. 56
- amoenum* Wall. 56
amplum Lindl. 64
anceps Sw. 56
aphyllum (Roxb.) C.E.C. Fischer. 56
- bicameratum* Lindl. 56
holboellus Falc. 56
caespitosum King & Pantl. 60
calceolaria Cary *ex* Hook.
candidum Wall. *ex* Hook. 56
- cathcartii* Hook. p. 57
chrysanthum Wall. *ex* Lindl. 57
clavatum Wall. *ex* Lindl. 57
crepidatum Lindl. 57
cumulatum Lindl. 57
demissum D. Don 97
denneanum A.F.G. Kerr. 57
densiflorum Wall. *ex* Lindl. 58
denudans D. Don 58
eriaeflorum Griff. 58
falconeri Hook. 58
farmeri Paxt. 58
fimbriatum Hook.
 var. *occulta* Hook.
- formosum* Roxb. *ex* Lindl. 59
fuscescens Griff. 64
gibsonii Lindl. 59
heterocarpum Wall. 59
hookerianum Lindl. 59
leopardianum 35
longicornu Lindl. 59
macraci Lindl. 72
moschatum Sw.
muscicola Lindl. 68
nobile Lindl. 60
ochreatum Lindl. 60
parishii Reichb. f. 60
pauciflorum King & Pantl. 60
peguanum Lindl. 61
pierardi Roxb. 56
porphyrochilum Lindl. 60
primulinum Lindl. 61
pubescens Hook. 69
ramosum Lindl. 61
rotundatum (Lindl.) Hook. f. 64
striatum Griff. 39
stuposum Lindl. 61
sulcatum Lindl. 61
teres Roxb. 98
terminale Par. & Reichb. 61
transparens Lindl. 62
wardianum Warner 62
williamsoni Day & Reichb. 62
- Dendrocalamus** Nees 238
hamiltonii Nees & Am. *ex* Munro 238
hookeri Munro 238
patellaris Gamble 238
sikkimensis Gamble 239
- Deschampsia** P. beauv. 286
caespitosa (L.) P. Beauv. 286
- Deyeuxia** P. Beauv. 286
abnormis Hook. f. 286
debilis (Hook. f.) Veldk. 286

- filiformis* (Griseb.) Hook. f. 287
nivicola Hook. f. 286
pulchella (Griseb.) Hook. f. 287
scabrescens (Griseb.) Duthie 287
 var. *humilis* (Griseb.) Hook. f. 287
tibetica Bor 287
treutleri (O. Ktze.) Stapf. 279
triariastata Hook. f. 279
Dianella Lamk. 148
eniifolia L. 148
caerulea Sim. 148
Dichanthium Willemet. 251
annulatum (Forssk.) Stapf. 251
assimilis (Steud.) Despande 247
bladhii (Retz.) W.D. Clayton 245
caricosum (L.) A. Camus 251
odoratum (Donna Liiboa) Jain &
 Despande 245
parviflorum (R. Br.) De Wet 247
Dictyospermum *scaberrimum* (Bl.) Morton
 ex Panigr. 172
Didiciea King & Prain ex King & Pantl. 62
cunninghami King & Prain ex King &
 Pantl. 62
Didymoplexis Griff. 63
pallens Griff. 63
Dienia *congesta* Lindl. 90
cylindrostachya Lindl.
muscifera Lindl. 91
Digitaria Haller. 251
adscendens H.B.K. 251
bicornis Willd. 251
ciliaris (Retz.) Koel. 251
compacta (Roth ex R. & S.) Veldk.
 251
cruciata (Nees ex Steud.) A. Camus 252
paspalodes Michx. 265
radicosa (C.B. Presl.) Miq. 252
sanguinalis (L.) Scop. 252
setigera Roth *apud* R. & S. 252
stricta Roth ex R. & S. 253
 var. *denudata* (Link.) Henr. 253
 var. *glabrescens* Bor. 253
timorensis Kunth 252
Diglyphosa Bl. 63
macrophylla (King & Pantl.) King &
 Pantl. 63
Dimeria R. Br. 253
ornithopoda Trin 253
DIOSCOREACEAE 141
Dioscorea L. 141
alata L. 141
deamona Roxb. 142
- esculenta* (Lour.) Burkill. 141
glabra Roxb. 141
hamiltonii Hook. f. 141
hispida Dennst. 142
kamoonensis Kunth. 142
prazeri Prain & Burkill 142
sativa L. 141
sikkimensis Prain & Burkill 142
Diphylax Hook. f. 63
urceolata (Clarke) Hook. f. 63
Diplochilos *hirsutus* Lindl. 63
Diplomeris D. Don 63
hirsuta (Lindl.) Lindl. 63
Diploprora Hook. f. 64
championii (Lindl.) Hook. f. 64
Diplosiphon *oryzetorum* Decne
Disporum Salisb. 148
calcaratum D. Don 148
cantonense 149
 var. *parviflorum* (Wall.) Hara 149
leucanthum Hara 149
wallichii D. Don 148
Doritis *taenialis* (Lindl.) Hook. f. 82
wightii (Reichb. f.) Benth. 82
Draceaena Vandelli ex L. 140
angustifolia Roxb. 140
ensifolia 148
Draconitum *spinosum* L. 192
Drepanostachyum Keng. f. 239
falconeri (Hook. f. ex Munro) J.
 Campbell 240
gracilis (Blanchard) Kumar 239
intermedium (Munro) Keng. 239
jainianum (Das & Pal) R. Majumdar
 239
polystachyum (Kurz ex Gamble) R.
 Majumdar 240
suberectum (Munro) R. Majumdar 240
Echinochloa P. Beauv. 253
colonum (L.) Link 253
crusgalli (L.) P. Beauv. 254
frumentacea Link 254
glabrescens Munro ex Eggel. 254
Echinoglossum *striatum* Reichb. f. 47
Echinolaena *polystachya* H.B.K. 268
Eichhornia Kunth
crassipes (Mart.) Solms. 166
Elate *sylvestris* L. 183
Eleocharis R. Br. 218
Eleocharis *afflata* Steud. 218
chaetaria 218
 var. *subvivipara* (Boeck.) C.E.C.
 Fisher 218

- congesta* D. Don 218
dulcis (Burm. f.) Trin *ex* Henschel 218
palustris (L.) R. Br. 218
plantaginea (Retz.) R. & S. 218
subvivipara Boeck. 218
tetraquetra Nees 219
Elettaria linguisomis Schult. 131
Eleusine Gaertn. 287
 aegyptia (L.) Desf. 285
 coracana (L.) Gaertn. 287
 indica (L.) Gaertn. 288
Elymus L. 288
 nutans Griseb. 288
 semicostatus (Nees *ex* Steud.) Melderis
 288
 sibiricus *auct.* 288
 sikkimensis (Melderis) Melderis 288
Elyna capillifolia Decne 222
 humilis C.A. Mey *apud* Tranty 226
 schoenoides C.A. Mey 226
Empusa paradoxa Lindl. 84
Epidendrum aloifolium L. 51
 ensifolium 53
 humile J.E. Smith 106
 pendulum Roxb. 52
 praecox J.E. Smith 106
 retusum L. 109
Epigeneium Gagnep. 64
 amplum (Lindl.) Summerh 64
 fuscescens (Griff.) Summerh 64
 rotundatum (Lindl.) Summerh 64
Epipactis Adans 64
 consimilis Wall. *ex* Hook. f. 65
 helleborine (L.) Crantz. 64
 latifolia All. 65
 veratrifolia Boiss. & Hohen. 65
Epipogium Gmelin *ex* Borkhausen 65
 aphyllum (F.W. Schmidt.) Swartz 65
 nutans Reichb. f. 65
 roseum (D. Don) Lindl. 65
Eragrostis Wolf. 289
 amabilis Stapf. 292
 atrovirens (Desf.) Trin. *ex* Steud. 289
 coarctata Stapf. 289
 curvula (Schrad.) Nees. 289
 diarrhena (Schult.) Steud. 290
 diplachnoides Steud. 290
 ferruginea (Thunb.) P. Beauv. 289
 gangetica (Roxb.) Steud. 289
 interrupta 290
 var. *tenuissima* Stapf. 290
 japonica (Thunb.) Trin. 290
 minor Host 290
 nigra Nees *ex* Steud. 290
 nutans (Retz.) Nees *ex* Steud. 290
 pappiana Chiov. 290
 papposa (Roem. & Schult.) Steud. 291
 pilosa (L.) P. Beauv. 291
 poaeoides P. Beauv. 290
 stenophylla Hochst *ex* Miq. 289
 stenophylla *auct.* 290
 tenella (L.) B. Beauv. *ex* R. & S. 291
 var. *plumosa* (Retz.) Stapf. 291
 var. *viscosa* (Retz.) Stapf. 292
 tremula (am.) Steud. 291
 unioloides (Retz.) Steud. 292
 viscosa (Retz.) Trin 292
Eremopogon (Hook.) Stapf. 284
Eremopogon foveolatus (Del.) Stapf. 254
Eria Lindl. 65
 acervata Lindl. 65
 alba Lindl. 66
 amica Reichb. f. 66
 andersonii Hook. f. 66
 bambusifolia Lindl. 66
 biflora Griff. 66
 bractescens Lindl. 66
 carinata Gibbs 67
 clausa King & Pantl. 67
 confusa Hook. f. 66
 convallarioides Lindl. 69
 coronaria (Lindl.) Reichb. f. 67
 dasyphylla Par. & Reichb. f. 116
 discolor Lindl. 117
 elongata Lindl. 68
 elwesii Reichb. f.
 excavata Lindl. 67
 ferruginea Lindl. 67
 fibuliformis King & Pantl. 108
 flava Lindl. 69
 fragrans Reichb. f. 68
 graminifolia Lindl. 67
 javanica (Sw.) Bl. 68
 laniceps Reichb. 68
 meirax Hook. f. 108
 muscicola (Lindl.) Lindl. 68
 paniculata Lindl. 68
 pannea Lindl. 68
 pubescens (Hook.) Lindl. 69
 pulchella *sensu* Hook. f. 117
 pulvinata Lindl. 116
 pumila Lindl. 69
 pusilla (Griff.) Lindl. 69
 rufinula Reichb. f. 116
 spicata (D. Don) Hand. - Mazz. 69
 stricta Lindl. 69

- vittata* Lindl. 70
Erianthus Nees ex Steud. ex Hack.
fastigatus Nees ex Steud. 254
fulvus Nees ex Hack. 271
hookeri Hack. 270
longisetosus Anders. ex Benth. 270
 var. *longisetosum* 270
 var. *hookeri* (Hack.) Bor 270
mollis Griseb. 255
narenga Nees ex Steud 270
nudipes Griseb. 259
rufipilus (Steud.) Griseb. 271
sikkimensis Hook. f. 271
versicolor Nees ex Steud. 271
 var. *palens* (Hack.) Hook. f. 271
ERIOCAULACEAE 197
Eriocaulon L. 197
Eriocaulon alpestre Hook. f. & Thoms. ex Koern. 197
luzulaefolium Mart. 197
nepalense Prescott. ex Bong. 197
oryzetorum Mart. 197
sieboldianum Steud. 197
truncatum Ham. ex Mart. 198
Eriophorum L. 219
 comosum (Wall.) Wall. ex Nees 219
 microstachyum Boeck. 219
Erythrorchis lindleyana Reichb. f. 73
Esmeralda 70
Clarkei Reichb. f. 70
Eulalia Kunth 254
cumingii (Nees) A. Camus 255
 fastigata (Nees ex Steud.) Haines 254
 hirtifolia (Hack.) A. Camus 255
 leschenaultiana (Decne) Ohwi 255
 mollis (Griseb.) O. Ktze. 255
 nepaleusis Trin. 259
 guadrinervis (Hack.) O. Ktze. 255
Eulaliopsis Honda 256
binata (Retz.) C.E. Hubb. 256
Eulophia R. Br. 70
 bicallosa (D. Don) P.F. Hunt & Summerh. 70
 bicarinata (Lindl.) Hook. f. 70
 var. *major* King & Pantl. 70
 var. *bicallosa* 70
 var. *major* (King & Pantl.) Sarkar 70
campestris Wall. ex Lindl.
candida (Lindl.) Hook. f. 71
dubia (D. Don) Hooker.
densiflora Lindl. 71
geniculata King & Pantl. 71
graminea Lindl. 71
macrobulbon Par. & Reichb. f. 71
macrorhizon Hook. f. 72
mannii (Reichb. f.) Hook. f. 71
nuda Lindl. 72
sanguinea (Lindl.) Hook. 72
Euproboscus pygmaea Griff. 115
Fargesia maling (Gamble) J. Campbell 242
pantlingii (Gamble) J. Campbell 243
racemosa (Munro) Yi 243
Festuca L. 292
 coelestis (St. Yves) Krecz. & Bobrov. 292
cumminsii Stapf. 292
leptopogon Stapf. 293
ovina L. 292
 ssp. *coelestis* St. Yves 292
 ssp. *sulcata* Hack. 294
 ssp. *polycolea* (Stapf.) St. Yves. 293
 ssp. *valesiaca* (Schleich. ex Gaud.) Koch. 293
parvigluma Steud. 293
polycolea Stapf. 293
rubra L. 293
sylvatica Huds. 282
undata Stapf. 294
 var. *undata* 294
 var. *aristata* Stapf. 294
valesiaca Schleich ex Gaud. 294
 var. *tibetica* Stapf. 292
Fimbristylis Vahl 219
aestivalis (Retz.) Vahl 219
annua R. & S. 220
 var. *paucispiculata* Blatt. & McC. 220
barbata 198
bisumbellata sensu C.E.C. Fischer 219
bisumbellata (Forssk.) Bub. Dodec. 220
capillaris
 var. *trifida* (Nees) Mitra 198
complanata (Retz.) Lin. 220
dichotoma sensu Clarke 220
dichotoma (L.) Vahl 220
diphylla Vahl 220
 var. *annua* (R. & S.) Clarke 220
litoralis Gaud. 220
miliacea sensu Benth. 220, 221
schoenoides (Retz.) Vahl 221
stolonifera Clarke 221
thomsonii Boeck. 221
Flickingeria Hawkes 72
macraei (Lindl.) Seid. 72
Floscopa Lour. 170
scandens Lour. 170
Fluggea dracaenoides Baker 155

- wallichiana* Kunth. 156
Forrestia glabrata Hassk. 168
hookeri Hassk. 167
marginata 168
 var. *rostrata* Clarke 168
mollis Hassk. 168
mollissima (Bl.) Kds. 168
Fritillaria L. 149
cirrhosa D. Don 149
delavayii Franch. 149
flavida Rendle 154
macrophylla D. Don 155
thomsonianum Royle 155
Furcraea Vent. 140
foetida 140
Gagea Salisb. 149
Gagea elegans Wall. ex D. Don 149
lutea auct. 149
Galeola Lour. 72
altissima Reichb. f.
cathcartii Hook. f. 72
falconeri Hook. f. 73
hydra Reichb. f. 73
lindleyana (Reichb. f.) Reichb. f. 73
nudiflora Lour. 73
Garnotia Brongn. 294
acutigluma (Steud.) Ohwi 294
 var. *acutigluma*
var. longiseta (Hack.) Ved Prakash & Jain
polypogonoides Munro ex Oliver 295
stricta auct. 295
stricta sensu Hook. f. 294
 var. *longiseta* Hack. 294
tenella (Arn. ex Miq.) Jan. 295
Gastrochilus D. Don. 73
acutifolius (Lindl.) O. Ktze. 73
affinis (King & Pantl.) Schltr. 73
calceolaris (J.E. Smith) O. Ktze. 74
dasygagan J.E. Smith 74
distichus (Lindl.) O. Ktze. 74
intermedius (Griff. ex Lindl.) O. Ktze. 74
longiflora Wall. 126
obtusifolius (Lindl.) A.S. Rao & Mukherjee 117
pseudodistichus (King & Pantl.) Seid. 75
Gastrodia R. Br. 75
dyeriana King & Pantl. 75
Geodorum G. Jackson 75
densiflorum (Lam.) Schltr. 75
dilatatum sensu Hook. f. 75
purpureum R. Br. 75
Georchis foliosa Lindl. 75
Gladiolus L. 136
Globba L. 126
andersonii Clarke ex Baker 126
clarkei Baker 126
expansa Wall. ex Baker 127
hookeri Clarke ex Baker 127
multiflora Wall. ex Baker 127
orixensis Roxb. 127
racemosa J.E. Smith 127
 var. *hookeri* 127
 var. *racemosa* 127
siamensium (Koen.) Rao & Verma 127
versicolor J.E. Smith 127
Gloriosa L. 150
superba L. 150
Glossula tentaculata Lindl. 100
Glyceria R. Br. 295
tentaculata Lindl. 295
kasmiriensis L. Kelso 295
tonglensis Clarke 295
Gonanthus P. Klotzsch.
pumilus (D. Don) Engler & Krause 193
sarmentosus Klotzsch & Otto. 193
Gonyanthes nepalensis Miers. 24
Goodyera R. Br. 75
dongchenii Lucksom
foliosa (Lindl.) Benth. ex Clarke 75
fumata Thw. 75
fusca (Lindl.) Hook. f. 76
grandis King & Pantl. 76
hemsleyana King & Pantl. 76
hispida Lindl. 76
procera (Wall. ex Ker-Gawl.) Hook. 76
repens (L.) R. Br. 77
schlechtendaliana Reichb. f. 77
secundiflora Lindl. 77
vittata Benth. ex Hook. f. 77
Govindovia nervosa Wt. 113
Gymnadenia R. Br. 77
orchidis Lindl. 77
spathulata Lindl.
Habenaria Willd. 77
affinis D. Don 99
atchinsonii Reichb. f. *apud* Aitch. 77
albomarginata King & Pantl. 78
arietina Hook. f. 78
aristata (Lindl.) Trin. 99
bakeriana King & Pantl. 104
biermanniana King & Pantl. 104
constricta (Lindl.) Hook. f. 99

- cumminsiana** King & Pantl. 78
densa Wall. ex Lindl. 104
dentata (Sw.) Schltr. 78
 var. *ecalcarata* R.C. Srivast. 78
 f. *ecalcarata* (King & Pantl.)
 Tuyama 78
dyeriana King & Pantl. 104
fallax (Lindl.) King & Pantl. 100
furfuracea Hook. f. 79
galendra auct. 31
geniculata D. Don 78
 var. *ecalcarata* King & Pantl.
goodyerooides D. Don 100
junccea King & Pantl. 104
latilabris (Lindl.) Hook. f. 105
leptocaulon Hook. f. 105
malleifera Hook. f. 79
nematocaulon Hook. f. 79
oligantha Hook. f. 104
orchidis (Lindl.) Hook. f. 77
parishii (Reichb. f.) Hook. f. 100
pectinata (J.E. Smith) D. Don 79
 var. *arietina* (Hook.) R. C. Srivast.
polytricha (Hook. f.) Pradhan 79
prainii Hook. f. 100
pseudophrys King & Pantl. 100
secundiflora Hook. f. 92
seshagiriana A.N. Rao 79
sikkimensis Hook. f. 105
stenantha Hook. f. 105
stenopetala Lindl. 79
 var. *polytricha* Hook. f. 79
 var. *stenopetala* 79
tentaculata (Lindl.) Reichb. f. 100
urceolata Clarke 63
Hackelochloa granularis (L.) Ktze. 260
porifera (Hack.) Rhind 260
Hedychium Koen. 128
 acuminatum Rosc. 130
aurantiacum Wall. ex Rosc. 128
coccineum Buch. - Ham. ex J.E. Smith
 128
 var. *coccineum* 128
 var. *longifolium* Baker 128
 var. *squarrosum* Buch. - Ham. ex Wall.
 128
coronarium Koen. 128
densiflorum Wall. 129
ellipticum Buch. - Ham. ex J.E. Smith
 129
gardnerianum Sheppard ex Ker-Gwl.
 129
gracile Roxb. 129
 var. *glaucum* Baker 129
greeni W.W. Smith 130
spicatum Buch. - Ham. ex J.E. Smith
 130
 var. *spicatum* 130
 var. *acuminatum* (Rosc.) Wall. 130
squarrosum Buch. - Ham. ex Wall. 129
thyrsiforme Buch. - Ham. ex J.E. Smith
 130
helictotrichon J.A. & J.H. Schultes 295
asperum (Munro ex Thw.) Bor 295
parviflorum (Hook. f.) Bor 295
virescens (Nees ex Steud.) Henr. 295
Hemarthria R. Br. 256
compressa (L.f.) R. Br. 256
Hemerocallis L. 150
fulva (L.) L. 150
Hemicarex filicina Clarke
pygmaea Clarke 225
Hemiorchis Kurz 130
Hemiorchis pantlingii King 130
Heritiera allughas Ketz.
Herminium L. 80
angustifolium (Lindl.) Benth. ex Hook.
 f. 80
angustilabre King & Pantl. 80
congestum Lindl. 80
constrictum Lindl. 99
fallax (Lindl.) Hook. f. 100
jaffreyanum King & Pantl. 80
josephii Reichb. f. 80
lanceum (Thunb. ex Swartz) J. Vuijk 80
macrophyllum (D. Don) Dandy 80
orbiculare Hook. f. 81
pugioniformae Lindl. ex Hook. f. 81
quinquaelobum King & Pantl. 81
Herpsma Lindl. 81
longicaulis Lindl. 81
Hetaeria Bl. 81
flava Lindl. ex Wall. 119
rubens (Lindl.) Benth. 81
Hierochloa flexuosa Hook. f. 280
gracillima Hook. f. 280
hookeri (Griseb.) Clarke ex Hook. f. 280
sikkimensis Maxim. 280
Himalayacalamus Keng f. 240
falconeri (Hook. f. ex Munro) Keng f.
 240
Holcus L. 296
lanatus L. 296
natans Roxb. ex Hook. f. 269
parviflorus R. Br. 247
Hordeum L. 296

- aegiceras** Nees *ex* Royle 296
vulgare L. 296
 ssp. *aegiceras* (Nees *ex* Royle) A. Love 296
 var. *aegiceras* Aitch. 296
 ssp. *vulgare* L. 296
 ssp. *agriocritthon* (Aberg) A. Love 296
- Hornstedtia** Retz. 131
 linguifomis K. Schum 131
- Hura** *siamensis* Koen. 127
- Hymenocallis** Salish 138
 littoralis Salisb. 138
- Hypolytrum** L.C. Rich.
 argenteum Vahl
 latifolium L.C. Rich
 nemorum (Vahl) Spr.
 trinervium Kunth
 wightianum Boeck.
- HYPOXIDACEAE** 139
- Hypoxis** *aurea* Lour.
Hysteria *veratrifolia* Reinw. 50
- Ichnanthus** P. Beauv. 256
 Cicinus (F.M. Bail.) Merr. 256
 Pallens Hook. f. 256
- Imperata** Cyr. 257
 arundinacea Cyr. 257
 cylindrica (L.) Reauschel 257
 var. *major* (Nees) C.E. Hub. *ex* C.E. Hubb. *et* Vaughan 257
- Indocourtosia** Bennet & Raiz. 221
 cyperoides Roxb. 221
- Iphigenia** Kunth 150
 indica (L.) Kunth 150
- Ione** *andersonii* King & Pantl. 112
 bicolor (Lindl.) Lindl. 112
 intermedia King & Pantl. 112
 paleacea Lindl. 112
 scariosa (Lindl.) King & Pantl. 113
- Ipsea** *wrayana* Hook. f. 115
- IRIDACEAE** 136
- Iris** L. 137
 clarkei Baker *ex* Hook. f. 137
 decora Wall. 137
 goniocarpa Baker 137
 himalaica W.R. Dykes 137
 nepalensis D. Don 137
 tectorum Maxim. 137
- Isachne** R. Br. 296
 albens Trin 296
 clarkei Hook. f. 297
 himalaica Hook. f. 297
 miliacea Roth *ex* R. & S. 297
 sikkimensis Bor 297
- Ischaemum** L. 257
 angustifolium (Trin) Hack. 256
 aristatum auct. 257
 aristatum sensu Hack. 257
 ssp. *imberbe* Hack. 257
 var. *imbricatum* Hack. 257
- barbatum** Retz. 257
 ciliare Retz. 257
 goebelii Hack. 257
 indicum (Houtt.) Merr. 257
 paleaceum Trin 244
 petiolaris (Trin) Hack. 258
 rugosum Salish. 257
- Isolepis** R. Br. 222
 setacea (L.) R. Br. 222
 trifida Nees 198
- Ixia** *chinensis* L. 136
- JUNCACEAE** 172
- Juncus** L. 172
 albescences Clarke 178
 alliodoides Franch. 172
 benghlensis Kunth 172
 bhutanensis Satake 173
 brachystigma Samuel. 173
 bracteatus Buchen. 173
 cephalostigma Samuel. 173
 chrysocarpus Buchen. 173
 clarkei Buchen. 174
 campestris L. 179
 concinus D. Don 174
 effusus L.
 elegans Royle *ex* D. Don 174
 glaucus Enrh. *ex* Sibth. 174
 var. *leptocarpus* (Buchen.) Buchen
 grisebachii Buchen 174
 himalensis Klotz. & Garcke 174
 inflexus L.
 var. *inflexus* 174
 var. *leptocarpus* (Buchen.) Sanjappa 174
 kingii Rendle 175
 latior Satake 175
 leschenaultii Gay 175
 leucanthus Royle *ex* D. Don 175
 var. *leucanthus* 175
 var. *alpinus* Buchen. 175
 leptocarpus Buchen. 175
 leucomelas Royle *ex* D. Don.
 leucomelas sensu Hook. f. 176, 177
 luteocarpus Satake 174
 membranaceus Royle *ex* D. Don 176
 membranaceus auct. 172
 minimus Buchen. 176

- monticola** Steud. 176
multiflorus Retz. 179
ochraceus Buchen. 176
prismatocarpus sensu Hook. f. 176
pseudocastaneus (Lingelsch) Samuel. 177
sikkimensis Hook. f. 177
 var. *pseudocastaneus* Lingelsch. *apud* Limpr. f. 177
sphacelatus Decne 177
sphenostemon Buchen. 172
thomsonii Buchen. 177
trichophyllum W.W. Smith 177
triglumis L. 178
uniflorus W.W. Smith 178
wallichianus Laharpe 178
yoskisukei Goel 178
Kaempferia L. 131
 angustifolia Rosc. 131
 invulnerata King *ex* Baker 131
 rotunda L. 131
 secunda Wall. 123
 sikkimensis King *ex* Baker 123
Kingidium P.F. Hunt 82
 deliosum (Reichb. f.) Sweet 82
 taenialis (Lindl.) P.F. Hunt 82
Kniphofia Moench. 150
Kobresia Willd. 222
 angusta 223
 capillifolia (Decne) Clarke 222
 clarkeana Kükenth. 222
 curticeps (Clarke) Kükenth. 223
 curvata (Boott) Clarke 223
 curvirostris Clarke 224
 duthiei Clarke 223
 esenbeckii (Kunth) Noltie 223
 var. *esenbeckii* 223
 var. *fissiglumis* (Clarke) Noltie 224
 gammieae Clarke 224
 govindarajalu Karthik.
 hookeri auct. 223
 laxa Nees 224
 var. *laxa* 224
 var. *curvirostris* (Clarke) R.C. Srivast. 224
 linearis Boott 224
 nepalensis (Nees) Kükenth. 224
 var. *elachista* Clarke 224
 var. *nepalensis* 225
 var. *vaginosa* (Clarke) Kükenth. 225
 persica Kükenth. & Bornm. 226
 prainii 225
 pygmaea (Clarke) Clarke 225
- royleana** (Nees) Boeck. 225
 var. *royleana* 225
 var. *humilis* (C.A. Mey) Kükenth. 226
schoenoides (C.A. Mey) Steud. 226
seticulmis Boeck. 223
sikkimensis Kükenth. 226
trinervis (Nees) Boeck. 223
 var. *foliosa* (Clarke) Kükenth
uncinoides (Boott) Clarke 226
vaginosa Clarke
williamisii T. Koyama
vidua (Boott *ex* Clarke) Kükenth. 227
Kyllinga Rottb. 227
 brevifolia Rottb. 227
 cylindrica Nees 227
 cyperina Retz. 228
 cypeoides Roxb. 221
 monocephala Rottb. 227
 nemorallis (J.R. & G. Forst.) Dandy *ex* Hutch. & Dalziel 227
 odorata Vahl 227
 ssp. *cylindrica* (Nees) T. Koyama 227
 panicea Rottb. 230
 sumatrensis Retz. 230
Lagurus cylindricus L. 257
Lasia Lour. 192
 heterophylla (Roxb.) Schott. 192
 spinosa (L.) Thwaites 192
LEMNACEAE
Lemna L.
 minor auct.
 perpusilla Torrey
 polyrhiza L.
Leucojum capitatum Lour. 139
LILIACEAE 142
Lilium L. 151
 giganteum Wall. 146
 nanum Klotz 154
 nepalense D. Don 151
 roseum Wall. *ex* Hook. f. 155
 wallichianum J.A. & J.H. Schultes 151
Limodorum aphyllum Roxb. 56
 densiflorum Lam. 75
 flavum Bl. 101
 roseum D. Don 65
 tankarvilleae Bank *ex* L' Herit 101
Liparis L.C. Rich 82
 bistriata Par. & Reichb. f. 82
 bituberculata (Hook.) Lindl. 84
 bhotanensis Griff. 82
 caespitosa (Thow.) Lindl. 83
 cathcartii Hook. f. 83
 cordifolia Hook. f. 83

- deflexa** Hook. f. 83
deliculata Hook. f. 83
gamblei Hook. f. 84
glossula Reichb. f. 84
griffithii Ridley 86
longipes Lindl. 86
lydialii Lucksom
mannii Reichb. f.
nepalensis Lindl. 85
nervosa (Thunb.) Lindl. 84
obscura Hook. f. 83
paradoxa (Lindl.) Reichb. f. 84
perpusilla Hook. f. 84
petiolata (D. Don) P.F. Hunt &
 Summerh 85
plantaginea Lindl. 85
platyrachis Hook. f. 85
pusilla Ridley 83
pygmaea King & Pantl. 85
resupinata Ridley 85
 var. *resupinata* 86
 var. *ridleyi* Hook. f. 86
stricklandia Reichb. f. 86
togashii Tuyama ex Hara 84
viridiflora (Bl.) Lindl. 86
- Lipocarpha** R. Br. 228
argentea (Vahl) R. Br. ex Nees 228
chinensis (Osb.) Kern 228
- Liriope** Lour.
graminifolia (L.) Baker
- Listera** R. Br. 86
alternifolia King & Pantl. 86
breviculmis King & Pantl. 87
dentata King & Pantl. 87
lindleyana King & Pantl. 92
longicaulis King & Pantl. 87
micrantha Lindl. 87
mucronata Panigr. & J.J. Wood 87
pinetorum Lindl. 87
tenuis Lindl. 88
- Lloydia** Salisb. 151
delavayi Franch. 151
deliculata Noltie 151
flavonutans Hara 151
himalensis Royle 152
longiscapa Hook. f. 152
serotina (L.) Reichb. L. 152
 var. *serotina* 152
 f. *parva* Marquand & Shaw 152
 var. *parva* (Marquand & Shaw) Hara
 var. *sikkimensis* 152
serotina auct. 151
tibetica Baker ex Oliver
- yunnanensis** Franch. 152
Lolium L. 297
 perenne L. 297
Luisia Gaud. 88
brachystachys (Lindl.) Bl. 88
inidivisa King & Pantl. 88
teretifolia 88
trichorhiza (Hook.) Bl. 88
volucris Lindl. 88
zeylanica Lindl. 88
- Luzula** DC. 179
australisica Steud.
chinensis N.E. Br. 179
campestris (L.) DC. 179
effusa Buchen. 179
multiflora (Retz.) Lejeune 179
oligantha Samuels. 179
parvifolia Buchen. 179
plumosa E. Meyer 180
rostrata Muchen. 180
 ssp. *darjeelingensis* Satake 180
sudetica DC. 179
- Malaxis** Soland ex Swartz. 89
acuminata D. Don 89
 f. *biloba* (Lindl.) Tuyama 89
 var. *acuminata*
 var. *biloba* (Lindl.) R.C. Srivast. 89
aphylla (King & Pantl.) Tang & Wang.
 89
caespitosa Thou. 83
calophylla (Reichb. f.) O. Ktze. 89
cylindrostachya (Lindl.) O. Ktze. 90
cyperoides Roxb. 272
ensiformis J.E. Smith 94
josephiana (Reichb. f.) O. Ktze.
khasiana (Hook. f) O. Ktze.
latifolia J.E. Smith
maximowicziana (King & Pantl.) Tang
 & Wang. 90
muscifera (Lindl.) O. Ktze. 91
saprophyta (King & Pantl.) Tang &
 Wang. 91
schottii Hook. f. 89
viridifolia Bl. 86
wallichii
 var. *brachycheila* Hook. f.
- Manisuris** granularis (L.) L.f. 260
porifera Hack. 260
- Maranta** malaccensis Bumf. 121
- Mariscus** Vahl 228
aristatus (Rottb.) Tang & Wang 230
compactus (Retz.) Boldingh 228

- cyperinus** (Retz) Vahl 228
 ssp. *laxatus* (Clarke) Koyama 228
- dilutus** (Vahl) Nees 228
- dregeanus** Kunth 229
- dubius** (Rottb.) C.E.C. Fischer 229
- hookerianus** Clarke 229
- microcephalus** Presl. 228
- paniceus** (Rottb.) Vahl 230
- sieberianus** Nees ex Clarke 230
 f. *evolutior* Clarke
 var. *sumatrensis* 230
 var. *evolutior* (Clarke) Karth. 230
- sikkimensis** Kükenth. 229
- sumatrensis** (Retz) Raynal 230
- tenuifolius** Schrad ex Nees 229
- Megastachya paposa** Roem. & Schult. 291
- Melanthium indicum** L. 150
- Melocalamus** Benth. 240
- Melocalamus indicus** R. Majumdar 240
 Mesoclastis uniflora Lindl. 98
- Micropera** Lindl. 91
 mannii (Hook. f.) Tang & Wang
 obtusa (Lindl.) Tang & Wang 91
- Microstegium** Nees 258
 ciliatum (Trin) A. Camus 258
 nudum (Trin) A. Camus 258
 petiolare (Trin) Bor. 258
 vagans (Nees ex Steud.) A. Camus 258
 vimineum (Trin.) A. Camus 259
 var. *willdenowianum* (Nees) Sur. 259
- Microstylis biloba** 89
 calophylla Reichb. f. 89
 congesta (Lindl.) Lindl. Reichb. f.
 cylindrostachya (Lindl.) Reichb. f. 90
 josephiana Reichb. f.
 khasiana Hook.
 latifalius J.E. Smith. 90
 maximoiwcziana King & Pantl. 90
 muscifera Lindl. 91
 sapropyhta King & Pantl. 91
 wallichii Lindl 89
- Milium treutleri** O. Ktze. 279
- Misanthus nepalensis** (Trin) Hack. 259
 nudipes Griseb.) Hack. 259
- Mnesithea** Kunth 260
 granularis (L.) Koning & Sosef 260
 khasiana (Hack.) Koning & Sosef 260
 striata (Nees ex Steud.) Koning & Sosef.
 260
- Molineria crassifolia** Baker 139
 gracilis Kurz 139
- Monochilus affine** Lindl. 119
 flavum Wall. ex Lindl. 119
- goodyerooides** Lindl. 119
- nervosum** Wall. 119
- Monochoria** K.B. Presl. 166
 vaginalis (Burm. f.) K.B. Presl. 166
- Monomeria** Lindl. 91
 barbata Lindl. 91
- Muhlenbergia** Sehreb. 298
 huegelii Trin 298
 viridissima Nees ex Steud. 298
- Murdannia** Royle 170
 malabarica (L.) Brueck.
 divergens (Clarke) Brueck. 170
 edulis (Stokes) Faden 170
 japonica (Thunb.) Faden 170
 malabarica (L.) Brueck. 171
 var. *compressa* (Dalz.) Sant. & Jain
 171
- nudiflora** (L.) Brenan 171
- viginata** (L.) Brueck. 171
- MUSACEAE** 135
- Musa** L. 135
 balbisiana Colla 135
 dasycarpa Kurz 136
 hookeri King ex A.M. Cowan & J.M.
 Cowan 135
 paradisiaca 135
 ssp. *sapientum* (L.) K. Schum. 135
 sapientum sensu Baker 135
 sikkimensis Kurz 136
 velutina Wendl. & Drude 136
- Myrmecallis** Lindl. 92
 pumila (Hook. f.) Tang & Wang 92
- Narenga porphyrocoma** (Hance ex Trin)
 Bor 270
- Neottia** Guettard 92
 australis R. Br. 111
 listeroides Lindl. 92
 macrophylla D. Don 80
 procera Wall. ex Ker - Gawl. 76
 sinensis Pers. 111
- Neottianthe** (Reichb.) Schltr. 92
 secundiflora (Hook. f.) Schultr. 92
- nudum** Hook. f. Bl. 92
 grandiflorum Hook. f. 115
 nudum Hook. f. 92
 pulchrum
 var. *sikkimensis* Hook. f.
- Neyraudia** Hook. f. 298
 madagascariensis (Kunth) Hook. f. 298
 var. *zollingeri* Hook. f. 298
- reynaudiana** (Kunth) Keng & Hitchc.
 298
- Nomocharis** Franch. 154

- flavidum** (Rendle) Hara 154
nana (Klotz) Wilson 154
Notholirion Wall. ex Boiss. 155
bulbuliferum (Lingelsh) Stearn. 155
macrophyllum (D. Don) Boiss. 155
thomsonianum (Royle) Stapf. 155
- Oberonia** Lindl. 93
acaulis Griff. 93
auriculata King & Pantl.
brachystachys Lindl. 93
caulescens Wall. ex Lindl. 93
croftiana King & Pantl. 96
demissa Lindl. 93
denticulata Wight 93
emarginata King & Pantl. 94
ensiformis (J.E. Smith) Lindl. 94
falcata King & Pantl. 94
falconeri Hook. f. 94
gammiei King & Pantl. 93
iridifolia Lindl. 93
 var. *angustifolia* Hook. f. 93
 var. *brevifolia* Hook. f. 93
 var. *denticulata* Hook. f. 93
jenkinsiana Griff. ex Lindl. 94
lobulata King & Pantl. 95
longilabris King & Pantl. 93
micrantha King & Pantl. 94
obcordata Lindl. 95
pachyrachis Reichb. f. ex. Hook. f. 95
parvula King & Pantl. 96
prainiana King & Pantl. 95
pyrulifera Lindl. 95
recurva Lindl. 96
treutleri Hook. f. 95
Octomeris spicata D. Don 69
Odontochilus crispus (Lindl.) Hook. f. 27
 elwesii Clarke ex Hook. f. 27
 grandiflorus (Lindl.) Benth. 28
 lanceolatus (Lindl.) Bl. 28
 pumilus Hook. f. 92
Oncus esculentus Lour. 141
Ophiopogon Ker-Gawl. 155
 clarkei Hook. f. 155
 dracaenoides (Baker) Hook. f. 155
 graminifolia L.
 intermedium D. Don 156
 var. *parviflora* Hook. f. 156
 parviflorus (Hook. f.) Hara 156
 pallidus Wall. ex Kunth. 161
 wallichianus Kunth 156
Ophiuros Gaertn. f. 260
 exaltatus (L.) O. Ktze 260
- corymbosus* Gaertn. 260
Ophrys lancea Thunb. ex Swartz 80
 nervosa Thunb. 84
Oplismenus B. Beauv. 261
 burmannii (Retz.) P. Beauv. 261
 compositus (L.) P. Beauv. 261
 undulatifolius (Ard.) R. & S. 261
- ORCHIDACEAE**
- Orchis aphylla* F.W. Schmidt. 65
 chusua D. Don 107
 var. *nana* King & Pantl. 108
 dentata Sw. 78
 habenarioides King & Pantl. 77
 latifolia auct.
 obcordata Buch. - Ham. ex D. Don 31
 pectinata J.E. Smith 79
 puberula King & Pantl. 27
 hatagirea D. Don
 strateumatica 120
- Oreorchis** Lindl. 96
 micrantha Lindl. 96
 foliosa (Lindl.) Lindl. 96
- Ornitharium** Lindl.
- striatum* Lindl. 108
- Ornithochilus** (Lindl.) Wall. ex Benth. 96]
 diformis (Wall. ex Lindl.) Schltr. 96
 fuscus Wall. ex Hook. f. 96
- Ornithogehn luteum* L. 149
Orthoraphium roylei Nees 304
Oryzopsis aequiglumis Duthie ex Hook. f. 299
- Otochilus** L. 96
 albus Lindl. 96
 fuscus Lindl. 97
 orrectus Lindl. 97
- Pachystoma** Bl. 97
 pubescens Bl.
 senile (Lindl.) Reichb. f. 97
- Pancratium** L. 138
 vereendum Aiton 138
- PANDANACEAE** 184
- Pandanus** 184
 minor Hook. f.
 furcatus auct.
 var. *indica* Kurz
 furcatus 184
 nepalensis St. John 184
 sikkimensis St. John 185
- PANICOIDEAE** 243
- Panicum** L. 201
 accrescens Trin 250
 americanum L. 265
 auritum Presl. ex Nees 261

- austroasiaticum* Ohwi 264
barbatum Lam. 272
bengalense Spreng. 280
brevifolium L. 262
burmannii Retz. 261
ciliare Tetz. 251
colonum L. 253
compositum L. 261
cruciatum Nees ex Steud. 252
crusgalli L. 254
 var. *furmentaceum* Hook. f. 254
dactylon L. 285
decempedale O. Ktze 281
flavescens sensu Hook. f. 272
flavidum Retz. 264
forbesianum Nees ex Steud. 273
glaucum L. 266
humidorum Buch. - Ham. ex Hook. f.
 262
humile Nees ex Steud. 263
indicum L. 271
 var. *angustatum* Hook. f. 271
 var. *gracile* Hook. f. 271
interruptum Willd. 272
italicum L. 273
khasianum Munro ex Hook. f. 262
longipes Wt. & Arn. ex Hook. f. 250
miliaceum L. 262
miliare Lam. 263
montanum Roxb. 262
myosuroides R. Br. 272
notatum Sw. 262
ovalifolium Poir. 262
oxyphyllum Steud. 250
palmifolium Koen. 273
paludosum Roxb. 263
patens L. 250
pilipes Nees & Arn. ex Buese 250
plicatum Lam. 273, 274
proliferum sensu Hook. f. 263
prostratum Lam. 246
psilotopodium Trin 263
 var. *coloratum* Hook. f. 263
pumillum Poir 274
punctatum Burm. f. 264
punctatum (Burm. f.) A. Camus 264
radicosum C.B. Presl. 252
ramosum L. 246
repens L. 263
reptans L. 246
rhachitrichum Hochst. 272
sanguinale L. 252
semialatum R. Br. 243
- simpliciusculum* Steud. 284
sumatrense Roth ex R. & S. 263
tomentosum Roxb. 273
uncinatum Raddi. 268
undulatifolium Ard. 261
verticillatum L. 274
vicinum F.M. Bail 256
villosum Lam. 246
walense Mez. 263
Panisea (Lindl.) Lindl. 97
demissa (D. Don) Pfitz. 97
parviflora (Lindl.) Lindl. 97
Pantlingia Prain 98
paradoxa Prain 98
Paphiopedilum Pfitz. 98
fairieanum (Lindl.) Stein
hirsutissimum (Lindl. ex Hook.) Stein
insigne (Wall. ex Lindl.) Pfitz.
venustum (Sims) Pfitz. 98
Papilionanthe Schltr. 98
teres (Roxb.) Schltr. 98
uniflora (Lindl.) Garay 98
vandarum (Reichb. f.) Garay 99
Paradisia *bulbulifera* Lingelsch 155
Paris L. 156
marmorata Stearn. 157
polyphylla Smith 156
 var. *brachystemon* Franch.
 ssp. *polyphylla*
 var. *wallichii* Hara 156
 var. *appendiculata* Hara
 var. *polyphylla*
 var. *Stenophylla* Franch. 157
 var. *Wallichii* Hara 156
 ssp. *marmorata* (Stearn.) Hara
- Paspalidium** Stapf. 264
flavidum (Retz.) A. Camus 264
punctatum Burm. f. 269
Paspalum L. 264
cartilagineum J.S. ex C.B. Presl. 265
commersonii Lam. 265
compactum Roth ex R. & S. 251
conjugatum Berg. 264
dilatatum Poir 264
distichum auct. 265
orbiculare Forst. 265
paspalodes (Michx.) Scribner 265
royleanum Nees ex Hook. f.
sanguinale (L.) Lam. 251, 252
 var. *calliblepharata* (Henr.) Veldk. 253
 var. *ciliare* (Retz.) Hook. f. 251
 var. *cruciatum* Nees ex Steud. Hook.
 f. 252

- var. extensum* Hook. f. 252
var. rottleri Hook. f.
scrobiculatum L. 265
Peliosanthes Andr. 157
griffithii Baker 157
macrophylla Wall. ex Baker 157
teta Andr. 157
Pennisetum Rich. ex Pres. 265
americanum (L.) K. Schum 265
clandestinum Hochst. ex chiov. 266
flaccidum Griseb. 266
glaucum (L.) R. Br. 266
orientale (L.C.) Rich. 266
pedicellatum Trin 266
purpureum K. Schum. 267
typhoideum L.C. Rich. 265
typhoides (Burm. f.) Stapf. & C.E. Hubb. 265
Peristylus Bl. 99
affinis (D. Don) Seid. 99
aristatus Lindl. 99
constrictus (Lindl.) Lindl. 99
fallax Lindl. 100
goodyeroides (D. Don) Lindl. 100
 var. *goodyeroides* 100
 var. *affinis* Lindl. 100
nematcaulon (Lindl.) Mitra
parishii Reichb. f. 100
prainii (Hook. f.) Kranze. 100
pseudophrys (King & Pantl.) Mitra 100
tentaculatus (Lindl.) J.J. Smith 100
Phacelurus Griseb. 267
 zey (Clarke) W.D. Clayton 267
Phaeneilema malabarica (L.) Narayanswamy & Biswas
Phaenospermum Benth. & Hook. f. 298
globosum Munro ex Oliver 298
Phaius Lour. 101
flavus (Bl.) Lindl. 101
albus Lindl. 116
blumei var. *puchra* King & Pantl. 101
maculatus Lindl. 101
mishmensis Reichb. f. 101
tancarvilleae (Banks ex L' Herit) Bl. 101
 var. *tancarvilleae* 101
 var. *pulchra* (King & Pantl.) Karth. 101
wallichii Lindl. 101
woodfordii (Hook.) Merr. 102
Phalaenopsis Bl. 102
deliciosa Reichb. f. 82
listeri Berkeley 102
lobbii (Reichb. f.) Sweet 102
mannii Reichb. f. 102
parishii var. *lobbii* Reichb. f. 102
wightii Reichb. f. 82
Phalangium nepalense Lindl. 147
Phalaris L. 298
arundinacea L. 298
hispida Thunb. 244
minor Retz. 299
zizanioides L. 276
Phleum L. 299
indicum Houtt.
alpinum L. 299
Phoenix L. 182
acaulis Roxb. ex Buch. - Ham. 182
humilis Royle ex Becc. & Hook. f. 182
lourierii Kunth 182
rupicola T. Anders. 183
sylvestris (L.) Roxb. 183
Pholidota Lindl. ex Hook. f. 102
articulata Lindl. 102
 var. *articulata*
 var. *griffithii* King & Pantl. 102
imbricata (Roxb.) Lindl. 103
protracta Hook. f. 103
recurva Lindl 103
rubra Lindl.
undulata Lindl. 103
Phragmites Adans 299
australis (Cav.) Trin ex Steud. 299
communis Trin 299
Phreatia Lindl. 103
elegans Lindl. 103
Phyllorchis helenae O. Ktze. 35
rolfei O. Ktze. 38
Phyllostachys Sieb. & Zucc. 241
assamica Gamble ex Brandis 241
bambusoides Gamble 241
Pinanga Bl. 183
gracilis (Roxb.) Bl. 183
Piptatherum P. Beauv. 299
aequiglume (Duthie ex HOok. f.) Duthie 299
Platanthera L.C. Rich. 104
bakeriana (King & Pantl.) Kranzl. 104
biermanniana (King & Pantl.) Kranzl. 104
clavigera Lindl. 104
dyeriana King & Pantl. 104
excelliana 800 104
junccea (King & Pantl.) Kranzl. 104
latilabris Lindl. 105
leptocaulon (Hook. f.) 800 105
sikkimensis (Hook. f.) Kranzl. 105

- stenantha** (Hook. f.) 500 105
Plectocomia Mart. & Bl. 183
himalayana Griff. 183
Pleione D. Don 105
 hookeriana (Lindl.) William. 105
 humilis (J.E. Smith) D. Don 106
 maculata (Lindl.) Lindl. 106
 praecox (J.E. Smith) D. Don 106
Poa L. 300
 alpina (Blytt.) Lindm. 302
 var. *sikkimensis* Stapf. 302
 annua L. 300
 atrovirens Derf. 289
 curvula Schrad. 289
 leanorae Bor. 300
 ferruginea Thunb. 289
 flexuosa Stapf. 302
 gammieana Hook. f. 300
 gangetica Roxb. 289
 himalayana Nees ex Steud. 300
 hirtiglumis Stepf. 300
 japonica Thunb. 290
 khasiana Stapf. 301
 ludens R.R. Stewart 301
 nemoralis L. 301
 var. *ligulata* Stapf. 301
 nitide-spiculata Bor 301
 nutans Retz. 290
 pagophila Bor 302
 pilosa L. 291
 polyneuron Bor 302
 poophagorum Bor 302
 pratensis var. *alpigena* Blytt.
 pseudo-pratensis Hook. f. 301
 sikkimensis (Stapf.) Bor 302
 tenella L. 291
 tibeticola Bor 302
 tremula Lam. 291
 unioloides Retz. 292
 viscosa Retz. 292
POACEAE 2 37
Podochilus Bl. 106
 culturatus Lindl. 106
 khasianus Hook. f. 107
Pogonatherum contortum Brongn. 267
 crinitum (Thunb.) Kunth 267
 paniceum (Lam.) Hack. 267
 saccharoideum P. Beauv. 267
Pogonia King & Pantl.
 falcata King & Pantl.
 gammieana Hook. f.
 hookeriana King & Pantl.
 macroglossa Hook. f.
 plicata Lindl.
- prainiana* King & Pantl.
 scottii Reichb. f.
Pollia Thunb. 171
 acisia Hassk. 171
 hasskarlii Rolla Rao 171
 subumbellata Clarke 171
Pollinia articulata 258
 ciliata Trin. 258
 cumingii Nees 255
 grata Hack. 258
 hirtifolia Hack. 255
 imberbis Nees ex Steud. 259
 mollis (Griseb.) Hack. 255
 monantha Nees ex Steud. 258
 nuda Trin 28
 quadrinervis Hack. 255
 vagans Nees ex Steud. 258
Polygonatum Nill. 158
 brevistylum Baker 158
 cathcartii Baker 158
 cirrhifolium (Wall.) Royle 158
 curvistylum Hua 159
 hookeri Baker 158
 hansuense 159
 leptopyllan D. Don 159
 norvulosum 159
 oppositifolium (Wall.) Royle 159
 punctatum Royle ex Kunth 159
 singalilense Hara 169
 verticillatum (L.) All. 180
Polypogon Desf. 303
 fugax Nees ex Steud. 303
 higagaweri Steud. 303
 litoralis 303
 var. *higagaweri* (Steud.) Hook. f. 303
Polytoca R. Br. 268
 bracteata R. Br. 268
 digitata (L.f.) Druce 268
 wallichiana (Nees ex Steud.) Benth. 268
Pomatocalpa Breda Kuhl & Hasselt 107
 armigerum (King & Pantl.) Tang &
 Wang 107
 wendlandorum (Reichb. f.) J.J. Smith
 107
Ponerorchis Reichb. f. 107
 chusua (D. Don) 107
 var. *chusua* 107
 var. *nana* (King & Pantl.) R.C. Srivast.
 108
 nana (King & Pantl.) 500 108
PONTEDERIACEAE 166
Pontederia *crassipes* Mart. 166
 vaginalis Burm. f. 166

- POOIDAE** 276
Porpax Lindl. 108
filiformis Persoon King & Pantl. 108
meirax (Par. & Reichb. f.) King & Pantl.
108
var. *meirax*
var. *elwesii* R.C. Srivast.
- POTAMOGETONACEAE** 196
Potamogeton L. 196
filiformis Persoon 196
indicus sensu Hook. f. 196
javanicus Hack. 196
nodosus Poir 196
octandrus Poir 196
pectinatus L. 46
- Pothos** L. 192
cathcartii Schott. 192
decurvata Roxb. 193
glaucus Wall. 193
heterophyllum Roxb. 192
lasia Roxb. 192
officinalis Roxb. 199
roxburghii Schott. 192
vriesianus Schott.
- Pseudechinolaena** (Hook. f.) Stapf. 268
polystachya (H.B.K.) Stapf. 268
- Pseudopogonatherum** A. Camus 268
articulata Trin 268
var. *pedicellata* Hack. ex Hook. f. 268
- bruniana** Griff 269
contortum (Brongn.) A. Camus 268
- Pseudoraphis** Griff.
brunoniana Griff.
- Pseudosorghum** A. Camus 269
fasciculare (Roxb.) A. Camus 269
polymorphum Munro 241
- Pteroceras** Hasselt. ex Hassk. 108
suaveolens (Roxb.) Holttum 108
- Pterygodium** *sulcatum* Roxb. 120
- Pycreus** P. Beauv. 231
capillaris (Koen. ex Roxb.) Nees ex Clarke 231
diaphanus (Schrad ex R. & S.) Hooker & T. Koyama 231
flavidus (Retz.) T. Koyama 231
var. *flavidus* 231
var. *erectus* (Clarke) Karthik. 231
globosus (All.) Reichb. f. 231
var. *erectus* Clarke 231
latespicatus (Boeck.) Clarke 231
polystachyos Rottb. 232
pumilus (L.) Nees ex Clarke 232
sanguinolentus (Vahl) Nees ex Clarke 232
- stramineus** Clarke 233
sulcinux (Clarke) Clarke 233
Pythonium *bulbiferum* Schott. 186
- Remusatia** Schott 192
hookerians Schott 192
pumila D. Don 193
vivipara (Roxb.) Schott
- Raphidophora** Hassk. 193
callophyllum Schott. 193
decurvata (Roxb.) Schott. 193
eximia Schott 193
glauca (Wall.) Schott. 193
grandis Schott. 194
hookeri Schott. 194
lancifolia Schott. 193
- Rhopalephora** Hassk. 172
scaberrima (Bl.) Faden 172
- Rhynchospora** Vahl 233
aurea Vahl 233
brownii R. & S. 234
corymbosa (L.) Britt. 233
glauca sensu Clarke 234
griffithii Boeck. 234
var. *levisetis* Clarke 234
var. *sikkimensis* (Clarke) Verma & Chandra 234
rugosa (Vahl) Gale 234
var. *rugosa*
var. *griffithii* (Boeck.) Verma & Chandra
ssp. *brownii* (R. & S.) T. Koyama 234
sikkimensis Clarke
- Rhychostylis** Bl. 109
retusa (L.) Bl. 109
- Risleya** King & Pantl. 109
atropurpurea King & Pantl. 109
- Ritaia** King & Pantl. 109
himalaica (Hook. f.) King & Pantl. 109
- Robiquetia** Gaud. 109
spathulata (Bl.) J.J. Smith 109
succisa (Lindl.) Scid. & Garay 110
- Roscoea** J.E. Smith 132
alpina Royle 132
var. *minor* Royle 132
auriculata K. Schum. 132
gracilis J.E. Smith 123
longifolia Baker 132
lutea Royle 124
purpurea J.E. Smith 132
spicata J.E. Smith 124
- Rottboellia** L. f. 269
cochininchinensis (Lour.) Clayton 269

- compressa* L. f. 256
exaltata L. f. 269
 var. *robusta* Hook. f. 269
striata Nees ex Steud. 260
 ssp. *khasiana* Hack. 260
zea Clarke 267
- Saccharum** L. 270
fastigatum Nees ex Steud. 254
longisetosum (Anders. ex Bengh.)
 Narayansw. ex Bor 270
narenga (Nees ex Steud.) Hack. 270
officinarum L. 270
paniceum Lam. 267
procerum Roxb. 270
rufipilum Steud. 271
sikkimensis (Hook. f.) Narayan sw. ex.
 Bor 271
spontaneum L. 271
- Sacciolepis** Nash 271
indica (L.) A. Chase 271
interrupta (Willd.) Stapf. 272
myosuroides (R. Br.) A. Camus 272
- Saccolabiopsis** J.J. Smith 110
pusilla (Lindl.) Seid. & Garay 110
- Saccolabium** acuminatum (Lindl.) Hook.
 f. 117
acutifolia Lindl. 73
affine King & Pantl. 73
ampullaceum (Roxb.) Lindl. 31
buccosum Reichb. f. 110
calceolare (J.E. Smith) Lindl. 74
dasypogon (J.E. Smith) Lindl. 74
distichum Lindl. 74
gemmae Lindl. 111
intermedium Griff. ex Lindl. 74
lancifolium King & Pantl. 117
longifolium (Lindl.) Hook. f. 25
micranthum Lindl. 111
obtusifolium (Lindl.) Hook. f. 117
ochraceum Lindl. 24
papillosum Lindl. 24
psendodistichum King & Pantl. 75
pumilio Reichb. f. 110
pusillum Lindl. 110
racemiferum Lindl. 46
rheedii Wt. 109
trichromum Reichb. f. 45
- Sarcopodium** griffithii Lindl. 34
rotundatum Lindl.
- Sagittaria** L.
guayanensis Kunth
 ssp. *lappula* (D. Don) Bogin
lappula D. Don
- Sansevieria** Mill. 160
cylindrica Boj. 160
trifasciata Hort. ex Prain 160
- Sarcanthus** appendiculatus (Lindl.) Par. 46
aspersus Reichb. f. 46
bicuspidatus (Hook. f.) J.J. Smith 46
filiformis Lindl. 46
insectifer Reichb. f.
pallidus Lindl. 46
secundus Griff. 47
subulatus (Bl.) Reichb. f. 47
succisus Lindl. 110
- Sarcochilus** bimaculatus King & Pantl. 31
crepidiformis King & Pantl. 114
hirtus (Lindl.) Benth. 112
luniferus (Reichb. f.) Benth. ex Hook. f.
 45
mannii Hook. f.
obtusus (Lindl.) Benth ex Hook. f. 91
pygmaeus King & Pantl. 113
retrospiculatus King & Pantl. 114
suaveolens (Roxb.) Hook. f. 108
- Sarcopodium** griffithii Lindl.
rotundatum Lindl. 64
- Satyrium** Swartz. 110
ciliatum Lindl. 110
nepalense D. Don 110
 var. *nepalense* 110
f. albiflorum Tuyama 110
 var. *ciliata* (Lindl.) Hook. f. 110
- repens* L. 77
- Sauromatum** Schott.
- brevipes** (Hook. f.) N.E. Br. 194
- Schizachyrium** Nees 272
brevifolium (Swartz) Nees ex Buese 272
delavayi (Hack) Bor 272
- Schizostachyum** Nees 241
capitatum (Munro) R. Majumdar 241
latifolium (Munro) R. Majumdar 241
munroi S. Kumar & P. Singh 241
polymorphum (Munro) R. Majumdar
 241
sharmae S. Kumar 241
- Schoenoplectus** (H.G.L. Reichb.) Palla 234
- Schoenoplectus goossus** (L. f.) Palla 234
juncoides (Roxb.) Palla 235
mucronatus (L.) Palla 235
supinus (L.) Palla 235
triangulatus (Roxb.) Sojak 235
- Schoenorchis** Bl. 111
gemmata (Lindl.) J.J. Smith 111
nemorum Vahl
rugosus Vahl

- Schoenoxiphium clarkeanum* Kükenth. 222
Scindapsus Schott 194
 officinalis (Roxb.) Schott 194
Scirpus L. 235
 aestivalis Retz. 219
 barbatus Rottb. 198
 bisumbellatus Forsk. f. 220
 caricis Retz. 198
 var. *sikkimensis* Clarke 198
chinensis Munro 235
chinensis Osb. 228
comosus Wall. 219
complanatus Retz. 220
corymbosus L. 230, 233
densus Wall. 198
dichotomus L. 220
erectus sensu Clarke 235
grossus L.f. 234
 var. *kysoor* (Roxb.) Clarke 234
juncoides Roxb. 235
kysoor Roxb. 234
lithospermus L.
miliaceus L. 220, 221
mucronatus L. 235
palustris L. 218
plantagineus Retz. 218
schoenoides Retz. 221
setaceus L. 222
supinus L. 235
ternatanus Reinw. ex Miq. 235
triangulatus Roxb. 235
Scleria Berg. 236
elata Thw. 236
 var. *decolorans* Clarke 236
 var. *latior* Clarke 236
hebecarpa Nees 236
 var. *pubescens* (Steud.) Clarke 236
levis Retz. 236
lithosperma (L.) Swartz. 236
parvula Steud. 236
pubescens Steud. 236
terrestris (L.) Fassett. 236
tessellata sensu clarke 236
Semiarundinaria pantlingii (Gamble) Nakai
 243
Serapias helleborine L. 44, 65
 var. *longifolia* L. 44
Setaria P. Beauv. 212
 barbata (Lam.) Kunth 272
forbesianas (Nees ex Steud.) Hook. f.
 273
glauca (L.) P. Beauv. 274
glauca auct. 266
intermedia R. & S. 273
italica (L.) P. Beauv. 273
pallida-fusca (K. Schum.) Stapf. & CE.
 Hubb. 274
palmifolia (Koen.) Stapf. 273
plicata (Lam.) T. Cooke 274
pumila (Poir) R. & S. 274
tomentosa (Roxb.) Kunth 273
verticillata (L.) P. Beauv. 274
SMILACACEAE 163
Smilacina Desf.
acuminata (Wang & Tang) Hara 153
fusca Wall. 153
oleracea (Baker) Hook. f. 153
 var. *acuminata* Wang & Tang 153
oligophylla (Baker) Hook. f. 153
pallida auct. 154
pallida Royle 154
purpurea Wall. 154
Smilax L. 163
aspera L. 163
aspericalis Wall. 163
bracteata Presl. 163
 var. *verruculosa* (Merr.) T. Koyama
 163
elegans Wall. ex Kunth 164
ferox Wall ex Kunth 164
glaucophylla Klotzsch. 163
hajrae R.C. Srivastava 164
lanceaefolia Roxb. 164
laurifolia auct. 165
longibracteata Hook. 164
macrophylla Roxb. 165
menispermoidea A. DC. 164
minutiflora A. DC.
ocreata A. DC. 165
osmastonii Wang & Tang. 165
orthoptera 165
ovalifolia Roxb. ex D. Don 165
parvifolia Wall. ex Hook. f. 164
perfoliata Lour. 165
prolifera Roxb. 165
rigida Wall. ex Kunth 164
roxburghiana Wall. ex Hook. f. 165
wallichii Kunth 165
wightii A. DC. 165
zeylanica L. 166
Smitinandia Holttum 111
micrantha (Lindl.) Holttum 111
Sorghum miliiforme (Hack.) Snowden 274
Sorghum Moench 274
SPARGANIACEAE 197
Sparganium L. 197

- fallax** (Graebner.) 197
simpex Hook. f. 197
- Spathoglottis** Bl. 111
ixioides (D. Don) Lindl. 111
- Spiranthes** L.C. Rich. 111
sinensis (Pers.) Ames 111
- Spirodela** Schleiden
polyrhiza (L.) Schleiden
- Spondiopogon** *petiolaris* Trin 258
- Sporobolus** R. Br. 303
diander (Retz.) P. Beauv. 303
fertilis (Steud.) W.D. Clayton 303
indicus auct. 303
piliferus (Trin) Kunth 303
- Stachypogon** *pauciflorus* Ktotsch 142
- Stauropsis** *undulatus* (Lindl.) Benth. ex Hook. f. 118
- Stegosia** *cochinchinensis* Lour. 269
- Stelis** *hirta* J.E. Smith 35
odoratissima J.E. Smith 36
- Stenophyllus** *barbata* (Rottb.) T. Cooke 198
- Stereochilus** Lindl. 112
bicuspidatum (Hook. f.) King & Pantl. 96
hirtus Lindl. 112
- Stipa** L. 304
concinna Hook. f. 304
consanguinea Trin & Rupr. 304
hookeri Stapf. 305
koelzii R.R. Stewart 304
mongholica Turcz. ex Trin 304
orthoraphium Steud. 304
purpurea Griseb. 304
roylei (Nees) Mez. 304
- Streptolirion** Edgew. 172
volubile Edgew. 172
- Streptopus** Michx. 161
parasimplex Hara & Ohashi 161
simplex D. Don 161
- Sunipia** Lindl. 112
andersonii King & Pantl. 112
intrmedia (King & Pantl.) P.F. Hunt 112
paleacea (Lindl.) P.F. Hunt 112
racemosa (J.E. Smith) Tang & Wang 113
scriosa Lindl. 113
- Taeniophyllum** Bl. 114
crepidiforme (King & Pantl.) King & Pantl. 114
retrospiculatum (King & Pantl.) King & Pantl. 114
- Tainia** Bl. 114
- cordata** Hook. f. 114
hookeriana King & Pantl. 114
latifolia (Lindl.) Benth. ex Hook. f. 114
minor Hook. f. 115
wrayana (Hook. f.) J.J. Smith 115
- Thamnocalamus** Munro 242
aristatus (Gamble) E.G. Camus 242
falconeri Hook. f. ex Munro 240
- Thelasis** Bl. 115
longifolia Hook. f. 115
pygmaea (Griff.) Lindl. 115
- Themeda** Forssk. 275
arundinacea (Roxb.) Ridley 275
hookeri (Griseb.) A. Camus 275
imberbis (Retz.) T. Cooke 276
quadrivalvis (L.) O. Ktze 275
triandra Forssk. 276
- Theropogon** Maxim 161
pallidus (Kunth) Maxim. 161
- Thomsonia** Wall.
nepalensis Wall.
- Thrixspermum** Lour. 113
luniferum Reichb. f. 45
pygmaeum (King & Pantl.) Holtt. 113
teres Reichb. f. 108
nemorale J.R. & G. Forst.
- Thryocephalon** *namorale* J.R. & G. Forst. 227
- Thunia** Reichb. f. 116
alba (Lindl.) Reichb. f. 116
- Thyrsia** *zea* (Clarke) Stapf. 267
- Thysanolaena** Nees 305
agrostis Nees 305
latifolia (Roxb. ex Hornem.) Honda 305
maxima (Roxb.) O. Ketz. 305
sikkimensis Gdger. 305
- Tipularia** Nutt. 116
josephii Reichb. f. ex. Lindl. 116
- Tofieldia** Huds. 161
himlaica Baker 161
- Tovaria** *finitima* W.W. Smith
fusca Baker 153
oleracea Baker 153
pallida Royle 154
purpurea (Wall.) Baker 154
- Trachycarpus** Wendl. 184
martianus (Mart.) Wendl. 184
- Tradescantia** *cordifolia* Griff. 172
cristata L. 169
malabarica L. 171
vaga Lour. 169
- Tribachia** *reptans* Lindl. 37
- Trichoasma** *suavis* Lindl. 67

- Trichotosia** Bl. 116
dasyphylla (Par. & Reichb. f.) Kranzl. 116
pulvinata (Lindl.) Kranzl. 116
- Tricyrtis** Wall. 162
maculata (D. Don) J.F. Mc Bride 162
pilosa Wall. 162
- Triglochin** L. 195
maritima L. 195
palustre L. 195
- Trikeria** Bor. 305
hookeri (Stapf.) Bor 305
- Trilepis royleana** Nees 225
- trillidium** Kunth 162
- tsechonoskii** Maxim 162
govanianum D. Don 162
- Trillium** L.
obovatum Pursh 162
- Tripogon** Roem. & Schult. 305
filiformis Nees ex Steud. 305
trifidus Munro 306
- Trisetum** Pres. 306
flavescens (L.) P. Beauv. 306
spicatum (L.) Richt 306
trifidus Munro ex Stapf.
scitulum Bor 306
subspicatum (L.) P. Beauv. 306
virescens Nees ex Steud. 295
- Triticum** L. 306
aestivum L. 306
semicostatum Nees ex Steud. 288
vulgare Vill. 306
- Tropidia** Lindl. 113
angulosa (Lindl.) Bl. 113
bellii Blatt. & Mc C. 113
curculigooides Lindl. 113
- Tupistra** L. 162
aurantiaca (Baker) Wall. ex Hook. f. 146
clarkii Hook. f. 162
natans Wall. ex Lindl. 163
- Tylostylis** Bl. 117
discolor Hook. f. 117
- Typhonium** Schott 195
brevipes Hook. f. 194
diversifolium Wall. ex Schott. 195
trilobatum (L.) Schott. 195
- Uncifera** Lindl. 117
acuminata Lindl. 117
lancefolia (King & Pantl.) Schltr. 117
obtusifolia Lindl. 117
- Uncinia nepalensis** Nees 224
- Urachne acutigluma** Steud. 294
Uvularia barviflora 149
- Vanda** W. Jones ex R. Br. 117
alpina Lindl. 117
cathcartii Lindl. 30
cristata Lindl. 118
longifolia Lindl. 25
multiflora Lindl.
parviflora Lindl. 118
pumila Hook. f. 118
teres (Roxb.) Lindl. 98
testacea (Lindl.) Reichb. f. 118
trichorhiza Hook. 88
undulata Lindl. 118
- Vandopsis** Pfitz. 118
undulata (Lindl.) J.J. Smith 118
- Vetiveria** Bory de St. Vincent 276
zizanioides (L.) Nash 276
- Vilfa pilifera** Trin. 303
- Wallichia** Roxb. 184
densiflora Mart. 184
disticha T. Anders. 184
oblongifolia Griff. 184
- Xanthosoma brasiliense** (Desf.) Engl.
- Xyridaceae** 166
Xyris L. 166
indica L. 166
pauciflora Willd. 167
schoenoides Mart. 167
- Yoania** Maxim. 118
prainii King & Pantl. 118
- Ypsilandra** Franch. 163
yunnanensis W.W. Smith & Jeffreys 163
var. *himalaica* Hara 163
- Yucca** Dill. ex L. 140
aloifolia L. 140
- Yushania** K.H. Keng 242
maling (Camble) R. Majumdar 242
microphylla (Griff.) R. Majumdar 242
pantlingii (Gamble) R. Majumdar 242
racemosa (Munro) R. Majumdar 243
- Zea** L. 276
mays L. 276
- Zephyranthes** Herb. 138
candida Herb. 138
carinata Herb. 138
- Zerna himalaica** Stapf. 282
- Zeuxine** Lindl. 119
affinis (Lindl.) Benth. ex Hook. f. 119

- flava** (Lindl.) Trimen. 119
franchetiana King & Pantl. 119
goodyeroides Lindl. 119
nervosa (Wall. ex Lindl.) Timen 119
pulchra King & Pantl. 120
pumila (Hook. f.) King & Pantl. 92
reflexa King & Pantl. 120
strateumatica (Lindl.) Schlechtr. 120
sulcata (Roxb.) Lindl. 120
ZINGIBERACEAE 120
Zingiber Boehm. 132
- capitatum** Roxb. 132
var. **capitatum** 132
chrysanthum Rosc. 133
clarkei King & Benth. 133
elatum Roxb. 133
nigrum Gaertn. 121
officinale Rosc. 133
purpureum Rosc. 134
rubens Roxb. 134
zerumbet (L.) Roxb. ex J.E. Smith 13
Zizania terrestris L. 236